


Concept Paper

# Following Excitation/Inhibition Ratio Homeostasis from Synapse to EEG in Monogenetic Neurodevelopmental Disorders

Lisa Geertjens <sup>1,2</sup>, Torben W. van Voorst <sup>3</sup> , Arianne Bouman <sup>4</sup> , Maaïke A. van Boven <sup>3</sup>, Tjitske Kleefstra <sup>4</sup>, Matthijs Verhage <sup>3,5</sup>, Klaus Linkenkaer-Hansen <sup>6</sup>, Nael Nadif Kasri <sup>4,†</sup>, L. Niels Cornelisse <sup>3,5,†</sup> and Hilgo Bruining <sup>1,2,7,\*</sup>

- <sup>1</sup> Child and Adolescent Psychiatry and Psychosocial Care, Emma Children's Hospital, Amsterdam UMC, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, The Netherlands; l.m.g.geertjens@amsterdamumc.nl
- <sup>2</sup> N=You Neurodevelopmental Precision Center, Amsterdam Neuroscience, Amsterdam Reproduction and Development, Amsterdam UMC, Meibergdreef 5, 1105 AZ Amsterdam, The Netherlands
- <sup>3</sup> Department of Functional Genomics Center for Neurogenomics and Cognitive Research (CNCR), VU University Amsterdam and Amsterdam UMC-Location VUmc, de Boelelaan 1085, 1081 HV Amsterdam, The Netherlands; t.w.van.voorst@vu.nl (T.W.v.V.); m.a.van.boven@vu.nl (M.A.v.B.); m.verhage@vu.nl (M.V.); l.n.cornelisse@vu.nl (L.N.C.)
- <sup>4</sup> Department of Human Genetics, Radboud University Medical Center, 6500 HB Nijmegen, The Netherlands; arianne.bouman@radboudumc.nl (A.B.); tjitske.kleefstra@radboudumc.nl (T.K.); n.nadif@donders.ru.nl (N.N.K.)
- <sup>5</sup> Department of Human Genetics, Center for Neurogenomics and Cognitive Research (CNCR), Amsterdam UMC-Location VUmc, de Boelelaan 1085, 1081 HV Amsterdam, The Netherlands
- <sup>6</sup> Department of Integrative Neurophysiology, Center for Neurogenomics and Cognitive Research (CNCR), Amsterdam Neuroscience, VU University Amsterdam, 1081 HV Amsterdam, The Netherlands; k.linkenkaerhansen@vu.nl
- <sup>7</sup> Levvel, Center for Child and Adolescent Psychiatry, Meibergdreef 5, 1105 AZ Amsterdam, The Netherlands
- \* Correspondence: h.bruining@amsterdamumc.nl
- † These authors contributed equally to this work.



**Citation:** Geertjens, L.; van Voorst, T.W.; Bouman, A.; van Boven, M.A.; Kleefstra, T.; Verhage, M.; Linkenkaer-Hansen, K.; Nadif Kasri, N.; Cornelisse, L.N.; Bruining, H. Following Excitation/Inhibition Ratio Homeostasis from Synapse to EEG in Monogenetic Neurodevelopmental Disorders.

*Genes* **2022**, *13*, 390. <https://doi.org/10.3390/genes13020390>

Academic Editors: Christine M. Freitag, Jacob A. S. Vorstman and Antonio Persico

Received: 28 December 2021

Accepted: 16 February 2022

Published: 21 February 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Pharmacological options for neurodevelopmental disorders are limited to symptom suppressing agents that do not target underlying pathophysiological mechanisms. Studies on specific genetic disorders causing neurodevelopmental disorders have elucidated pathophysiological mechanisms to develop more rational treatments. Here, we present our concerted multi-level strategy 'BRAINMODEL', focusing on excitation/inhibition ratio homeostasis across different levels of neuroscientific interrogation. The aim is to develop personalized treatment strategies by linking iPSC-based models and novel EEG measurements to patient report outcome measures in individual patients. We focus our strategy on chromatin- and SNAREopathies as examples of severe genetic neurodevelopmental disorders with an unmet need for rational interventions.

**Keywords:** neurodevelopmental disorders; iPSC-based models; EEG; SNAREopathies; chromatinopathies


## 1. Introduction

Neurodevelopmental disorders (NDDs) are highly heterogeneous in etiology and manifestation, and cause tremendous suffering for patients and caregivers. Current treatments are limited to generic symptom suppressing medications that do not take heterogeneity into account. There is a need for mechanism-based therapeutic options to remedy the life-long suffering of patients and caregivers often associated with NDDs.

The identification of risk genes for NDDs provides new starting points for mechanism-based therapies. For example, in recent work, 102 risk genes for autism spectrum disorder (ASD) have been identified [1]. The discovery of NDD-associated de novo variants in genes

with roles in synaptic plasticity has provided an entry to start developing rational interventions. Indeed, NDDs caused by variations in single genes, so called monogenetic NDDs (mNDDs), seem to converge on a disturbed balance between excitatory and inhibitory inputs (E/I) in neuronal networks in the brain [2–8] that occur early (1st/2nd trimester), or in early postnatal stages [1]. Although the concept is rather generic and applied in many contexts, it is well established that cortical networks require a finely tuned coordination of excitatory and inhibitory inputs for normal information processing [9], and that changes in both directions (increasing or decreasing E/I ratio) may compromise processing and lead to NDD clinical symptoms. The E/I-balance concept is further supported by NDD mouse model studies that show E/I ratio disturbances [10] and EEG abnormalities in NDD patients that suggest E/I ratio imbalances [11–13]. Finally, we have recently reported initial successes with off-label medication targeting E/I-regulation [14–16]. Thus, influencing E/I ratios is regarded as a promising target for pharmaceutical interventions, but is complicated by the multifaceted heterogeneity of underlying mechanisms and thus requires personalized treatment strategies [17].

Here, we first outline the heterogeneous nature and consequences of E/I ratio disturbances observed in NDD model research, which emphasizes the need for personalized treatment development strategies. We put forward that induced pluripotent stem-cell (iPSC)-based models provide new opportunities for translatability of E/I ratios to network activity homeostasis as proposed by BRAINMODEL. This project is conducted by a publicly funded Dutch consortium of neuroscientists and clinicians, and aims to develop personalized E/I targeting treatments through the linking iPSC-based models, EEG data, and clinical assessments in patients with two forms of genetic NDDs, chromatin-, and SNAREopathies (Figure 1).



**Figure 1.** BRAINMODEL's multi-level strategy.

## 2. Molecular and Physiological Heterogeneity of E/I Ratio Homeostasis in NDDs

Alterations in E/I ratio homeostasis in NDDs can result from aberrations in several processes, including synapse development, synaptic transmission, and neuronal excitability [18]. For example, synaptic E/I ratio changes have been described in human iPSC-derived neurons from Rett-Syndrome patients carrying *MECP2* loss-of-function mutations, resulting in decreased excitatory synaptic activity with no change in inhibitory activity [19]. Likewise, iPSC-derived neurons from individuals with Phelan–McDermid syndrome (PMDs) and autism showed selective defects in excitatory, but not inhibitory, synaptic transmission [20], and disruption of the autism-associated gene *SYNGAP1* increased excitatory synapse numbers in developing human neurons [21]. Alternatively, neurological phenotypes associated with E/I ratios changes have been observed as the broadening of action potentials in neurons derived from individuals with Timothy syndrome [22], while the genetic deletion of the Angelman Syndrome-associated gene *UBE3A* was shown to increase the excitability of induced human neurons [23]. Similarly, hu-

man neurons KO for the Fragile X Syndrome-associated gene *FMR1* exhibited increased intrinsic excitability, with no discernible synaptic phenotype [24]. In addition, altered synaptic E/I ratios and neuronal excitability phenotypes often co-occur. For instance, NDD-associated variants of the synaptic protein CASK appear to reduce the size of inhibitory presynaptic compartments, while simultaneously reducing spiking activity [25]. Likewise, the conditional deletion of the PMDS-associated gene *SHANK3* did not only produce hyperexcitability through modulation of intrinsic membrane properties, but also produced extensive synaptic impairments [26].

### 3. The Potential and Pitfalls of E/I Ratio Measurements in iPSC Models

The above findings underline the potential of models consisting of neurons and neuronal networks derived from *patient-own* tissues using iPSC-technology. These may bridge the gap between in vitro models and in vivo manifestations [27], and for instance, link different levels of E/I ratio homeostasis in response to treatment. The generation of glutamatergic and GABAergic neurons from patient-derived iPSCs can be achieved either using dual-SMAD inhibition [28] or through the ectopic expression of transcription factors [29–31], of which the latter is better suited to robust high-throughput assays in terms of both scalability, and cellular and maturational homogeneity [32–34]. The use of single-neuron (“autapse”) cultures [35,36] generated from induced glutamatergic and GABAergic neurons (iNeurons) grown in isolation on microdot arrays, allows for robust and standardized analysis of cell-autonomous synaptic functioning and intrinsic excitability without the interference of homeostatic mechanisms. Conversely, co-culturing patient-derived glutamatergic and GABAergic iNeurons on multi-electrode arrays (MEAs) allows for recording of network activity and analysis of network E/I ratio [37]. Moreover, by recording the development of neuronal network behavior over time, the interactions between the primary disfunctions in synaptic activity or excitability and maturation checkpoints can be studied, such as the development of mature intracellular chloride levels [31].

It is important to note that methodologies to develop human-neurons from iPSCs are still actively being developed and optimized, and it has become clear that neurons generated in vitro differ substantially from those found in the human brain [38]. Thus, one should be aware that while these cells resemble human neurons, they are not necessarily identical to those found in the human brain. Nevertheless, it is clear that induced neurons have a clear neuronal morphology, are electrophysiologically active, express markers for specific neuronal lineages also found in the human brain, and form interconnected networks able to integrate into the developing brain in mice in vivo [39,40], illustrating its validity as a model for (developing) human neurons. In addition, in the developing brain, neurons receive many inputs from many different cell-types, which are under strict spatiotemporal control. This cannot be accurately modeled in in vitro systems. Thus, readouts obtained from these systems should always be interpreted in its context as a simplified model.


Furthermore, these methodologies come paired with moral and ethical issues. As this novel strategy has yet to prove its efficacy as a tool for the selection of therapeutic interventions, patients and other stakeholders might be reluctant to participate in this study. Another concern is that cultured human neurons are often portrayed as ‘miniature brains’, suggesting that there is a possibility that consciousness could be generated in cultured neurons. This brings forward concerns regarding the moral status of said cultured neuronal networks. However, its highly improbable that cultured neurons in the methodologies currently employed could be conscious, as the number of neurons and the complexity of the networks in these systems are low, even compared to cultured neuronal organoids, which resemble the human cortex to a greater extent and for which this concern is more relevant [41,42].

### 4. Focus on Chromatinopathies and SNAREopathies

In the BRAINMODEL project, we will initially focus our multi-level phenotyping on two classes of mNDDs. We thereby aim to collect observations from multiple individuals

with variants in the same gene to identify cellular and network hallmarks and targets for these disorders, as we have previously done for Kleefstra-syndrome and *STXBP1*-encephalopathy, part of the so-called chromatinopathies and SNAREopathies, respectively [43,44]. Indeed, using CRISPR/Cas9-technology, the genotype of the iPSCs can be altered, inducing disease-associated variants in healthy cells or repairing the variant in patient-derived cells [45], demonstrating causal relationships between detected phenotypes and patient genotype. After the identification of cellular and network deficits, therapeutic strategies to correct these aberrations can be selected and tested in vitro as performed by Marchetto et al. and Yahata et al. for Rett syndrome and Alzheimer's disease, respectively [19,46]. In BRAINMODEL, we will expand our expertise on iPSC characterization of chromatinopathies and SNAREopathies and focus on treatment development for these two classes of mNDDs.


Chromatinopathies are found to be major contributors to NDDs [1,47]. Chromatin remodeling determines whether genes are available for transcription and is crucial in active regulation of gene expression. We focus on four different monogenic disorders caused by a pathological mutation or deletion in genes (*EHMT1*, *KMT2C*, *KMT2D* and *SETD1A*) coding for enzymes which carry out chromatin remodeling (e.g., methylation) (Figure 2). Although mNDDs caused by mutations in these genes have the same neurobiological etiology (altered chromatin remodeling), there is a high variability in clinical presentation. Apart from intellectual disability (ID) and/or developmental delay (DD) presenting in almost 100% of cases, core symptoms are childhood hypotonia, psychiatric disorders (including autism-spectrum disorder (ASD), attention-deficit disorder (ADHD), and anxiety), epilepsy, and sleep disorders. In addition, facial dysmorphisms are present, and anomalies are found in several organ systems [48–51]. As for the E/I ratio homeostasis, previous studies showed that the Loss of function (LoF) of *EHMT1* results in delayed GABAergic maturation, reduced inhibition, and hence increased E/I ratio [52–54].



**Figure 2.** Transcription regulation is controlled by histone modifications. Schematic representation of the deposition of histone methylation, performed by histone methyltransferases and histone demethylases on histone 3 (H3), which have been linked to neurodevelopmental disorders.

SNAREopathies are another group of pathobiological well-defined mNDDs. These disorders, caused by mutations that disturb SNARE function, are a subset of the previously defined synaptopathies. The neuronal SNARE complex (soluble NSF attachment protein receptor complex) is an important molecular machine driving synaptic vesicle exocytosis and secretion of neuropeptides and neuromodulators from dense core vesicles [55]. We focus on four of these genes that are associated with NDDs (*STXBP1*, *SYT1*, *SNAP25*, *RIMS1*) (Figure 3). Although the pathogenic starting point of these disorders is well defined, clinical

phenotype and disease severity is very diverse. Moreover, high clinical variety is found in the same amino acid changes between different individuals [55]. Most common clinical aspects found in SNAREopathies are ID and/or DD, seizures, ASD, and neurological motor problems. Even though almost all cases present with ID, the mechanisms through which mutations in SNARE genes lead to neurodevelopmental impairments remain unexplained. Additional genetic and/or environmental factors might contribute substantially to disease presentation and should be considered when studying disease mechanisms [55]. Based on mouse models, mutations in SNAREopathy genes also create a disturbed E/I ratio setpoints [55]. However, it is unknown to which extent the different components in the E/I microcircuits in the brain are susceptible to gene mutations [55].



**Figure 3.** Schematic Representation of the eight SNAREopathy genes with their orientation relative to the synaptic vesicle and the plasma membrane and their interaction.

## 5. Connecting the Dots

There is need for caution in interpreting neuronal and network phenotypes, as an apparent E/I phenotype might be the result of homeostatic compensation mechanisms [6] rather than a cell-autonomous NDD phenotype [56]. This is illustrated by the finding that the pharmacological induction of hyperexcitability was sufficient to phenocopy *SHANK*<sup>-/-</sup>-associated synaptic defects in wild type neurons [26], similar to findings in mouse models, where an altered E/I ratio was found to be a compensatory mechanism to stabilize the circuit [10]. Indeed, the applicability for therapeutic screening in iPSC based models is limited, as the model does not represent a full organism. Pharmacodynamics and -kinetics are different, for example, due to the incapacity to model the blood brain barrier. Thus, whether a potential therapeutic intervention can reach the target cells in vivo cannot be determined in these models. Furthermore, off-target effects at other areas of the body cannot be studied. iPSC-based models do, however, provide the opportunity to test novel compounds in vitro in advance of clinical trials. This enables the identification of more therapeutic options, either based upon existing or new compounds, where the former has the advantage of knowledge on pharmacodynamics and -kinetics.

To complement the multi-level strategy on a neurophysiological level, resting state electroencephalography (rsEEG) recordings can be analyzed in the same patients. We have put forward that the concept of critical brain dynamics is a steppingstone to derive E/I ratios from neuronal oscillations measured with conventional EEG [57–60] (represented in Figure 4B) [58,61]. Our computational modeling [60] (Figure 4A), as well as pharmacological challenges [61], have indicated that the so-called ‘critical’ regime between low and high activity requires balance between excitation and inhibition. Therefore, the basis of the E/I method is the statistical character of activity in this critical state where long-range temporal correlations (LRTC) [60] weaken when network E/I is out of balance [61,62] (Figure 4C,D). Therefore, we could use LRTCs to estimate E/I ratios leading to a functional E/I measure (fE/I) (Figure 4E–G). We validated this fE/I method at rest (rsEEG) and after GABAergic treatment. In children with ASD, we corroborated that both increased and decreased E/I





problematic behavior or affective dysregulation and how disturbed E/I ratio homeostasis may be translated into clinical readout measures [63]. This resulted in the PROM for the repeated and reliable measurement of patient-relevant consequences of sensory reactivity alterations [63] that we developed by following the FDA steps for (parent proxy) PROM for clinical trials [66,67]. According to this protocol, we initiated focus groups and interviews with caregivers to elicit the most impactful and most relevant symptoms and then sought to measure these PROs with large item banks of the Patient-Reported Outcomes Measurement Information System® (PROMIS) [68], initiated by the “NIH Roadmap Initiative”, based on Item Response Theory (IRT) with the possibility to use Computerized Adaptive Testing (CAT) [69].

## 6. Conclusions

Previous research has provided crucial starting points to understand pathophysiological mechanisms in NDDs to develop therapeutic options. iPSC-based models for mNDDs have unprecedented promise to bridge the gap between well-established animal and cellular models towards human treatment development. In BRAINMODEL, we will employ a multi-level strategy in which iPSC based-models, neurophysiological parameters, and PROMs are combined within the framework of chromatin- and SNAREopathies in order to develop personalized mechanism-based therapeutic strategies targeting E/I ratio homeostasis.

**Author Contributions:** Conceptualization, H.B. and L.N.C.; writing—original draft preparation, L.G., M.A.v.B., T.W.v.V. and A.B.; writing—review and editing, H.B., L.N.C., M.V., K.L.-H., N.N.K. and T.K.; All authors have read and agreed to the published version of the manuscript.

**Funding:** BRAINMODEL ZonMW PSIDER program 10250022110003. BECAUSE ZonMW TOP 91216064. NewTDEC Netherlands Organization for Scientific Research (NWO) Dutch National Research Agenda, NWA-ORC Call (NWA.1160.18.200). Aspasia Grant of the Dutch Research Council (015.014.036). Netherlands Organization for Health Research and Development (91718310).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** K.L.-H. is a shareholder of NBT. Analytics BV, which provides EEG-analysis services for clinical trials. H.B. and K.L.-H. are shareholders of Aspect Neuroprofiles BV, which develops physiology-informed prognostic measures for neurodevelopmental disorders. K.L.-H. has filed the patent claim (PCT/NL2019/050167) “Method of determining brain activity”; with priority date 16 March 2018.

## References

1. Satterstrom, F.K.; Kosmicki, J.A.; Wang, J.; Breen, M.S.; De Rubeis, S.; An, J.Y.; Peng, M.; Collins, R.; Grove, J.; Klei, L.; et al. Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiology of Autism. *Cell* **2020**, *180*, 568–584.e23. [[CrossRef](#)] [[PubMed](#)]
2. Bozzi, Y.; Provenzano, G.; Casarosa, S. Neurobiological bases of autism-epilepsy comorbidity: A focus on excitation/inhibition imbalance. *Eur. J. Neurosci.* **2018**, *47*, 534–548. [[CrossRef](#)]
3. Foss-Feig, J.H.; Adkinson, B.D.; Ji, J.L.; Yang, G.; Srihari, V.H.; McPartland, J.C.; Krystal, J.H.; Murray, J.D.; Anticevic, A. Searching for Cross-Diagnostic Convergence: Neural Mechanisms Governing Excitation and Inhibition Balance in Schizophrenia and Autism Spectrum Disorders. *Biol. Psychiatry* **2017**, *81*, 848–861. [[CrossRef](#)] [[PubMed](#)]
4. Jeste, S.S.; Frohlich, J.; Loo, S.K. Electrophysiological biomarkers of diagnosis and outcome in neurodevelopmental disorders. *Curr. Opin. Neurol.* **2015**, *28*, 110–116. [[CrossRef](#)] [[PubMed](#)]
5. Naaijen, J.; Bralten, J.; Poelmans, G.; Glennon, J.C.; Franke, B.; Buitelaar, J.K. Glutamatergic and GABAergic gene sets in attention-deficit/hyperactivity disorder: Association to overlapping traits in ADHD and autism. *Transl. Psychiatry* **2017**, *7*, e999. [[CrossRef](#)] [[PubMed](#)]
6. Nelson, S.B.; Valakh, V. Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. *Neuron* **2015**, *87*, 684–698. [[CrossRef](#)] [[PubMed](#)]

7. Selten, M.; van Bokhoven, H.; Kasri, N.N. Inhibitory control of the excitatory/inhibitory balance in psychiatric disorders. *F1000Research* **2018**, *7*, 23. [[CrossRef](#)]
8. Rubenstein, J.L.; Merzenich, M.M. Model of autism: Increased ratio of excitation/inhibition in key neural systems. *Genes Brain Behav.* **2003**, *2*, 255–267. [[CrossRef](#)]
9. Isaacson, J.S.; Scanziani, M. How inhibition shapes cortical activity. *Neuron* **2011**, *72*, 231–243. [[CrossRef](#)]
10. Antoine, M.W.; Langberg, T.; Schnepel, P.; Feldman, D.E. Increased Excitation-Inhibition Ratio Stabilizes Synapse and Circuit Excitability in Four Autism Mouse Models. *Neuron* **2019**, *101*, 648–661.e4. [[CrossRef](#)]
11. Boutros, N. Epileptiform discharges in psychiatric patients: A controversy in need of resurrection. *Clin. EEG Neurosci.* **2009**, *40*, 239–244. [[CrossRef](#)]
12. Bruining, H.; Hardstone, R.; Juarez-Martinez, E.L.; Sprengers, J.; Avramiea, A.E.; Simpraga, S.; Houtman, S.J.; Poil, S.S.; Dallares, E.; Palva, S.; et al. Measurement of excitation-inhibition ratio in autism spectrum disorder using critical brain dynamics. *Sci. Rep.* **2020**, *10*, 9195. [[CrossRef](#)]
13. Spence, S.J.; Schneider, M.T. The role of epilepsy and epileptiform EEGs in autism spectrum disorders. *Pediatr. Res.* **2009**, *65*, 599–606. [[CrossRef](#)]
14. Bruining, H.; Passtoors, L.; Goriounova, N.; Jansen, F.; Hakvoort, B.; De Jonge, M.; Poil, S.S. Paradoxical Benzodiazepine Response: A Rationale for Bumetanide in Neurodevelopmental Disorders? *Pediatrics* **2015**, *136*, e539–e543. [[CrossRef](#)]
15. Sprengers, J.J.; Van Andel, D.M.; Zuithoff, N.P.; Keijzer-Veen, M.G.; Schulp, A.J.; Scheepers, F.E.; Lilien, M.R.; Oranje, B.; Bruining, H. Bumetanide for Core Symptoms of Autism Spectrum Disorder (BAMBI): A Single Center, Double-Blinded, Participant-Randomized, Placebo-Controlled, Phase Two, Superiority Trial. *J. Am. Acad. Child Adolesc. Psychiatry* **2020**, *60*, 865–876. [[CrossRef](#)]
16. Van Andel, D.M.; Sprengers, J.J.; Oranje, B.; Scheepers, F.E.; Jansen, F.E.; Bruining, H. Effects of bumetanide on neurodevelopmental impairments in patients with tuberous sclerosis complex: An open-label pilot study. *Mol. Autism* **2020**, *11*, 30. [[CrossRef](#)]
17. Charman, T.; Loth, E.; Tillmann, J.; Crawley, D.; Wooldridge, C.; Goyard, D.; Ahmad, J.; Auyeung, B.; Ambrosino, S.; Banaschewski, T.; et al. The EU-AIMS Longitudinal European Autism Project (LEAP): Clinical characterisation. *Mol. Autism* **2017**, *8*, 27. [[CrossRef](#)]
18. Trobiani, L.; Meringolo, M.; Diamanti, T.; Bourne, Y.; Marchot, P.; Martella, G.; Dini, L.; Pisani, A.; De Jaco, A.; Bonsi, P. The neuroligins and the synaptic pathway in Autism Spectrum Disorder. *Neurosci. Biobehav. Rev.* **2020**, *119*, 37–51. [[CrossRef](#)]
19. Marchetto, M.C.; Carromeu, C.; Acab, A.; Yu, D.; Yeo, G.W.; Mu, Y.; Chen, G.; Gage, F.H.; Muotri, A.R. A Model for Neural Development and Treatment of Rett Syndrome Using Human Induced Pluripotent Stem Cells. *Cell* **2010**, *143*, 527–539. [[CrossRef](#)]
20. Shcheglovitov, A.; Shcheglovitova, O.; Yazawa, M.; Portmann, T.; Shu, R.; Sebastiano, V.; Krawisz, A.; Froehlich, W.; Bernstein, J.A.; Hallmayer, J.F.; et al. SHANK3 and IGF1 restore synaptic deficits in neurons from 22q13 deletion syndrome patients. *Nature* **2013**, *503*, 267–271. [[CrossRef](#)]
21. Llamosas, N.; Arora, V.; Vij, R.; Kilinc, M.; Bijoch, L.; Rojas, C.; Reich, A.; Sridharan, B.; Willems, E.; Piper, D.R.; et al. SYNGAP1 Controls the Maturation of Dendrites, Synaptic Function, and Network Activity in Developing Human Neurons. *J. Neurosci.* **2020**, *40*, 7980–7994. [[CrossRef](#)]
22. Paşca, S.P.; Portmann, T.; Voineagu, I.; Yazawa, M.; Shcheglovitov, A.; Paşca, A.M.; Cord, B.; Palmer, T.D.; Chikahisa, S.; Nishino, S.; et al. Using iPSC-derived neurons to uncover cellular phenotypes associated with Timothy syndrome. *Nat. Med.* **2011**, *17*, 1657–1662. [[CrossRef](#)]
23. Sun, A.X.; Yuan, Q.; Fukuda, M.; Yu, W.; Yan, H.; Lim, G.G.Y.; Nai, M.H.; D’agostino, G.A.; Tran, H.D.; Itahana, Y.; et al. Potassium channel dysfunction in human neuronal models of Angelman syndrome. *Science* **2019**, *366*, 1486–1492. [[CrossRef](#)]
24. Susco, S.G.; Arias-García, M.A.; López-Huerta, V.G.; Beccard, A.; Bara, A.M.; Moffitt, J.; Korn, J.; Fu, Z.; Barrett, L.E. FMR1 loss in a human stem cell model reveals early changes to intrinsic membrane excitability. *Dev. Biol.* **2020**, *468*, 93–100. [[CrossRef](#)]
25. Becker, M.; Mastropasqua, F.; Reising, J.P.; Maier, S.; Ho, M.L.; Rabkina, I.; Li, D.; Neufeld, J.; Ballenberger, L.; Myers, L.; et al. Presynaptic dysfunction in CASK-related neurodevelopmental disorders. *Transl. Psychiatry* **2020**, *10*, 312. [[CrossRef](#)]
26. Yi, F.; Danko, T.; Botelho, S.C.; Patzke, C.; Pak, C.; Wernig, M.; Südhof, T.C. Autism-associated SHANK3 haploinsufficiency causes Ih channelopathy in human neurons. *Science* **2016**, *352*, aaf2669. [[CrossRef](#)]
27. Dolmetsch, R.; Geschwind, D.H. The human brain in a dish: The promise of iPSC-derived neurons. *Cell* **2011**, *145*, 831–834. [[CrossRef](#)]
28. Chambers, S.M.; Fasano, C.A.; Papapetrou, E.P.; Tomishima, M.; Sadelain, M.; Studer, L. Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat. Biotechnol.* **2009**, *27*, 275–280. [[CrossRef](#)]
29. Zhang, Y.; Pak, C.; Han, Y.; Ahlenius, H.; Zhang, Z.; Chanda, S.; Marro, S.; Patzke, C.; Acuna, C.; Covy, J.; et al. Rapid single-step induction of functional neurons from human pluripotent stem cells. *Neuron* **2013**, *78*, 785–798. [[CrossRef](#)]
30. Yang, N.; Chanda, S.; Marro, S.; Ng, Y.H.; Janas, J.A.; Haag, D.; Ang, C.E.; Tang, Y.; Flores, Q.; Mall, M.; et al. Generation of pure GABAergic neurons by transcription factor programming. *Nat. Methods* **2017**, *14*, 621–628. [[CrossRef](#)]
31. Mossink, B.; Van Rhijn, J.R.; Wang, S.; Linda, K.; Vitale, M.R.; Zöller, J.E.; van Hugte, E.J.; Bak, J.; Verboven, A.H.; Selten, M.; et al. Cadherin-13 is a critical regulator of GABAergic modulation in human stem-cell-derived neuronal networks. *Mol. Psychiatry* **2021**, 1–18. [[CrossRef](#)]



32. Busskamp, V.; Lewis, N.E.; Guye, P.; Ng, A.H.; Shipman, S.L.; Byrne, S.M.; Sanjana, N.E.; Murn, J.; Li, Y.; Li, S.; et al. Rapid neurogenesis through transcriptional activation in human stem cells. *Mol. Syst. Biol.* **2014**, *10*, 760. [[CrossRef](#)] [[PubMed](#)]
33. Wang, C.; Ward, M.E.; Chen, R.; Liu, K.; Tracy, T.E.; Chen, X.; Xie, M.; Sohn, P.D.; Ludwig, C.; Meyer-Franke, A.; et al. Scalable Production of iPSC-Derived Human Neurons to Identify Tau-Lowering Compounds by High-Content Screening. *Stem Cell Rep.* **2017**, *9*, 1221–1233. [[CrossRef](#)] [[PubMed](#)]
34. Deneault, E.; Faheem, M.; White, S.H.; Rodrigues, D.C.; Sun, S.; Wei, W.; Piekna, A.; Thompson, T.; Howe, J.L.; Chalil, L.; et al. CNTN5-/+or EHMT2-/+human iPSC-derived neurons from individuals with autism develop hyperactive neuronal networks. *eLife* **2019**, *8*, e40092. [[CrossRef](#)]
35. Meijer, M.; Rehbach, K.; Brunner, J.W.; Classen, J.A.; Lammertse, H.C.; van Linge, L.A.; Schut, D.; Krutenko, T.; Hebisch, M.; Cornelisse, L.N.; et al. A Single-Cell Model for Synaptic Transmission and Plasticity in Human iPSC-Derived Neurons. *Cell Rep.* **2019**, *27*, 2199–2211.e6. [[CrossRef](#)]
36. Rhee, H.J.; Shaib, A.H.; Rehbach, K.; Lee, C.; Seif, P.; Thomas, C.; Gideons, E.; Guenther, A.; Krutenko, T.; Hebisch, M.; et al. An Autaptic Culture System for Standardized Analyses of iPSC-Derived Human Neurons. *Cell Rep.* **2019**, *27*, 2212–2228.e7. [[CrossRef](#)] [[PubMed](#)]
37. Mossink, B.; Verboven, A.H.; van Hugte, E.J.; Gunnewiek, T.M.K.; Parodi, G.; Linda, K.; Schoenmaker, C.; Kleefstra, T.; Kozicz, T.; van Bokhoven, H.; et al. Human neuronal networks on micro-electrode arrays are a highly robust tool to study disease-specific genotype-phenotype correlations in vitro. *Stem Cell Rep.* **2021**, *16*, 2182–2196. [[CrossRef](#)]
38. Lin, H.C.; He, Z.; Ebert, S.; Schörnig, M.; Santel, M.; Nikolova, M.T.; Weigert, A.; Hevers, W.; Kasri, N.N.; Taverna, E. NGN2 induces diverse neuron types from human pluripotency. *Stem Cell Rep.* **2021**, *16*, 2118–2127. [[CrossRef](#)] [[PubMed](#)]
39. D’Alessio, R.; Koukoulis, F.; Blanchard, S.; Catteau, J.; Raïs, C.; Lemonnier, T.; Féraud, O.; Bennaceur-Griscelli, A.; Groszer, M.; Maskos, U. Long-term development of human iPSC-derived pyramidal neurons quantified after transplantation into the neonatal mouse cortex. *Dev. Biol.* **2020**, *461*, 86–95. [[CrossRef](#)]
40. Vitrac, A.; Pons, S.; Balkota, M.; Lemièrre, N.; Raïs, C.; Bourgeois, J.P.; Maskos, U.; Bourgeron, T.; Cloëz-Tayarani, I. A chimeric mouse model to study human iPSC-derived neurons: The case of a truncating SHANK3 mutation. *Sci. Rep.* **2020**, *10*, 13315. [[CrossRef](#)]
41. Lavazza, A. Human cerebral organoids and consciousness: A double-edged sword. *Monash Bioeth. Rev.* **2020**, *38*, 105–128. [[CrossRef](#)] [[PubMed](#)]
42. Lavazza, A. Potential ethical problems with human cerebral organoids: Consciousness and moral status of future brains in a dish. *Brain Res.* **2021**, *1750*, 147146. [[CrossRef](#)] [[PubMed](#)]
43. Lammertse, H.C.A.; van Berkel, A.A.; Iacomino, M.; Toonen, R.F.; Striano, P.; Gambardella, A.; Verhage, M.; Zara, F. Homozygous STXBP1 variant causes encephalopathy and gain-of-function in synaptic transmission. *Brain* **2019**, *143*, 441–451. [[CrossRef](#)]
44. Frega, M.; Linda, K.; Keller, J.M.; Gümüş-Akay, G.; Mossink, B.; van Rhijn, J.-R.; Negwer, M.; Gunnewiek, T.K.; Foreman, K.; Kompier, N.; et al. Neuronal network dysfunction in a model for Kleefstra syndrome mediated by enhanced NMDAR signaling. *Nat. Commun.* **2019**, *10*, 4928. [[CrossRef](#)] [[PubMed](#)]
45. Adli, M. The CRISPR tool kit for genome editing and beyond. *Nat. Commun.* **2018**, *9*, 1911. [[CrossRef](#)]
46. Yahata, N.; Asai, M.; Kitaoka, S.; Takahashi, K.; Asaka, I.; Hioki, H.; Kaneko, T.; Maruyama, K.; Saido, T.C.; Nakahata, T.; et al. Anti-A $\beta$  drug screening platform using human iPSC cell-derived neurons for the treatment of Alzheimer’s disease. *PLoS ONE* **2011**, *6*, e25788. [[CrossRef](#)] [[PubMed](#)]
47. Ciptasari, U.; van Bokhoven, H. The phenomenal epigenome in neurodevelopmental disorders. *Hum. Mol. Genet.* **2020**, *29*, R42–R50. [[CrossRef](#)] [[PubMed](#)]
48. Kleefstra, T.; de Leeuw, N. Kleefstra Syndrome. In *GeneReviews*<sup>®</sup>; Adam, M.P., Ed.; University of Washington: Seattle, WA, USA, 1993.
49. Gallagher, D.; Voronova, A.; Zander, M.A.; Cancino, G.I.; Bramall, A.; Krause, M.P.; Abad, C.; Tekin, M.; Neilsen, P.; Callen, D.; et al. Ankrd11 is a chromatin regulator involved in autism that is essential for neural development. *Dev. Cell* **2015**, *32*, 31–42. [[CrossRef](#)]
50. Kummeling, J.; Stremmelaar, D.E.; Raun, N.; Reijnders, M.R.F.; Willemsen, M.H.; Ruiterkamp-Versteeg, M.; Schepens, M.; Man, C.C.O.; Gilissen, C.; Cho, M.T.; et al. Characterization of SETD1A haploinsufficiency in humans and Drosophila defines a novel neurodevelopmental syndrome. *Mol. Psychiatry* **2021**, *26*, 2013–2024. [[CrossRef](#)]
51. Adam, M.P.; Hudgins, L.; Hannibal, M. Kabuki syndrome. In *GeneReviews*<sup>®</sup>; Adam, M.P., Ed.; University of Washington: Seattle, WA, USA, 1993.
52. Martens, M.B.; Frega, M.; Classen, J.; Epping, L.; Bijvank, E.; Benevento, M.; van Bokhoven, J.; Tiesinga, P.; Schubert, D.; Kasri, N.N. Euchromatin histone methyltransferase 1 regulates cortical neuronal network development. *Sci. Rep.* **2016**, *6*, 35756. [[CrossRef](#)]
53. Negwer, M.; Piera, K.; Heslen, R.; Lütje, L.; Aarts, L.; Schubert, D.; Kasri, N.N. EHMT1 regulates Parvalbumin-positive interneuron development and GABAergic input in sensory cortical areas. *Brain Struct. Funct.* **2020**, *225*, 2701–2716. [[CrossRef](#)]
54. Frega, M.; Selten, M.; Mossink, B.; Keller, J.M.; Linda, K.; Moerschen, R.; Qu, J.; Koerner, P.; Jansen, S.; Oudakker, A.; et al. Distinct Pathogenic Genes Causing Intellectual Disability and Autism Exhibit a Common Neuronal Network Hyperactivity Phenotype. *Cell Rep.* **2020**, *30*, 173–186.e6. [[CrossRef](#)] [[PubMed](#)]
55. Verhage, M.; Sorensen, J.B. SNAREopathies: Diversity in Mechanisms and Symptoms. *Neuron* **2020**, *107*, 22–37. [[CrossRef](#)] [[PubMed](#)]

56. Pintacuda, G.; Martín, J.M.; Eggan, K.C. Mind the translational gap: Using iPSC cell models to bridge from genetic discoveries to perturbed pathways and therapeutic targets. *Mol. Autism* **2021**, *12*, 10. [[CrossRef](#)] [[PubMed](#)]
57. Hardstone, R.; Poil, S.-S.; Schiavone, G.; Jansen, R.; Nikulin, V.V.; Mansvelder, H.D.; Linkenkaer-Hansen, K. Detrended fluctuation analysis: A scale-free view on neuronal oscillations. *Front. Physiol.* **2012**, *3*, 450. [[CrossRef](#)]
58. Linkenkaer-Hansen, K.; Nikouline, V.V.; Palva, J.M.; Ilmoniemi, R. Long-range temporal correlations and scaling behavior in human brain oscillations. *J. Neurosci.* **2001**, *21*, 1370–1377. [[CrossRef](#)]
59. Poil, S.-S.; de Haan, W.; van der Flier, W.M.; Mansvelder, H.D.; Scheltens, P.; Linkenkaer-Hansen, K. Integrative EEG biomarkers predict progression to Alzheimer's disease at the MCI stage. *Front. Aging Neurosci.* **2013**, *5*, 58. [[CrossRef](#)]
60. Poil, S.-S.; Hardstone, R.; Mansvelder, H.; Linkenkaer-Hansen, K. Critical-state dynamics of avalanches and oscillations jointly emerge from balanced excitation/inhibition in neuronal networks. *J. Neurosci. Off. J. Soc. Neurosci.* **2012**, *32*, 9817–9823. [[CrossRef](#)]
61. Beggs, J.M.; Plenz, D. Neuronal avalanches in neocortical circuits. *J. Neurosci.* **2003**, *23*, 11167–11177. [[CrossRef](#)]
62. Poil, S.S.; van Ooyen, A.; Linkenkaer-Hansen, K. Avalanche dynamics of human brain oscillations: Relation to critical branching processes and temporal correlations. *Hum. Brain Mapp.* **2008**, *29*, 770–777. [[CrossRef](#)]
63. van Andel, D.M.; van Stel, H.F.; Scheepers, F.E.; Oostrom, K.J.; Haverman, L.; Bruining, H. The sensory-reactivity PROM set: Identification of a parent reported outcome measure set for autism spectrum disorder. *J. Patient Rep. Outcomes* **2021**, *5*, 123. [[CrossRef](#)] [[PubMed](#)]
64. Schauder, K.B.; Bennetto, L. Toward an Interdisciplinary Understanding of Sensory Dysfunction in Autism Spectrum Disorder: An Integration of the Neural and Symptom Literatures. *Front. Neurosci.* **2016**, *10*, 268. [[CrossRef](#)] [[PubMed](#)]
65. Wigham, S.; Rodgers, J.; South, M.; McConachie, H.; Freeston, M. The interplay between sensory processing abnormalities, intolerance of uncertainty, anxiety and restricted and repetitive behaviours in autism spectrum disorder. *J. Autism Dev. Disord.* **2015**, *45*, 943–952. [[CrossRef](#)] [[PubMed](#)]
66. Food and Drug Administration. Guidance for Industry: Patient-Reported Outcome Measures: Use in Medical Product Development to Support Labelling Claims. Available online: [www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM193282.pdf](http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM193282.pdf) (accessed on 5 December 2020).
67. Streiner, D.L.; Norman, G.R. *Health Measure Scales: A Practical Guide to Their Development and Use*, 4th ed.; Oxford University Press: New York, NY, USA, 2008.
68. Cella, D.; Yount, S.; Rothrock, N.; Gershon, R.; Cook, K.; Reeve, B.; Ader, D.; Fries, J.F.; Bruce, B.; Rose, M. The Patient-Reported Outcomes Measurement Information System (PROMIS): Progress of an NIH Roadmap cooperative group during its first two years. *Med. Care* **2007**, *45*, S3–S11. [[CrossRef](#)] [[PubMed](#)]
69. HealthMeasures. Computer Adaptive Tests (CATs). Available online: <https://www.healthmeasures.net/resource-center/measurement-science/computer-adaptive-tests-cats> (accessed on 26 February 2021).