



Review

# The Unforeseen Non-Coding RNAs in Head and Neck Cancer

Alexandra Iulia Irimie <sup>1</sup>, Alina-Andreea Zimta <sup>2</sup>, Cristina Ciocan <sup>2</sup> , Nikolay Mehterov <sup>3,4</sup> ,  
Diana Dudea <sup>1</sup>, Cornelia Braicu <sup>5</sup> and Ioana Berindan-Neagoe <sup>2,5,6,\*</sup>

- <sup>1</sup> Department of Prosthetic Dentistry and Dental Materials, Division Dental Propaedeutic, Aesthetic, “Iuliu-Hatieganu” University of Medicine and Pharmacy, Cluj-Napoca, 23 Marinescu Street, 40015 Cluj-Napoca, Romania; irimie.alexandra@umfcluj.ro (A.I.I.); ddudea@umfcluj.ro (D.D.)
- <sup>2</sup> MEDFUTURE—Research Center for Advanced Medicine, University of Medicine and Pharmacy Iuliu-Hatieganu, 23 Marinescu Street, 40015 Cluj-Napoca, Romania; andreea.zimta@umfcluj.ro (A-A.Z.); cristina.ciocan@umfcluj.ro (C.C.)
- <sup>3</sup> Department of Medical Biology, Medical University Plovdiv, BulVasilAprilov 15-A, Plovdiv 4002, Bulgaria; ni\_ki82@abv.bg
- <sup>4</sup> Technological Center for Emergency Medicine, BulVasilAprilov 15-A, Plovdiv 4002, Bulgaria
- <sup>5</sup> Research Center for Functional Genomics and Translational Medicine, “Iuliu-Hatieganu” University of Medicine and Pharmacy, 23 Marinescu Street, 40015 Cluj-Napoca, Romania; braicucornelia@yahoo.com
- <sup>6</sup> Department of Functional Genomics and Experimental Pathology, The Oncology Institute “Prof. Dr. Ion Chiricuta”, Republicii 34 Street, 40015 Cluj-Napoca, Romania
- \* Correspondence: ioana.neagoe@umfcluj.ro; Tel.: +40-264-598-885

Received: 31 January 2018; Accepted: 22 February 2018; Published: 1 March 2018

**Abstract:** Previously ignored non-coding RNAs (ncRNAs) have become the subject of many studies. However, there is an imbalance in the amount of consideration that ncRNAs are receiving. Some transcripts such as microRNAs (miRNAs) or small interfering RNAs (siRNAs) have gained much attention, but it is necessary to investigate other “pieces of the RNA puzzle”. These can offer a more complete view over normal and pathological cell behavior. The other ncRNA species are less studied, either due to their recent discovery, such as stable intronic sequence RNA (sisRNA), YRNA, miRNA-offset RNAs (moRNA), telomerase RNA component (TERC), natural antisense transcript (NAT), transcribed ultraconserved regions (T-UCR), and pseudogene transcript, or because they are still largely seen as non-coding transcripts with no relevance to pathogenesis. Moreover, some are still considered housekeeping RNAs, for instance small nucleolar RNAs (snoRNAs) and TERC. Our review summarizes the biogenesis, mechanism of action and potential role of less known ncRNAs in head and neck cancer, with a particular focus on the installment and progress for this particular cancer type.

**Keywords:** head and neck cancer; rare non-coding RNAs; biological mechanisms

## 1. Introduction

Head and neck cancer encompasses a group of malignancies developing in the areas of the head outside the central nervous system [1]. Esophageal cancer is associated with head and neck cancer due to its anatomical proximity and common risk factors. The major risk factors are increased alcohol consumption, tobacco smoking, human papilloma virus (HPV) infection and betel chewing [2–5].

Squamous cell carcinoma is related to *NOTCH1*, *PIK3CA* and *PTEN* mutations, while esophageal cancer is related to high rates of *KRAS* mutations and *HER2* amplification [6]. Accompanying these mutations, there are alterations in the transcriptomics landscape of the cells [7].

The present study is an incursion on rare RNA. Their biogenesis, mechanism of action and biological effects are analyzed, in relation to their implication in head and neck cancer [8–12].

The importance of microRNAs (miRNAs) and small interfering RNAs (siRNAs) cannot be ignored [9,13]. It is however important to study other molecular players of carcinogenesis that would offer a more complete view [13,14]. Each class of these lesser known transcripts is presented separately, with a particular focus on the relationship with head and neck cancer.

## 2. The Unforeseen RNAs


A few decades ago, all molecular biology studies were based on the central dogma that stated there is an exchange of information between DNA and RNA, and that this unequivocal information encodes the protein sequence of amino acids [10,15,16]. With the discovery of miRNAs in *Caenorhabditis elegans* and small interfering RNAs (siRNAs) in plants [17–19], the central dogma was challenged. This was the first time that the issue of non-coding RNAs (ncRNAs) with regulatory functions entered the mainstream of molecular research [17–19]. RNAs are nowadays classified as protein-coding RNAs and ncRNAs. Nevertheless, some RNAs were found in the grey area, as having both regulatory and protein-coding abilities [20].

miRNAs are the most studied transcripts. Thus far, there are numerous preclinical studies that are trying to formulate better diagnostic, prognostic and therapeutic options based on miRNAs [10,14,21–26]. The miRNAs that are up-regulated in cancer cells are called oncomiRs. The down-regulated miRNAs are named tumor suppressor miRNAs [27]. In addition, there is an abundance of data regarding the role of miRNAs in the diagnosis and prognosis of head and neck cancer [7,28].

siRNAs are also a well-known ncRNAs class. These transcripts are used in preclinical study as therapeutic strategy or to study the role of a specific gene, mainly due to their ability to target a single gene [13,29–31]. siRNAs can also become the basis for future therapeutic strategies, targeting cancer-promoting gene transcripts in various cancers [29,31–35], such as head and neck cancer [36,37].


miRNA-offset RNAs (moRNAs) are located in the nucleus [38], where they interact with messenger RNA (mRNA) [39,40]. Transcribed ultraconserved regions (T-UCR) have two mechanisms of action; changing the methylation of CpG island in the DNA [41,42] and miRNA inhibition through direct interaction [43–45]. The stable intronic sequence RNA (sisRNA) mechanism of action interacts post-transcriptionally with natural antisense transcript (NAT) in *Drosophila melanogaster* [46]. sisRNAs are also thought to regulate the expression of their own gene [47]. Some sisRNAs interact with the small nucleolar RNAs (snoRNAs)-specific protein FOX2, hence they are also involved in alternative splicing [48]. snoRNAs are involved primarily in ribosomal RNA (rRNA) nucleotide changes, but also in the splicing machinery [49]. YRNAs decide the fate of misfolded RNAs [50] or bind to the chromatin during the early stages of DNA replication [51]. Small YRNAs (sYRNAs), even if derived from the YRNAs, seem to have an unknown molecular mechanism of action. They sustain, however, the toll-like receptor-mediated cell apoptosis [52]. Enhancer RNAs (eRNAs) interact with transcription factors and the promoter region of their target gene in a specific spatial-temporal manner that results in the transcription stimulation [53]. Telomerase RNA component (TERC) interacts with telomerase and helps in telomere length maintenance [54]. In plants, it was proven that tRNA (transfer RNA)-derived fragments (tRFs) interact with the argonaute 1 (AGO1) and Dicer-like proteins and repress the transportable elements expression [55], but, in animal cells, their mechanism of action is still unknown [56]. NAT can act as source of endogenous siRNA, as post-transcriptional repressors for the mRNA of the antisense strand [57] and as modulator of heterochromatin [58]. The pseudogenes are transcribed in approximately 20% of cases. Some pseudogene transcripts were even found to be translated into peptides [52,59,60]. Whether these peptides have a functional role inside the cell remains to be studied [60,61]. On the other hand, pseudogene transcripts can also have regulatory roles inside the cell. They can be sources of endogenous siRNA and miRNA [62], can sponge miRNAs [63] or regulate gene expression through epigenetic modifications [62]. Moreover, recently, based on bioinformatics tools, it was found that small RNAs transcripts interact with long non-coding RNAs (lncRNAs) [64,65]. The main characteristics of these transcripts are summarized in Table 1. Their biogenesis is illustrated in Figure 1 and their mechanism of action in Figure 2.

It is worth noting that some ncRNAs are located in the intronic region of the mRNA and that they are involved in splicing. This correlates with the observation that some synonymous mutations, located in the intronic region of the DNA, can cause translation repression or intronic maintenance in the transcript. This “intronic retention” was observed in several tumor suppressors transcripts in human cancer [66].



**Figure 1.** A schematic representation of the biogenesis and the main characteristics of some rare RNA species. The bidirectional arrow represents the length of the transcript, the red arrow stands for transcription and the blue arrow stands for RNA processing. (A) The enhancer RNA (eRNA) is transcribed from the enhancer DNA (eDNA) region. The transcription can be unidirectional; in this case,

eRNA is polyadenylated, has a length of more than 4 kb and is named 1D-eRNA. If the eRNA is transcribed bidirectionally, it is not polyadenylated, has a length of less than 2 kb and is called 2D-eRNA. (B) The stable intronic RNA (sisRNA) is transcribed as primary messenger RNA (mRNA) transcript. After splicing, the intronic sequence remains stable thus forming the sisRNA. (C) The YRNA is transcribed from special regions of the DNA, called YDNA. The YRNA can remain in the nucleus or it can be cleaved into smaller fragments associated with Rho proteins. The smaller fragments are called small YRNA (sYRNA) and are sometimes exported into the cytoplasm. (D) From the promoter region of the RNA polymerase II (Pol II), the telomerase RNA component (TERC) is transcribed. TERC helps with telomerase activity. (E) From the antisense strand of protein-coding genes, the natural antisense transcript (NAT) is synthesized. NAT bears many similarities to the mRNA. (F) the small nucleolar RNA (snoRNA) can be originated from the intronic regions of protein-coding genes or from the promoter region of RNA Pol II. (G) From transporter genes, transfer RNA (tRNA) is formed; this transcript can be further fragmented and it forms the tRNA-derived fragments (tRFs). (H) The endogens are former protein-coding genes that have accumulated major mutations throughout evolution and become non-functional. The pseudogenes can be transcribed and the pseudogenes transcript bears many similarities to the mRNA. (I) The primary form of miRNA is processed by Argonaute (AGO) proteins. After processing, some part of the primary transcript remains in the nucleus, becomes stable and is very similar to miRNAs. These transcripts are named miRNA-offset RNA (moRNA). (J) Some regions of DNA, also considered non-functional, have high similarity across different species. These regions are called ultraconserved regions and can be transcribed. The transcribed ultraconserved region (T-UCR) is also similar in some aspects to mRNA.



**Figure 2.** The lesser-known non-coding RNAs (ncRNAs) analyzed in this review have different mechanisms of action. NAT and pseudogene transcripts function as miRNA sponge, esiRNA and miRNA sources and can possess coding capacity. moRNA, T-UCR and sisRNA regulate gene expression through post-translational interaction with different coding or non-coding RNA species. In the alternative-splicing process, in addition to snoRNA, sisRNA might also play a role. YRNA is implicated in DNA replication and in deciding the fate of misfolded RNA. T-UCR and eRNA control gene expression before transcription: T-UCR by modulating the methylation pattern of CpG island and eRNA by chromatin structural changes and interaction with the transcription factors. TERC, through association with telomerase, maintains the length of telomeres, thus enabling replicative immortality. sYRNA were shown to be implicated in cell apoptosis, but the exact mechanism remains unknown. In addition, tRFs are involved in cell proliferation through an unknown mechanism.

**Table 1.** The main features of less known non-coding RNAs (ncRNAs).

ncRNA Type	Length (nt)	DNA Region	Mechanism of Action	Reference
eRNA	800–2000	enhancer DNA	histone methylation	[67–69]
sisRNA	intronic length	protein-coding genes	alternative splicing, transcription stimulation, antibody isotype class switch	[70,71]
YRNA	84–112	yDNA	misfolded RNA degradation, DNA replication	[50,72,73]
sYRNA	~24 to ~34	yDNA	interaction with TLR	[74]
TERC/hTR	451	Pol II promotor region	telomere maintenance	[54,75]
NAT	depending of the coding gene length	antisense strand of protein-coding genes	mRNA inhibition	[76]
snoRNA	60–200 C/D snoRNAs, 120–250 for H/ACA snoRNAs	introns, promoter region of Pol II	rRNA processing, splicing	[77]
tRF	20	tRNA coding genes	Unknown	[78]
Pseudogenestranscripts	gene length	Pseudogenes	translation repression, miRNA sponge, multiple miRNAs, siRNA origin	[62,79]
moRNAs	20	miRNA-coding genes	Unknown	[80]
T-UCR	>200	Ultra-conserved regions of the DNA	miRNA sponging	[81]

eRNA: enhancer RNAs; nt: nucleotides; sisRNA: stable intronic sequence RNAs; sYRNA: small YRNA; TLR: Toll-like receptors; TERC: telomerase RNA component; hTR: human telomerase RNA; Pol II: polymerase II; NAT: natural antisense transcript; mRNA: messenger RNA; snoRNA: small nucleolar RNAs; H/ACA: specific H and ACA template boxes; rRNA: ribosomal RNA; tRF: tRNA-derived fragments; tRNA: transfer RNA; miRNA: micro RNA; siRNA: small interfering RNA; moRNAs: miRNA-offset RNAs; T-UCR: transcribed ultraconserved regions.

### 3. Enhancer RNAs

eRNAs were initially described in neuronal cells by Kim et al. in 2010, by using genome-wide sequencing approaches [82]. eRNAs are a type of ncRNAs of 50 bp–2 kb length that reside in the nucleus and are transcribed from the enhancer DNA. The enhancer DNA represents various areas of the DNA with the main characteristic of controlling gene expression. The enhancers interact especially with Lineage Determining Transcription Factors (LDTFs). LDTFs determine a cell type and act by modifying the chromatin structure through histone methylation. A chromatin loop is formed that allows the interaction between the enhancer, the transcription factor and the promoter region of the targeted gene. eRNAs are produced to facilitate loop formation. eRNAs arise as unidirectional (1D-eRNAs) or bidirectional transcripts (2D-eRNAs) [83]. Generally, 1D-eRNAs are long (over 4 kb) and polyadenylated, while 2D-eRNAs are relatively short (<2 kb) and are not polyadenylated [84–87]. eRNAs play crucial roles in pathogenesis [82]. The basal cell carcinoma is closely linked to the *ACTRT1* gene transcript ARP-T1 [88]. The Cancer Genome Atlas (TCGA) data analysis present molecular data, including those related to eRNAs, for different types of cancer. It was demonstrated that the metastatic niche preparation is reflected by the differentially methylated enhancer regions, followed by stimulated transcription of oncogenes and oncomiRs [89]. Enhancer RNAs can be *cis*-regulators and act on the same chromatin fiber or *trans*-regulators and induce the transcription of a gene found on another chromosome [87]. An interesting finding was the fact that p53 protein was bound mostly in the enhancer regions of the DNA. As follows, p53 modulates the expression of eRNA, in human and mouse primary fibroblast [90]. The p53 binding regions and the dysregulation of eRNAs in head and neck neoplasia may constitute a fertile ground for further studies [31,91].

### 4. Stable Intronic Sequence RNAs

SisRNAs are a class of ncRNAs originated from the spliced introns that are not degraded by the nucleases [92]. Previous studies have postulated their existence, but a study from 2012 was the one that has offered the best description yet [92]. sisRNAs are transcribed from the protein-coding genes as primary mRNA transcripts. After splicing, part of the remaining intronic RNA remains stable in the nucleus for days [92], which is why these RNA species were given the name of stable intronic sequence RNA. After the lariat is formed during splicing, the intronic sequence can remain in the circular form as circular sisRNAs (also known under the name of circular intronic RNA) or the 2′–5′ phosphodiester bond can be cleaved by the lariat debranching enzyme, and the linear sisRNA is generated [47]. In H9 and HeLa cells, it was discovered that the sisRNA *ci-ankrd52* can bind to the Pol II and stimulate the transcription of their own gene of origin [70]. A linear sisRNA directs the activation induced cytidine deaminase (AID) enzyme in the DNA of B cells that helps the immunoglobulin class switch [71]. A linear sisRNA regulates the alternative splicing process in pluripotent cells by interacting with the FOX2 protein [48]. Epstein-Barr virus (EBV) presence was correlated among other cancers, to oral squamous cell carcinoma [93]. A RNA-Seq analysis in EBV transcriptome reveal the presence of a sisRNA, *ebv-sisRNA-1* [94].

### 5. YRNAs

YRNAs were sequenced and characterized in 1980 [95]. YRNAs are versatile transcripts involved in small RNA quality control, replication mechanism and are related with cellular processes that sustains cell proliferation. They are also synthesized in response to cellular stress [96]. In addition, YRNAs are transcribed from a special part of the DNA, called YDNA. They have a stem loop structure and a length of approximately 84–112 nt. As such, there are four types of yRNAs in humans, hY1, hY3, hY4 and hY5, which have one thing in common, i.e. they are bound to the Ro60 and La proteins [97,98]. These ncRNAs are encoded by the *Y* genes located in close proximity of one another and the polymerase III (Pol III) is responsible for their transcription [72] in the form of a stem-loop secondary structure [99].



YRNAs are recognized as a component of soluble ribonucleoproteins (RNPS). These are related with many pathological conditions [100]. The Ro60 protein bound to the YRNAs controls the fate of the misfolded RNAs. The misfolded RNAs are refolded or degraded [50]. YRNAs are also a part of the initial phases of DNA replication by interacting with the chromatin and some of the replication proteins, such as Cdc6, Cdt1, DUE-B and ORC [72].

These small RNAs were, for a long period of time, categorized as miRNAs, but now they are considered as a subtype of yRNAs [101,102]. sYRNAs are particularly present in apoptotic cells. sYRNAs were initially discovered in tumor tissue. These RNAs are not processed by Dicer and do not interact with the Ago proteins [98]. The sYRNAs have a size range of 25–29 nt [103]. Their biological functions are still largely unknown. In primary cell culture of macrophage from C57BL/6J mice and THP.1 monocytes, RNY1 or RNY3 was related with the apoptosis activation through the cleavage of caspase-3 and NFκB signaling pathway [74]. Several sYRNAs were found to be down-regulated in the sera of head and neck cancer patients, compared to healthy donors, 30.2% vs. 44.2%. Using next generation sequencing (NGSseq) approaches, a specific yRNA was identified along with miRNA and tRNA signature related to head and neck squamous cell carcinoma (HNSCC), revealing a complex regulation of oncogenic processed. The most relevant altered yRNAs transcripts in head and neck cancer are summarized in Table 2 [104]. In addition, sYRN an apoptotic regulator of apoptosis is proposed as diagnostic marker [105]. s-YRNAs can be detected in tandem with other short transcripts length transcripts leading to the identification of novel signature with clinical application [105].

## 6. The Human Telomerase RNA Component or Human Telomerase RNA

The telomerase enzyme comprises two crucial constituents that operate together. The first one represents a reverse transcriptase (TERT) that has a catalytic unit associated and the other one is the TERC. TERC is a RNA template from which the missing part of the DNA is synthesized [106]. TERC was first discovered in eukaryotic cells. The human TERC (hTR) is overexpressed in the germline and in the cancerous cells [107].

hTR contains two hairpin structures separated by an H box and a single stranded ACA box. hTR is encoded by a sequence located in the promoter region of Pol II gene. Following the production of a primary transcript that has 1451 nt, the cap-binding complex (CBC) links with the ARS2 protein and directs the hTR towards processing or degradation [54]. The nuclear exosome targeting (NEXT) complex is formed of several proteins which recognize alterations in the human telomerase primary form, such as 3' extension or oligoadenylation. The NEXT complex mediates the hTR degradation by the nuclear RNA exosome complex [54]. The nuclear RNA exosome complex is a multi-protein complex that is involved in the quality control of several transcripts [108]. The final hTR transcript has 451 nts [54,75]. It was proven that it is possible for the hTR to be exported into the cytoplasm. The exported hTR was either degraded by the XRN1 and DCP2 enzymes, or multiple hTR formed complexes called cytoplasmic TERC (cyTERC) [109].

The hTR is upregulated in oral cancer tissue from patients and was associated with the malignancy installment. With the help of reverse transcription polymerase chain reaction (RT-PCR), an over 6.9-fold increase in the expression of hTR in oral squamous cell carcinoma was proven, when compared to the normal tissue. It seems that the expression of this RNA increases with the malignant transformation of cells, since it had low levels in normal epithelium and mild dysplasia, higher in moderate and severe dysplasia and the highest in the invasive oral carcinoma [110]. The Fragile X-related protein 1 (FXR1) is upregulated in HPV-negative tongue squamous cell carcinoma cell lines UMSCC74A and UMSCC74B, in which it was found to play an essential role in cell cycle progression and senescence evasion. One way in which FXR1 can cause this pathological shift is through the interaction and stabilization of hTR. The induced overexpression of hTR in these cell lines caused an amplification of immortality [111,112].

## 7. Natural Antisense Transcripts

NATs were discovered a few decades ago [113]. Some of them function as protein-coding RNAs and others as ncRNAs. They are transcribed from the antisense strand of a protein-coding gene located at the same DNA locus (cis-NAT) or from a distant site of the DNA (trans-NAT) [114]. NAT has a full complementarity with the protein-coding strand. It can stimulate or repress the transcription of a gene by modulating the heterochromatin interaction [58] or by functioning as a source of endogenous small interfering RNAs (esiRNAs) [57].

WRAP53 (WD40-encoding RNA antisense to p53) is transcribed from the antisense strand of p53 gene. It can interact with the p53 mRNA and repress its translation. Through real-time quantitative reverse transcription PCR (qRT-PCR) and Western blotting, the WRAP53 NAT was detected to have an elevated level in the esophageal carcinoma tissue samples and in the cell lines EC109, EC9706, KYSE150, and KYSE180. Its expression increases with disease progression and it was associated with the lymph node metastasis [76]. These natural antisense transcripts to p53 play a significant role in the development and progression of oral squamous cell carcinoma (OSCC), representing an important therapeutic target or biomarker [76]. It was demonstrated that ATM it is an upstream regulator of these transcript, being implicated not only in DNA damage response, but in the regulation of other functions as well [115].

## 8. The Small Nucleolar RNAs

snoRNAs are a type of linear transcripts with protein-guiding function. Depending on their sequence characteristics, the snoRNAs are divided into C/D snoRNA, H/ACA snoRNA, and U snoRNA [116,117]. The C/D snoRNA and the H/ACA snoRNA are transcribed from protein-coding genes, as primary mRNAs. After splicing, the intronic lariat is cleaved into a linear form by the Dbr1p protein. Then the 3' and the 5' end are shortened by the local endonucleases, until the mature C/D or H/ACA snoRNA is generated. The U3, U8 and U13 snoRNAs are transcribed from the promoter regions of RNA Pol II [116].

The mature forms of snoRNAs have the sequence length of 60–300 nt [118]. Based on their structure, the snoRNAs are classified into C/D box snoRNAs and H/ACA box snoRNAs. The C/D box RNAs have a double C box (RUGAUGA, R = A/G) and a double D box (CUGA). They are arranged in the C-D-C-D order and bind specific proteins, such as Snu13p, Nop56p, Nop58p and fibrillarin [119,120]. A small subclass of C/D snoRNAs has only the C or the D box and it has been reported to be subjected to mutation accumulation across species [120]. The H/ACA snoRNAs contain the H sequence (ANANNA, N = A/T/C/G) and the ACA sequence and two stem regions. These H/ACA boxes mediate the association with Gar1p, Cbf5p, Nhp2p, and Nop10p proteins [118,119]. Generally, snoRNAs are involved especially in rRNA processing. The C/D snoRNAs cause 2'-O-methylation of rRNA. The H/ACA snoRNAs convert rRNA uridine into pseudouridine [49,119]. These ncRNAs, such as HBII-52, are also components of the splicing machinery [49].


Apart from these functions there are several snoRNAs whose functions remain unknown. They are called orphan snoRNAs. Orphan SNORD32A (U32A), SNORD33 (U33) and SNORD35A (U35A) accumulate in the cytosol under lipotoxic and oxidative stress, hence it was theorized that snoRNAs can be exported in the cytoplasm and fulfill other roles [78].

By analyzing the TCGA RNA sequencing (RNASeq) data, it was discovered that snoRNA SNORD35B was upregulated in some head and neck cancer and its expression was positively associated with a lower survival rate among the patients. The overexpression of this snoRNAs was confirmed by qRT-PCR in the cell carcinoma cell lines UMSCC-10B, UMSCC-22B, HN-1, HN-12, and HN-30 [121]. In another subtype of rare oral cancer named ameloblastoma, several snoRNAs were found to be dysregulated. LINC340, SNORD116-25, SNORA11, SNORA21, SNORA47 and SNORA65 were proposed as biomarkers [122].

The microarray comparison made between OSCC tissue and keratinized gums, proved that 14 snoRNAs were down-regulated (displayed in Figure 3) and one snoRNA was up-regulated in



OSCC. An accentuated down-regulation of mgh18S-121 and ACA17 can be the cause of a worse prognostic [123].



**Figure 3.** A short list of the most studied down-regulated and up-regulated rare ncRNAs in head and neck cancer discovered to this date.

### 9. tRNA Derived Small RNAs

tRFs are transcripts with regulatory role, originated from tRNA. They are 20 nt in length and are divided into nine categories, depending on their biogenesis and mechanism of action. There are specific RNase which cleaves the primary tRNA transcript. At the tRNA 5' end, cleavage is catalyzed by the ubiquitous endoribonuclease RNase P, and tRNA 3' end cleavage is specific for RNase P [124,125].

The cleaved fragments, that have poly-U repeats at the 3' end, are categorized as tRF-1. The Dicer enzyme may further process the mature tRNA1 by a cut in the T loop and generate the tRF-3. The tRF-3 has a CCA sequence at its 3' end as a remnant of the original tRNA acceptor sequence. The mature tRNA can also be cleaved by the angiogenin enzyme and give rise to the tRF-5 [56,126].

The tRFs biological function is still unidentified, but until now, studies indicate that it may have a profound role in cancer. The tRF-1001 is responsible for increased cell proliferation [56]. They are also dysregulated in breast cancer and B cell lymphoma [127]. To the best of our knowledge, no study was done regarding the evaluation of these RNAs in head and neck cancer.

### 10. Pseudogene Transcripts

Pseudogenes are paralogous to coding genes and are the result of duplication associated with mutation. Because they are lacking the promoter and regulatory regions of the coding genes, for a long time, pseudogenes were viewed only as vestigial sequences without any function [128]. However, during the latest studies, it was proven that the pseudogenes are transcribed and that their codified transcripts are a new type of RNAs which can either regulate other RNA or are translated into proteins. Pseudogene RNA can cause epigenetic modifications of the genome by interacting with the chromatin folding around histones. They can be processed into miRNAs or can sponge other miRNAs.

Pseudogene transcripts are a source of esiRNAs that targets their paralogous genes. There are pseudogenes regulated by the same promoter as the protein-coding gene. These can be transcribed and translated into a chimerical protein [62,79]. The pseudogene transcripts ability to be translated

was proved through proteomic analysis from human tissue [52], however whether these proteins fulfill any function remains to be determined [61]. Because of their length, pseudogene RNA are sometimes regarded as lncRNAs [79].

In head and neck cancer, it was proven that lncPTENP1, which is in fact a pseudotranscript of the *PTEN* gene, is under-expressed in the HN4, HN6, HN13, HN30 and Cal27 cell lines, being associated with lower survival rate in HNSCC patients. Through lentiviral transfection and restoration of the expression level of PTENP1, it was proven that, in vitro, the cells had decreased proliferation, invasion and migration capabilities, while, in vivo, they failed to form tumors [62]. The PTENP1 pseudogene transcript can bind the same micro-RNAs as the original gene *PTEN*, more precisely the miR-17, 21, 214, 9, and 26 families and act as a decoy mechanism that allows the translation of the *PTEN* gene [129]. The pseudogenes have *cis*-regulating sequences that compete with the *trans*-regulators of the protein-coding mRNAs [130].

Some rare ncRNAs are transcribed from pseudogenes. For instance, apart from the 5 YDNA genes located in the proximity of one another, it was reported that there are some YDNA pseudogenes that are spread throughout the whole genome. These pseudogenes existence was probably due to the retrotransposition of *trans*-mutated human YRNAs by the long-interspersed element-1 (L1). These hY-pseudogenes (hYRNAs) are also transcribed into a novel type of human yRNAs that may add a higher degree of complexity to the five hYRNAs [131].

### 11. miRNA-Offset RNAs

moRNAs were discovered in the chordate *Ciona intestinalis* through small RNAseq in 2009 [132]. They are enriched in the nucleus, being related with the regulation of nuclear processes, particularly transcription initiation and splicing activity [133]. moRNAs are similar in length and structure with miRNA. They have a length close to 20 nt. The biogenesis mechanisms are related to Drosha/DGCR8 complex that processes the multiple stem-loops of pri-miRNA transcript into the stem-loop form of pre-miRNA. The sequences that were thought to be discarded are kept in the nucleolus, giving rise to moRNAs [80].

With the small RNAseq technique, which compared the human embryonic stem cells with the normal fibroblast cell line, it was found that 326 moRNAs were in stem cells compared with 65 found in normal fibroblasts. The most abundant human moRNAs include moR-367-3p, moR-103a-2-3p, moR-92a-1-3p, moR-16-2-3p, moR-21-3p, moR-221-3p, moR-103a-1-3p, moR-20b-3p, moR-421-5p, and moR-21-5p [80]. Although their exact role in oral cancer remains unknown, they have potential to become well-studied ncRNAs because of their similarity with miRNAs.

### 12. Ultraconserved Regions of the DNA

UCR are sequences of the genome located in the exonic, intronic or even intergenic regions. In 2007, through microarray analysis, it was discovered that these UCRs are transcribed, giving rise to a new type of long non-coding RNAs, called T-UCRs [43]. They are highly similar across species and can bear up to 100% similarity in the same species. Surprisingly, it was discovered that the T-UCRs are dysregulated in multiple human cancers, such as bladder cancer, colorectal cancer, hepatocellular carcinoma, prostate cancer and chronic lymphocytic leukemia (reviewed in [81]).

In an esophageal cancer subtype, Barrett's adenocarcinoma, T-UCRs were evaluated. The down-regulated T-UCRs, i.e., uc.214+, uc.328+, uc.329+ and uc.356+, and up-regulated T-UCRs, i.e., uc.202−, uc.223−, and uc.269−, were found to be associated with the carcinogenesis of this malignancy through tissue microarray analysis [134].

All the above mentioned new information about the lesser-studied transcripts has impacted cancer research. The pseudogene transcripts are dysregulated in the head and neck malignancies (Figure 3 and Table 2), suggesting that they can become valuable diagnostic, prognostic or even therapeutic markers [62].

**Table 2.** Rare types of ncRNAs retrieved to be dysregulated in head and neck cancer.

Type of ncRNA	Name/Code	Expression	Determination method	Clinical/In Vitro/In Vivo	Observation	Reference
syRNAs	RNY1, RNY4P17	Up				
	RNY4P1, RNY4P5 RNY4P8, RNY4P11 Y_RNA.725, Y_RNA.122 RNY4P20, RNY4P25 RNY4P27, Y_RNA.182 Y_RNA.257, RNY4P18 Y_RNA.44, RNY4P6 Y_RNA.668, Y_RNA.7 Y_RNA.796, Y_RNA.292 Y_RNA.662	Down	Deep sequencing with IlluminaHiSeq 2000	Clinical blood samples	Altered expression level in head and neck cancer	[104]
TERC/hTR		Up Up Up	qRT-PCR Gene copy number—FISH RT-PCR	in vitro UMSCC74A, UMSCC74B/TCGA OSCC—tissue samples vs. normal tissue oral cancer tissue vs. normal tissue	cell senescence inhibited installment of oral lesions associated with higher grade dysplasia or carcinoma	[111] [135] [110]
	SNORD35B	Down	RNA-Seq for lncRNA, (TCGA)/RT-PCR	in vitro normal cell lines OKF4 and OKF6 vs. oral cancer cell lines UMSCC-10B, UMSCC-22B, HN-1, HN-12, and HN-30	associated with poor prognostic mediate the pattern of 28S rRNA	[121]
snoRNAs	4qI-4,14qII-22,ACA17,U84, mgh18S-121, U18A, U8, 14qII-12, U28 ENSG00000263442, ENSG00000264591, ENSG00000265325, ENSG00000265607, ENSG00000266646, ENSG00000266755,	Down	Microarray	OSCC tissue vs. keratinized gums preserved at −80 °C	Potential biomarkers, prognostic value	[123]
	LINC340, SNORD116-25, SNORA11, SNORA21, SNORA47 and SNORA65	Up	microarray, RT-PCR	Formalin fixed and paraffin embedded (FFPE) samples of ameloblastoma clinical	better diagnostic tool through biomarkers and potential therapeutic targets	[122]
NATs	WRAP53	Up	qRT-PCR, Western-blot	ESCC cell lines EC109, EC9706, KYSE150, and KYSE180 and tissue from 134 oesophageal cancer patients	higher expression of this transcript is progression of oesophageal cancer	[76]
Pseudogenes transcript	PTENP1	Down	HN4, HN6, HN13Cal27, HN30 cell lines		proliferation, invasion and migration capabilitiesand decreased survival rate	[37]
T-UCRs	uc.214+, uc.328+, uc.329+, uc.356+, uc.202-, uc.223-uc.269-	Down Up	Microarray, RT-PCR, immunohistochemistry	oesophageal cancer	cancerous transformation	[134]

qRT-PCR: real-time quantitative reverse transcription PCR; FISH: fluorescent hybridization in situ; OSCC: oral squamous cell carcinoma; RT-PCR: reverse transcription polymerase chain reaction; lncRNA: long non-coding RNA.


### 13. Exosomes as Important Sources of Rare ncRNAs

Exosomes are small vesicles of 40–100 nanometers that can carry information from one cell to another [136]. These small vehicles are important due to their cargo content [137,138]. They carry miRNAs [139], siRNAs [140], mRNA and DNA fragments [141].

It was discovered that the exosomes can also carry yRNAs associated with small fragments of mRNA thus proving the fact that the initial stages of exosomes formation can include the whole machinery of RNA degradation [142]. The extracellular vesicles were proven to be loaded with a greater amount of RNY5, in different cancer cell lines [143]. In fact, the YRNA concentration in exosomes was detected to be in a greater quantity than in the intracellular environment. The RNAY5 is the second most abundant RNA species in the human epithelial cell line and breast cancer cell line [144]. In the exosomes, RNY1 and RNY4 were also found [142]. The small YRNA serum levels in healthy donors was comparable to that of miRNAs [73].

By using LRRC24, MDM2, and CDKN1A primers in a RT-PCR analysis, after RNA isolation from colorectal cancer cells-derived exosomes, it was established for the first time that the exosomes can also carry NATs [145]. Another important class of ncRNAs found in the exosomes were snoRNAs, compared even with their profile from the cell of origin [142].

Exosomes are also secreted in the saliva and the molecular profiling of exosomes was proposed as a better option for evaluation of oral health status (Figure 4), diagnosing oral cancer and assessing the progression of this malignancy [138,146]



**Figure 4.** Salivary exosome-mediated intracellular communication in oral cancer. (1) In the tumor cell, by the budding of membrane, the multivesicular endosome (MVE) is formed, containing small vesicles of 40–100 nm called exosomes. The MVE can have an endogenous or an exogenous origin. (2) The MVE can fuse with a lysosome and its content is destroyed. (3) The MVE can also fuse with the plasma membrane and release the exosomes in the saliva. (4) The exosomes circulate through the body fluids such as the saliva or the blood and reach a targeted cell. (5) In the targeted cell, the exosomes are recognized by the surface proteins and are internalized through endocytosis. (6) The exosome can also bind directly to the plasma membrane and release its content into the cytoplasm. (7) An exosome contains surface proteins necessary for the engulfment by the targeted cell. In its interior, the exosome contains ncRNAs, DNA fragment and other proteins. This cargo can influence the behavior of their targeted cell. The exosome can contain also viruses; hence, it can mediate the spreading of human papilloma virus or Epstein-Barr virus.

A focus should be on the cargo content related to viral exposure such as HPV or EBV, which is a predisposition factor for some cancer types. These small salivary vesicles can be used as an alternative model in biomarkers discovery.

Although currently exosomes from saliva are evaluated mostly for their miRNA and siRNA cargo, the future analysis of salivary exosomes can also examine their cargo of the lesser-studied ncRNAs.

#### 14. Conclusion and Perspective

Based on past experience with siRNA and miRNA, the current knowledge regarding the ncRNA world could see dramatic changes in the following years, by increasing the importance of these rare transcripts in the clinical application. However, for sYRNA, snoRNA, NAT and T-UCR involvement in head and neck cancer, data are scarce. As for the others ncRNAs, such as eRNAs, sisRNAs, tRFs and pseudogene transcripts, there are indications of their dysregulation in other cancer types.

Notwithstanding the fact that there is a great diversity of studies about ncRNAs, we have to bear in mind that it is still in its infancy. Consequently, there is yet much to be learned about the sources of ncRNAs; their various maturation steps; mechanism of action; and interactions with the DNA, proteins, other coding or ncRNAs, all of which could offer the scientific community a better view of head and neck cancers and new ideas for oncological personalized therapy. These are sustained by the latest progress related to the approaches used for the evaluation.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

1. Chung, C.H.; Dietz, A.; Gregoire, V.; Guzzo, M.; Hamoir, M.; Leemans, R.; Lefebvre, J.-L.; Licitra, L.; El-Naggar, A.K.; O'Sullivan, B.; et al. Head and Neck Cancer. In *Oxford Textbook of Oncology*, 3rd ed.; Kerr, D., Haller, D., Velde, C., Baumann, M., Eds.; Oxford Medicine: Oxford, UK, 2016.
2. Lim, H.; Kim, D.H.; Jung, H.Y.; Gong, E.J.; Na, H.K.; Ahn, J.Y.; Kim, M.Y.; Lee, J.H.; Choi, K.S.; Choi, K.D.; et al. Clinical significance of early detection of esophageal cancer in patients with head and neck cancer. *Gut Liver* **2015**, *9*, 159–165. [[CrossRef](#)] [[PubMed](#)]
3. Yete, S.; D'Souza, W.; Saranath, D. High-Risk Human Papillomavirus in Oral Cancer: Clinical Implications. *Oncology* **2018**, *94*, 133–141. [[CrossRef](#)] [[PubMed](#)]
4. Jiang, S.; Dong, Y. Human papillomavirus and oral squamous cell carcinoma: A review of HPV-positive oral squamous cell carcinoma and possible strategies for future. *Curr. Probl. Cancer* **2017**, *41*, 323–327. [[CrossRef](#)] [[PubMed](#)]
5. Catana, C.S.; Calin, G.A.; Berindan-Neagoe, I. Inflammation-miRNAs in Aging and Breast Cancer: Are They Reliable Players? *Front. Med.* **2015**, *2*. [[CrossRef](#)] [[PubMed](#)]
6. Wang, V.E.; Grandis, J.R.; Ko, A.H. New Strategies in Esophageal Carcinoma: Translational Insights from Signaling Pathways and Immune Checkpoints. *Clin. Cancer Res.* **2016**, *22*, 4283–4290. [[CrossRef](#)] [[PubMed](#)]
7. Irimie, A.I.; Braicu, C.; Cojocneanu-Petric, R.; Berindan-Neagoe, I.; Campian, R.S. Novel technologies for oral squamous carcinoma biomarkers in diagnostics and prognostics. *Acta Odontol. Scand.* **2015**, *73*, 161–168. [[CrossRef](#)] [[PubMed](#)]
8. Gulei, D.; Magdo, L.; Jurj, A.; Raduly, L.; Cojocneanu-Petric, R.; Moldovan, A.; Moldovan, C.; Florea, A.; Pasca, S.; Pop, L.A.; et al. The silent healer: miR-205-5p up-regulation inhibits epithelial to mesenchymal transition in colon cancer cells by indirectly up-regulating E-cadherin expression. *Cell Death Dis.* **2018**, *9*. [[CrossRef](#)] [[PubMed](#)]
9. Ling, H.; Vincent, K.; Pichler, M.; Fodde, R.; Berindan-Neagoe, I.; Slack, F.J.; Calin, G.A. Junk DNA and the long non-coding RNA twist in cancer genetics. *Oncogene* **2015**, *34*, 5003–5011. [[CrossRef](#)] [[PubMed](#)]
10. Berindan-Neagoe, I.; Monroig Pdel, C.; Pasculli, B.; Calin, G.A. MicroRNAome genome: A treasure for cancer diagnosis and therapy. *Cancer J. Clin.* **2014**, *64*, 311–336. [[CrossRef](#)] [[PubMed](#)]
11. Berindan-Neagoe, I.; Braicu, C.; Gulei, D.; Tomuleasa, C.; Calin, G.A. Noncoding RNAs in Lung Cancer Angiogenesis. In *Physiologic and Pathologic Angiogenesis—Signaling Mechanisms and Targeted Therapy*; Simionescu, D., Simionescu, A., Eds.; InTech: Rijeka, Croatia, 2017.



12. Braicu, C.; Calin, G.A.; Berindan-Neagoe, I. MicroRNAs and cancer therapy— from bystanders to major players. *Curr. Med. Chem.* **2013**, *20*, 3561–3573. [[CrossRef](#)] [[PubMed](#)]
13. Irimie, A.I.; Braicu, C.; Sonea, L.; Zimta, A.A.; Cojocneanu-Petric, R.; Tonchev, K.; Mehterov, N.; Diudea, D.; Buduru, S.; Berindan-Neagoe, I. A Looking-Glass of Non-coding RNAs in oral cancer. *Int. J. Mol. Sci.* **2017**, *18*, 2620. [[CrossRef](#)] [[PubMed](#)]
14. Pop-Bica, C.; Gulei, D.; Cojocneanu-Petric, R.; Braicu, C.; Petrut, B.; Berindan-Neagoe, I. Understanding the Role of Non-Coding RNAs in Bladder Cancer: From Dark Matter to Valuable Therapeutic Targets. *Int. J. Mol. Sci.* **2017**, *18*, 1514. [[CrossRef](#)] [[PubMed](#)]
15. Shapiro, J.A. Revisiting the central dogma in the 21st century. *Ann. N. Y. Acad. Sci.* **2009**, *1178*, 6–28. [[CrossRef](#)] [[PubMed](#)]
16. Crick, F. Central dogma of molecular biology. *Nature* **1970**, *227*, 561–563. [[CrossRef](#)] [[PubMed](#)]
17. Esteller, M. Non-coding RNAs in human disease. *Nat. Rev. Genet.* **2011**, *12*, 861–874. [[CrossRef](#)] [[PubMed](#)]
18. Catana, C.S.; Pichler, M.; Giannelli, G.; Mader, R.M.; Berindan-Neagoe, I. Non-coding RNAs, the Trojan horse in two-way communication between tumor and stroma in colorectal and hepatocellular carcinoma. *Oncotarget* **2017**, *8*, 29519–29534. [[CrossRef](#)] [[PubMed](#)]
19. Braicu, C.; Catana, C.; Calin, G.A.; Berindan-Neagoe, I. NCRNA combined therapy as future treatment option for cancer. *Curr. Pharm. Des.* **2014**, *20*, 6565–6574. [[CrossRef](#)] [[PubMed](#)]
20. Kumari, P.; Sampath, K. cncRNAs: Bi-functional RNAs with protein coding and non-coding functions. *Semin. Cell Dev. Biol.* **2015**, *47–48*, 40–51. [[CrossRef](#)] [[PubMed](#)]
21. Fabbri, M.; Garzon, R.; Andreeff, M.; Kantarjian, H.M.; Garcia-Manero, G.; Calin, G.A. MicroRNAs and noncoding RNAs in hematological malignancies: Molecular, clinical and therapeutic implications. *Leukemia* **2008**, *22*, 1095–1105. [[CrossRef](#)] [[PubMed](#)]
22. Braicu, C.; Cojocneanu-Petric, R.; Chira, S.; Truta, A.; Floares, A.; Petrut, B.; Achimas-Cadariu, P.; Berindan-Neagoe, I.; et al. Clinical and pathological implications of miRNA in bladder cancer. *Int. J. Nanomed.* **2015**, *10*, 791–800. [[CrossRef](#)] [[PubMed](#)]
23. Ling, H.; Pickard, K.; Ivan, C.; Isella, C.; Ikuo, M.; Mitter, R.; Spizzo, R.; Bullock, M.; Braicu, C.; Pileczki, V.; et al. The clinical and biological significance of MIR-224 expression in colorectal cancer metastasis. *Gut* **2016**, *65*, 977–989. [[CrossRef](#)] [[PubMed](#)]
24. Berindan-Neagoe, I.; Calin, G.A. Molecular pathways: microRNAs, cancer cells, and microenvironment. *Clin. Cancer. Res.* **2014**, *20*, 6247–6253. [[CrossRef](#)] [[PubMed](#)]
25. Ambros, V. microRNAs: Tiny regulators with great potential. *Cell* **2001**, *107*, 823–826. [[CrossRef](#)]
26. Lee, R.C.; Feinbaum, R.L.; Ambros, V. The *C. elegans* heterochronic gene *LIN-4* encodes small RNAs with antisense complementarity to *LIN-14*. *Cell* **1993**, *75*, 843–854. [[CrossRef](#)]
27. Rupaimoole, R.; Slack, F.J. MicroRNA therapeutics: Towards a new era for the management of cancer and other diseases. *Nat. Rev. Drug Discov.* **2017**, *16*, 203–222. [[CrossRef](#)] [[PubMed](#)]
28. Sannigrahi, M.K.; Sharma, R.; Panda, N.K.; Khullar, M. Role of non-coding RNAs in head and neck squamous cell carcinoma: A narrative review. *Oral Dis.* **2017**, *21*. [[CrossRef](#)] [[PubMed](#)]
29. Pileczki, V.; Braicu, C.; Gherman, C.D.; Berindan-Neagoe, I. *TNF- $\alpha$*  gene knockout in triple negative breast cancer cell line induces apoptosis. *Int. J. Mol. Sci.* **2012**, *14*, 411–420. [[CrossRef](#)] [[PubMed](#)]
30. Jurj, A.; Braicu, C.; Pop, L.A.; Tomuleasa, C.; Gherman, C.D.; Berindan-Neagoe, I. The new era of nanotechnology, an alternative to change cancer treatment. *Drug Des. Dev. Ther.* **2017**, *11*, 2871–2890. [[CrossRef](#)] [[PubMed](#)]
31. Irimie, A.I.; Braicu, C.; Pileczki, V.; Petrushev, B.; Soritau, O.; Campian, R.S.; Berindan-Neagoe, I. Knocking down of p53 triggers apoptosis and autophagy, concomitantly with inhibition of migration on SSC-4 oral squamous carcinoma cells. *Mol. Cell. Biochem.* **2016**, *419*, 75–82. [[CrossRef](#)] [[PubMed](#)]
32. Singh, A.; Trivedi, P.; Jain, N.K. Advances in siRNA delivery in cancer therapy. *Artif. Cells Nanomed. Biotechnol.* **2018**, *46*, 274–283. [[CrossRef](#)] [[PubMed](#)]
33. Dana, H.; Chalbatani, G.M.; Mahmoodzadeh, H.; Karimloo, R.; Rezaiean, O.; Moradzadeh, A.; Mehmandoost, N.; Moazzen, F.; Mazraeh, A.; Marmari, V. Molecular Mechanisms and Biological Functions of siRNA. *Int. J. Biomed. Sci.* **2017**, *13*, 48–57. [[PubMed](#)]
34. Irimie, A.I.; Sonea, L.; Jurj, A.; Mehterov, N.; Zimta, A.A.; Budisan, L.; Braicu, C.; Berindan-Neagoe, I. Future trends and emerging issues for nanodelivery systems in oral and oropharyngeal cancer. *Int. J. Nanomed.* **2017**, *12*, 4593–4606. [[CrossRef](#)] [[PubMed](#)]

35. Berindan-Neagoe, I.; Braicu, C.; Irimie, A. Combining the chemotherapeutic effects of epigallocatechin 3-gallate with siRNA-mediated p53 knock-down results in synergic pro-apoptotic effects. *Int. J. Nanomed.* **2012**, *7*, 6035–6047.
36. Lecaros, R.L.; Huang, L.; Lee, T.C.; Hsu, Y.C. Nanoparticle Delivered VEGF-A siRNA Enhances Photodynamic Therapy for Head and Neck Cancer Treatment. *Mol. Ther.* **2016**, *24*, 106–116. [[CrossRef](#)] [[PubMed](#)]
37. Chen, W.H.; Lecaros, R.L.; Tseng, Y.C.; Huang, L.; Hsu, Y.C. Nanoparticle delivery of HIF1 $\alpha$  siRNA combined with photodynamic therapy as a potential treatment strategy for head-and-neck cancer. *Cancer Lett.* **2015**, *359*, 65–74. [[CrossRef](#)] [[PubMed](#)]
38. Taft, R.J.; Simons, C.; Nahkuri, S.; Oey, H.; Korbie, D.J.; Mercer, T.R.; Holst, J.; Ritchie, W.; Wong, J.J.; Rasko, J.E.; et al. Nuclear-localized tiny RNAs are associated with transcription initiation and splice sites in metazoans. *Nat. Struct. Mol. Biol.* **2010**, *17*, 1030–1034. [[CrossRef](#)] [[PubMed](#)]
39. Umbach, J.L.; Strelow, L.L.; Wong, S.W.; Cullen, B.R. Analysis of rhesus rhadinovirus microRNAs expressed in virus-induced tumors from infected rhesus macaques. *Virology* **2010**, *405*, 592–599. [[CrossRef](#)] [[PubMed](#)]
40. Zhao, J.; Schnitzler, G.R.; Iyer, L.K.; Aronovitz, M.J.; Baur, W.E.; Karas, R.H. MicroRNA-Offset RNA Alters Gene Expression and Cell Proliferation. *PLoS ONE* **2016**, *11*. [[CrossRef](#)] [[PubMed](#)]
41. Jacquot, J.; Tournier, J.M.; Puchelle, E. In vitro evidence that human airway lysozyme is cleaved and inactivated by *Pseudomonas aeruginosa* elastase and not by human leukocyte elastase. *Infect. Immun.* **1985**, *47*, 555–560. [[PubMed](#)]
42. Lujambio, A.; Portela, A.; Liz, J.; Melo, S.A.; Rossi, S.; Spizzo, R.; Croce, C.M.; Calin, G.A.; Esteller, M. CpG island hypermethylation-associated silencing of non-coding RNAs transcribed from ultraconserved regions in human cancer. *Oncogene* **2010**, *29*, 6390–6401. [[CrossRef](#)] [[PubMed](#)]
43. Calin, G.A.; Liu, C.G.; Ferracin, M.; Hyslop, T.; Spizzo, R.; Sevignani, C.; Fabbri, M.; Cimmino, A.; Lee, E.J.; Wojcik, S.E.; et al. Ultraconserved regions encoding ncRNAs are altered in human leukemias and carcinomas. *Cancer Cell* **2007**, *12*, 215–229. [[CrossRef](#)] [[PubMed](#)]
44. Fabris, L.; Calin, G.A. Understanding the Genomic Ultraconservations: T-UCRs and Cancer. *Int. Rev. Cell. Mol. Biol.* **2017**, *333*, 159–172. [[PubMed](#)]
45. Terreri, S.; Durso, M.; Colonna, V.; Romanelli, A.; Terracciano, D.; Ferro, M.; Perdon, S.; Castaldo, L.; Febbraio, F.; de Nigris, F.; et al. New Cross-Talk Layer between Ultraconserved Non-Coding RNAs, MicroRNAs and Polycomb Protein YY1 in Bladder Cancer. *Genes* **2016**, *7*, 127. [[CrossRef](#)] [[PubMed](#)]
46. Pek, J.W.; Osman, I.; Tay, M.L.; Zheng, R.T. Stable intronic sequence RNAs have possible regulatory roles in *Drosophila melanogaster*. *J. Cell Biol.* **2015**, *211*, 243–251. [[CrossRef](#)] [[PubMed](#)]
47. Osman, I.; Tay, M.L.; Pek, J.W. Stable intronic sequence RNAs (sisRNAs): A new layer of gene regulation. *Cell. Mol. Life Sci.* **2016**, *73*, 3507–3519. [[CrossRef](#)] [[PubMed](#)]
48. Yin, Q.F.; Yang, L.; Zhang, Y.; Xiang, J.F.; Wu, Y.W.; Carmichael, G.G.; Chen, L.L. Long noncoding RNAs with snoRNA ends. *Mol. Cell.* **2012**, *48*, 219–230. [[CrossRef](#)] [[PubMed](#)]
49. Ono, M.; Yamada, K.; Avolio, F.; Scott, M.S.; van Koningsbruggen, S.; Barton, G.J.; Lamond, A.I. Analysis of human small nucleolar RNAs (snoRNA) and the development of snoRNA modulator of gene expression vectors. *Mol. Biol. Cell* **2010**, *21*, 1569–1584. [[CrossRef](#)] [[PubMed](#)]
50. Stein, A.J.; Fuchs, G.; Fu, C.; Wolin, S.L.; Reinisch, K.M. Structural insights into RNA quality control: The Ro autoantigen binds misfolded RNAs via its central cavity. *Cell* **2005**, *121*, 529–539. [[CrossRef](#)] [[PubMed](#)]
51. Christov, C.P.; Gardiner, T.J.; Szuts, D.; Krude, T. Functional requirement of noncoding Y RNAs for human chromosomal DNA replication. *Mol. Cell. Biol.* **2006**, *26*, 6993–7004. [[CrossRef](#)] [[PubMed](#)]
52. Kim, M.S.; Pinto, S.M.; Getnet, D.; Nirujogi, R.S.; Manda, S.S.; Chaerkady, R.; Madugundu, A.K.; Kelkar, D.S.; Isserlin, R.; Jain, S.; et al. A draft map of the human proteome. *Nature* **2014**, *509*, 575–581. [[CrossRef](#)] [[PubMed](#)]
53. Rahman, S.; Zorca, C.E.; Traboulsi, T.; Noutahi, E.; Krause, M.R.; Mader, S.; Zenklusen, D. Single-cell profiling reveals that eRNA accumulation at enhancer-promoter loops is not required to sustain transcription. *Nucleic Acids Res.* **2017**, *45*, 3017–3030. [[CrossRef](#)] [[PubMed](#)]
54. Tseng, C.K.; Wang, H.F.; Burns, A.M.; Schroeder, M.R.; Gaspari, M.; Baumann, P. Human Telomerase RNA Processing and Quality Control. *Cell Rep.* **2015**, *13*, 2232–2243. [[CrossRef](#)] [[PubMed](#)]
55. Martinez, G.; Choudury, S.G.; Slotkin, R.K. tRNA-derived small RNAs target transposable element transcripts. *Nucleic Acids Res.* **2017**, *45*, 5142–5152. [[CrossRef](#)] [[PubMed](#)]

56. Lee, Y.S.; Shibata, Y.; Malhotra, A.; Dutta, A. A novel class of small RNAs: tRNA-derived RNA fragments (tRFs). *Genes Dev.* **2009**, *23*, 2639–2649. [[CrossRef](#)] [[PubMed](#)]
57. Werner, A.; Swan, D. What are natural antisense transcripts good for? *Biochem. Soc. Trans.* **2010**, *38*, 1144–1149. [[CrossRef](#)] [[PubMed](#)]
58. Li, N.; Joska, T.M.; Ruesch, C.E.; Coster, S.J.; Belden, W.J. The frequency natural antisense transcript first promotes, then represses, frequency gene expression via facultative heterochromatin. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 4357–4362. [[CrossRef](#)] [[PubMed](#)]
59. Johnson, T.S.; Li, S.; Kho, J.R.; Huang, K.; Zhang, Y. Network analysis of pseudogene-gene relationships: From pseudogene evolution to their functional potentials. *Pac. Symp. Biocomput.* **2018**, *23*, 536–547. [[PubMed](#)]
60. Ji, Z.; Song, R.; Regev, A.; Struhl, K. Many lncRNAs, 5'UTRs, and pseudogenes are translated and some are likely to express functional proteins. *eLife* **2015**, *4*. [[CrossRef](#)] [[PubMed](#)]
61. Xu, J.; Zhang, J. Are Human Translated Pseudogenes Functional? *Mol. Biol. Evol.* **2016**, *33*, 755–760. [[CrossRef](#)] [[PubMed](#)]
62. Poliseno, L.; Marranci, A.; Pandolfi, P.P. Pseudogenes in Human Cancer. *Front. Med.* **2015**, *2*. [[CrossRef](#)] [[PubMed](#)]
63. Chan, J.J.; Kwok, Z.H.; Chew, X.H.; Zhang, B.; Liu, C.; Soong, T.W.; Yang, H.; Tay, Y. A *FTH1* gene: Pseudogene: microRNA network regulates tumorigenesis in prostate cancer. *Nucleic Acids Res.* **2017**, gkx1248. [[CrossRef](#)]
64. Fang, Y.; Fullwood, M.J. Roles, Functions, and Mechanisms of Long Non-coding RNAs in Cancer. *Genom. Proteom. Bioinform.* **2016**, *14*, 42–54. [[CrossRef](#)] [[PubMed](#)]
65. Paraskevopoulou, M.D.; Hatzigeorgiou, A.G. Analyzing miRNA-lncRNA Interactions. *Methods Mol. Biol.* **2016**, *1402*, 271–286. [[PubMed](#)]
66. Jung, H.; Lee, D.; Lee, J.; Park, D.; Kim, Y.J.; Park, W.Y.; Hong, D.; Park, P.J.; Lee, E. Intron retention is a widespread mechanism of tumor-suppressor inactivation. *Nat. Genet.* **2015**, *47*, 1242–1248. [[CrossRef](#)] [[PubMed](#)]
67. Shlyueva, D.; Stampfel, G.; Stark, A. Transcriptional enhancers: From properties to genome-wide predictions. *Nat. Rev. Genet.* **2014**, *15*, 272–286. [[CrossRef](#)] [[PubMed](#)]
68. Lai, F.; Shiekhattar, R. Enhancer RNAs: The new molecules of transcription. *Curr. Opin. Genet. Dev.* **2014**, *25*, 38–42. [[CrossRef](#)] [[PubMed](#)]
69. Melamed, P.; Yosefzon, Y.; Rudnizky, S.; Pnueli, L. Transcriptional enhancers: Transcription, function and flexibility. *Transcription* **2016**, *7*, 26–31. [[CrossRef](#)] [[PubMed](#)]
70. Zhang, Y.; Zhang, X.O.; Chen, T.; Xiang, J.F.; Yin, Q.F.; Xing, Y.H.; Zhu, S.; Yang, L.; Chen, L.L. Circular intronic long noncoding RNAs. *Mol. Cell* **2013**, *51*, 792–806. [[CrossRef](#)] [[PubMed](#)]
71. Zheng, S.; Vuong, B.Q.; Vaidyanathan, B.; Lin, J.Y.; Huang, F.T.; Chaudhuri, J. Non-coding RNA Generated following Lariat Debranching Mediates Targeting of AID to DNA. *Cell* **2015**, *161*, 762–773. [[CrossRef](#)] [[PubMed](#)]
72. Kowalski, M.P.; Krude, T. Functional roles of non-coding Y RNAs. *Int. J. Biochem. Cell. Biol.* **2015**, *66*, 20–29. [[CrossRef](#)] [[PubMed](#)]
73. Dhahbi, J.M.; Spindler, S.R.; Atamna, H.; Boffelli, D.; Mote, P.; Martin, D.I. 5'-YRNA fragments derived by processing of transcripts from specific YRNA genes and pseudogenes are abundant in human serum and plasma. *Physiol. Genom.* **2013**, *45*, 990–998. [[CrossRef](#)] [[PubMed](#)]
74. Hizir, Z.; Bottini, S.; Grandjean, V.; Trabucchi, M.; Repetto, E. RNY (YRNA)-derived small RNAs regulate cell death and inflammation in monocytes/macrophages. *Cell Death Dis.* **2017**, *8*. [[CrossRef](#)] [[PubMed](#)]
75. Rubtsova, M.P.; Vasilkova, D.P.; Naraykina, Y.V.; Dontsova, O.A. Peculiarities of Yeasts and Human Telomerase RNAs Processing. *Acta Nat.* **2016**, *8*, 14–22. [[PubMed](#)]
76. Rao, X.; Huang, D.; Sui, X.; Liu, G.; Song, X.; Xie, J.; Huang, D. Overexpression of WRAP53 is associated with development and progression of esophageal squamous cell carcinoma. *PLoS ONE* **2014**, *9*. [[CrossRef](#)] [[PubMed](#)]
77. Scott, M.S.; Ono, M. From snoRNA to miRNA: Dual function regulatory non-coding RNAs. *Biochimie* **2011**, *93*, 1987–1992. [[CrossRef](#)] [[PubMed](#)]
78. Michel, C.I.; Holley, C.L.; Scruggs, B.S.; Sidhu, R.; Brookheart, R.T.; Listenberger, L.L.; Behlke, M.A.; Ory, D.S.; Schaffer, J.E. Small nucleolar RNAs U32a, U33, and U35a are critical mediators of metabolic stress. *Cell Metab.* **2011**, *14*, 33–44. [[CrossRef](#)] [[PubMed](#)]

79. Liu, J.; Xing, Y.; Xu, L.; Chen, W.; Cao, W.; Zhang, C. Decreased expression of pseudogene PTENP1 promotes malignant behaviours and is associated with the poor survival of patients with HNSCC. *Sci. Rep.* **2017**, *7*. [[CrossRef](#)] [[PubMed](#)]
80. Asikainen, S.; Heikkinen, L.; Juhila, J.; Holm, F.; Weltner, J.; Trokovic, R.; Mikkola, M.; Toivonen, S.; Balboa, D.; Lampela, R.; et al. Selective microRNA-Offset RNA expression in human embryonic stem cells. *PLoS ONE* **2015**, *10*. [[CrossRef](#)] [[PubMed](#)]
81. Terracciano, D.; Terreri, S.; de Nigris, F.; Costa, V.; Calin, G.A.; Cimmino, A. The role of a new class of long noncoding RNAs transcribed from ultraconserved regions in cancer. *Biochim. Biophys. Acta* **2017**, *1868*, 449–455. [[CrossRef](#)] [[PubMed](#)]
82. Kim, T.K.; Hemberg, M.; Gray, J.M.; Costa, A.M.; Bear, D.M.; Wu, J.; Harmin, D.A.; Laptewicz, M.; Barbara-Haley, K.; Kuersten, S.; et al. Widespread transcription at neuronal activity-regulated enhancers. *Nature* **2010**, *465*, 182–187. [[CrossRef](#)] [[PubMed](#)]
83. Lam, M.T.; Li, W.; Rosenfeld, M.G.; Glass, C.K. Enhancer RNAs and regulated transcriptional programs. *Trends Biochem. Sci.* **2014**, *39*, 170–182. [[CrossRef](#)] [[PubMed](#)]
84. Ren, C.; Liu, F.; Ouyang, Z.; An, G.; Zhao, C.; Shuai, J.; Cai, S.; Bo, X.; Shu, W. Functional annotation of structural ncRNAs within enhancer RNAs in the human genome: Implications for human disease. *Sci. Rep.* **2017**, *7*. [[CrossRef](#)] [[PubMed](#)]
85. Kim, T.K.; Hemberg, M.; Gray, J.M. Enhancer RNAs: A class of long noncoding RNAs synthesized at enhancers. *Cold Spring Harb. Perspect. Biol.* **2015**, *7*. [[CrossRef](#)] [[PubMed](#)]
86. Lai, F.; Gardini, A.; Zhang, A.; Shiekhata, R. Integrator mediates the biogenesis of enhancer RNAs. *Nature* **2015**, *525*, 399–403. [[CrossRef](#)] [[PubMed](#)]
87. Li, W.; Notani, D.; Rosenfeld, M.G. Enhancers as non-coding RNA transcription units: Recent insights and future perspectives. *Nat. Rev. Genet.* **2016**, *17*, 207–223. [[CrossRef](#)] [[PubMed](#)]
88. Bal, E.; Park, H.S.; Belaid-Choucair, Z.; Kayserili, H.; Naville, M.; Madrange, M.; Chiticariu, E.; Hadj-Rabia, S.; Cagnard, N.; Kuonen, F.; et al. Mutations in ACTRT1 and its enhancer RNA elements lead to aberrant activation of Hedgehog signaling in inherited and sporadic basal cell carcinomas. *Nat. Med.* **2017**, *23*, 1226–1233. [[CrossRef](#)] [[PubMed](#)]
89. Bell, R.E.; Golan, T.; Sheinboim, D.; Malcov, H.; Amar, D.; Salamon, A.; Liron, T.; Gelfman, S.; Gabet, Y.; Shamir, R.; et al. Enhancer methylation dynamics contribute to cancer plasticity and patient mortality. *Genome Res.* **2016**, *26*, 601–611. [[CrossRef](#)] [[PubMed](#)]
90. Younger, S.T.; Kenzelmann-Broz, D.; Jung, H.; Attardi, L.D.; Rinn, J.L. Integrative genomic analysis reveals widespread enhancer regulation by p53 in response to DNA damage. *Nucleic Acids Res.* **2015**, *43*, 4447–4462. [[CrossRef](#)] [[PubMed](#)]
91. Rowley, H.; Sherrington, P.; Helliwell, T.R.; Kinsella, A.; Jones, A.S. p53 expression and p53 gene mutation in oral cancer and dysplasia. *Otolaryngol. Head Neck Surg.* **1998**, *118*, 115–123. [[CrossRef](#)]
92. Gardner, E.J.; Nizami, Z.F.; Talbot, C.C., Jr.; Gall, J.G. Stable intronic sequence RNA (sisRNA), a new class of noncoding RNA from the oocyte nucleus of *Xenopus tropicalis*. *Genes Dev.* **2012**, *26*, 2550–2559. [[CrossRef](#)] [[PubMed](#)]
93. Shimakage, M.; Horii, K.; Tempaku, A.; Kakudo, K.; Shirasaka, T.; Sasagawa, T. Association of Epstein-Barr virus with oral cancers. *Hum. Pathol.* **2002**, *33*, 608–614. [[CrossRef](#)] [[PubMed](#)]
94. Moss, W.N.; Steitz, J.A. Genome-wide analyses of Epstein-Barr virus reveal conserved RNA structures and a novel stable intronic sequence RNA. *BMC Genom.* **2013**, *14*. [[CrossRef](#)] [[PubMed](#)]
95. Farris, A.D.; O'Brien, C.A.; Harley, J.B. Y3 is the most conserved small RNA component of Ro ribonucleoprotein complexes in vertebrate species. *Gene* **1995**, *154*, 193–198. [[CrossRef](#)]
96. Kohn, M.; Pazaitis, N.; Huttelmaier, S. Why YRNAs? About Versatile RNAs and Their Functions. *Biomolecules* **2013**, *3*, 143–156. [[CrossRef](#)] [[PubMed](#)]
97. Mosig, A.; Guofeng, M.; Stadler, B.M.; Stadler, P.F. Evolution of the vertebrate Y RNA cluster. *Theory Biosci.* **2007**, *126*, 9–14. [[CrossRef](#)] [[PubMed](#)]
98. Hall, A.E.; Turnbull, C.; Dalmay, T. Y RNAs: Recent developments. *Biomol. Concepts* **2013**, *4*, 103–110. [[CrossRef](#)] [[PubMed](#)]
99. Perreault, J.; Perreault, J.P.; Boire, G. Ro-associated Y RNAs in metazoans: Evolution and diversification. *Mol. Biol. Evol.* **2007**, *24*, 1678–1689. [[CrossRef](#)] [[PubMed](#)]



100. Tolkach, Y.; Stahl, A.F.; Niehoff, E.-M.; Zhao, C.; Kristiansen, G.; Müller, S.C.; Ellinger, J. YRNA expression predicts survival in bladder cancer patients. *BMC Cancer* **2017**, *17*. [[CrossRef](#)] [[PubMed](#)]
101. Catalanotto, C.; Cogoni, C.; Zardo, G. MicroRNA in Control of Gene Expression: An Overview of Nuclear Functions. *Int. J. Mol. Sci.* **2016**, *17*, 1712. [[CrossRef](#)] [[PubMed](#)]
102. Hendrick, J.P.; Wolin, S.L.; Rinke, J.; Lerner, M.R.; Steitz, J.A. Ro small cytoplasmic ribonucleoproteins are a subclass of La ribonucleoproteins: Further characterization of the Ro and La small ribonucleoproteins from uninfected mammalian cells. *Mol. Cell. Biol.* **1981**, *1*, 1138–1149. [[CrossRef](#)] [[PubMed](#)]
103. Dhahbi, J.M.; Spindler, S.R.; Atamna, H.; Boffelli, D.; Martin, D.I.K. Deep Sequencing of Serum Small RNAs Identifies Patterns of 5' tRNA Half and YRNA Fragment Expression Associated with Breast Cancer. *Biomark. Cancer* **2014**, *6*, 37–47. [[CrossRef](#)] [[PubMed](#)]
104. Victoria Martinez, B.; Dhahbi, J.M.; Nunez Lopez, Y.O.; Lamperska, K.; Golusinski, P.; Luczewski, L.; Kolenda, T.; Atamna, H.; Spindler, S.R.; Golusinski, W.; et al. Circulating small non-coding RNA signature in head and neck squamous cell carcinoma. *Oncotarget* **2015**, *6*, 19246–19263. [[PubMed](#)]
105. Repetto, E.; Lichtenstein, L.; Hizir, Z.; Tekaya, N.; Benahmed, M.; Ruidavets, J.-B.; Zaragosi, L.-E.; Perret, B.; Bouchareychas, L.; Genoux, A.; et al. RNY-derived small RNAs as a signature of coronary artery disease. *BMC Med.* **2015**, *13*. [[CrossRef](#)] [[PubMed](#)]
106. Baena-Del Valle, J.A.; Zheng, Q.; Esopi, D.M.; Rubenstein, M.; Hubbard, G.K.; Moncaliano, M.C.; Hruszkewycz, A.; Vaghasia, A.; Yegnasubramanian, S.; Wheelan, S.J.; et al. MYC drives overexpression of telomerase RNA (hTR/TERC) in prostate cancer. *J. Pathol.* **2018**, *244*, 11–24. [[CrossRef](#)] [[PubMed](#)]
107. Feng, J.; Funk, W.D.; Wang, S.S.; Weinrich, S.L.; Avilion, A.A.; Chiu, C.P.; Adams, R.R.; Chang, E.; Allsopp, R.C.; Yu, J.; et al. The RNA component of human telomerase. *Science* **1995**, *269*, 1236–1241. [[CrossRef](#)] [[PubMed](#)]
108. Kilchert, C.; Wittmann, S.; Vasiljeva, L. The regulation and functions of the nuclear RNA exosome complex. *Nat. Rev. Mol. Cell Biol.* **2016**, *17*, 227–239. [[CrossRef](#)] [[PubMed](#)]
109. Shukla, S.; Schmidt, J.C.; Goldfarb, K.C.; Cech, T.R.; Parker, R. Inhibition of telomerase RNA decay rescues telomerase deficiency caused by dyskerin or PARN defects. *Nat. Struct. Mol. Biol.* **2016**, *23*, 286–292. [[CrossRef](#)] [[PubMed](#)]
110. Kim, H.R.; Christensen, R.; Park, N.H.; Sapp, P.; Kang, M.K.; Park, N.H. Elevated expression of hTERT is associated with dysplastic cell transformation during human oral carcinogenesis in situ. *Clin. Cancer Res.* **2001**, *7*, 3079–3086. [[PubMed](#)]
111. Majumder, M.; House, R.; Palanisamy, N.; Qie, S.; Day, T.A.; Neskey, D.; Diehl, J.A.; Palanisamy, V. RNA-Binding Protein FXR1 Regulates p21 and TERC RNA to Bypass p53-Mediated Cellular Senescence in OSCC. *PLoS Genet.* **2016**, *12*. [[CrossRef](#)]
112. Fernandez, E.; Mallette, F.A. The Rise of FXR1: Escaping Cellular Senescence in Head and Neck Squamous Cell Carcinoma. *PLoS Genet.* **2016**, *12*. [[CrossRef](#)] [[PubMed](#)]
113. Lacatena, R.M.; Cesareni, G. Base pairing of RNA I with its complementary sequence in the primer precursor inhibits ColE1 replication. *Nature* **1981**, *294*, 623–626. [[CrossRef](#)] [[PubMed](#)]
114. Rosikiewicz, W.; Makalowska, I. Biological functions of natural antisense transcripts. *Acta Biochim. Pol.* **2016**, *63*, 665–673. [[CrossRef](#)] [[PubMed](#)]
115. Coucoravas, C.; Dhanjal, S.; Henriksson, S.; Bohm, S.; Farnebo, M. Phosphorylation of the Cajal body protein WRAP53beta by ATM promotes its involvement in the DNA damage response. *RNA Biol.* **2017**, *14*, 804–813. [[CrossRef](#)] [[PubMed](#)]
116. Li, M.W.; Sletten, A.C.; Lee, J.; Pyles, K.D.; Matkovich, S.J.; Ory, D.S.; Schaffer, J.E. Nuclear export factor 3 regulates localization of small nucleolar RNAs. *J. Biol. Chem.* **2017**, *292*, 20228–20239. [[CrossRef](#)] [[PubMed](#)]
117. Holley, C.L.; Li, M.W.; Scruggs, B.S.; Matkovich, S.J.; Ory, D.S.; Schaffer, J.E. Cytosolic accumulation of small nucleolar RNAs (snoRNAs) is dynamically regulated by NADPH oxidase. *J. Biol. Chem.* **2015**, *290*, 11741–11748. [[CrossRef](#)] [[PubMed](#)]
118. Falaleeva, M.; Stamm, S. Processing of snoRNAs as a new source of regulatory non-coding RNAs: snoRNA fragments form a new class of functional RNAs. *Bioessays* **2013**, *35*, 46–54. [[CrossRef](#)] [[PubMed](#)]
119. Ellis, J.C.; Brown, D.D.; Brown, J.W. The small nucleolar ribonucleoprotein (snoRNP) database. *RNA* **2010**, *16*, 664–666. [[CrossRef](#)] [[PubMed](#)]



120. Kishore, S.; Gruber, A.R.; Jedlinski, D.J.; Syed, A.P.; Jorjani, H.; Zavolan, M. Insights into snoRNA biogenesis and processing from PAR-CLIP of snoRNA core proteins and small RNA sequencing. *Genome Biol.* **2013**, *14*. [[CrossRef](#)] [[PubMed](#)]
121. Zou, A.E.; Ku, J.; Honda, T.K.; Yu, V.; Kuo, S.Z.; Zheng, H.; Xuan, Y.; Saad, M.A.; Hinton, A.; Brumund, K.T.; et al. Transcriptome sequencing uncovers novel long noncoding and small nucleolar RNAs dysregulated in head and neck squamous cell carcinoma. *RNA* **2015**, *21*, 1122–1134. [[CrossRef](#)] [[PubMed](#)]
122. Davanian, H.; Balasiddaiah, A.; Heymann, R.; Sundstrom, M.; Redenstrom, P.; Silfverberg, M.; Brodin, D.; Sallberg, M.; Lindskog, S.; Kruger Weiner, C.; et al. Ameloblastoma RNA profiling uncovers a distinct non-coding RNA signature. *Oncotarget* **2017**, *8*, 4530–4542. [[CrossRef](#)] [[PubMed](#)]
123. Chamorro-Petronacci, C.; Perez-Sayans, M.; Padin-Iruegas, M.E.; Marichalar-Mendia, X.; Gallas-Torreira, M.; Garcia Garcia, A. Differential expression of snoRNAs in oral squamous cell carcinomas: New potential diagnostic markers. *J. Enzyme Inhib. Med. Chem.* **2018**, *33*, 424–427. [[CrossRef](#)] [[PubMed](#)]
124. Kunzmann, A.; Brennicke, A.; Marchfelder, A. 5' end maturation and RNA editing have to precede tRNA 3' processing in plant mitochondria. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 108–113. [[CrossRef](#)] [[PubMed](#)]
125. Betat, H.; Long, Y.; Jackman, J.E.; Mörl, M. From End to End: tRNA Editing at 5'- and 3'-Terminal Positions. *Int. J. Mol. Sci.* **2014**, *15*, 23975–23998. [[CrossRef](#)] [[PubMed](#)]
126. Pederson, T. Regulatory RNAs derived from transfer RNA? *RNA* **2010**, *16*, 1865–1869. [[CrossRef](#)] [[PubMed](#)]
127. Diebel, K.W.; Zhou, K.; Clarke, A.B.; Bemis, L.T. Beyond the Ribosome: Extra-translational Functions of tRNA Fragments. *Biomark. Insights* **2016**, *11*, 1–8. [[CrossRef](#)] [[PubMed](#)]
128. Guo, X.; Lin, M.; Rockowitz, S.; Lachman, H.M.; Zheng, D. Characterization of human pseudogene-derived non-coding RNAs for functional potential. *PLoS ONE* **2014**, *9*. [[CrossRef](#)] [[PubMed](#)]
129. Spizzo, R.; Almeida, M.I.; Colombatti, A.; Calin, G.A. Long non-coding RNAs and cancer: A new frontier of translational research? *Oncogene* **2012**, *31*, 4577–4587. [[CrossRef](#)] [[PubMed](#)]
130. Tutar, Y. Pseudogenes. *Comp. Funct. Genom.* **2012**, *2012*. [[CrossRef](#)] [[PubMed](#)]
131. Perreault, J.; Noel, J.F.; Briere, F.; Cousineau, B.; Lucier, J.F.; Perreault, J.P.; Boire, G. Retropseudogenes derived from the human Ro/SS-A autoantigen-associated hY RNAs. *Nucleic Acids Res.* **2005**, *33*, 2032–2041. [[CrossRef](#)] [[PubMed](#)]
132. Shi, W.; Hendrix, D.; Levine, M.; Haley, B. A distinct class of small RNAs arises from pre-miRNA-proximal regions in a simple chordate. *Nat. Struct. Mol. Biol.* **2009**, *16*, 183–189. [[CrossRef](#)] [[PubMed](#)]
133. Bortoluzzi, S.; Biasiolo, M.; Bisognin, A. MicroRNA-offset RNAs (moRNAs): By-product spectators or functional players? *Trends Mol. Med.* **2011**, *17*, 473–474. [[CrossRef](#)] [[PubMed](#)]
134. Fassan, M.; Dall'Olmo, L.; Galasso, M.; Braconi, C.; Pizzi, M.; Realdon, S.; Volinia, S.; Valeri, N.; Gasparini, P.; Baffa, R.; et al. Transcribed ultraconserved noncoding RNAs (T-UCR) are involved in Barrett's esophagus carcinogenesis. *Oncotarget* **2014**, *5*, 7162–7171. [[CrossRef](#)] [[PubMed](#)]
135. Boscolo-Rizzo, P.; Da Mosto, M.C.; Rampazzo, E.; Giunco, S.; Del Mistro, A.; Menegaldo, A.; Baboci, L.; Mantovani, M.; Tirelli, G.; De Rossi, A. Telomeres and telomerase in head and neck squamous cell carcinoma: From pathogenesis to clinical implications. *Cancer Metastasis Rev.* **2016**, *35*, 457–474. [[CrossRef](#)] [[PubMed](#)]
136. Gulei, D.; Irimie, A.I.; Cojocneanu-Petric, R.; Schultze, J.L.; Berindan-Neagoe, I. Exosomes-Small Players, Big Sound. *Bioconjug. Chem.* **2018**, *9*. [[CrossRef](#)] [[PubMed](#)]
137. Zhao, X.; Wu, Y.; Duan, J.; Ma, Y.; Shen, Z.; Wei, L.; Cui, X.; Zhang, J.; Xie, Y.; Liu, J. Quantitative proteomic analysis of exosome protein content changes induced by hepatitis B virus in Huh-7 cells using SILAC labeling and LC-MS/MS. *J. Proteome Res.* **2014**, *13*, 5391–5402. [[CrossRef](#)] [[PubMed](#)]
138. Yu, X.; Odenthal, M.; Fries, J.W.U. Exosomes as miRNA Carriers: Formation–Function–Future. *Int. J. Mol. Sci.* **2016**, *17*, 2028. [[CrossRef](#)] [[PubMed](#)]
139. Shahabipour, F.; Banach, M.; Sahebkar, A. Exosomes as nanocarriers for siRNA delivery: Paradigms and challenges. *Arch. Med. Sci.* **2016**, *12*, 1324–1326. [[CrossRef](#)] [[PubMed](#)]
140. Takahashi, A.; Okada, R.; Nagao, K.; Kawamata, Y.; Hanyu, A.; Yoshimoto, S.; Takasugi, M.; Watanabe, S.; Kanemaki, M.T.; Obuse, C.; et al. Exosomes maintain cellular homeostasis by excreting harmful DNA from cells. *Nat. Commun.* **2017**, *8*. [[CrossRef](#)] [[PubMed](#)]
141. Van Balkom, B.W.; Eisele, A.S.; Pegtel, D.M.; Bervoets, S.; Verhaar, M.C. Quantitative and qualitative analysis of small RNAs in human endothelial cells and exosomes provides insights into localized RNA processing, degradation and sorting. *J. Extracell. Vesicles* **2015**, *4*. [[CrossRef](#)] [[PubMed](#)]

142. Chakraborty, S.K.; Prakash, A.; Nechooshtan, G.; Hearn, S.; Gingeras, T.R. Extracellular vesicle-mediated transfer of processed and functional RNY5 RNA. *RNA* **2015**, *21*, 1966–1979. [[CrossRef](#)] [[PubMed](#)]
143. Tosar, J.P.; Gambaro, F.; Sanguinetti, J.; Bonilla, B.; Witwer, K.W.; Cayota, A. Assessment of small RNA sorting into different extracellular fractions revealed by high-throughput sequencing of breast cell lines. *Nucleic Acids Res.* **2015**, *43*, 5601–5616. [[CrossRef](#)] [[PubMed](#)]
144. Chiba, M.; Kimura, M.; Asari, S. Exosomes secreted from human colorectal cancer cell lines contain mRNAs, microRNAs and natural antisense RNAs, that can transfer into the human hepatoma HepG2 and lung cancer A549 cell lines. *Oncol. Rep.* **2012**, *28*, 1551–1558. [[CrossRef](#)] [[PubMed](#)]
145. Zlotogorski-Hurvitz, A.; Dayan, D.; Chaushu, G.; Korvala, J.; Salo, T.; Sormunen, R.; Vered, M. Human Saliva-Derived Exosomes: Comparing Methods of Isolation. *J. Histochem. Cytochem.* **2015**, *63*, 181–189. [[CrossRef](#)] [[PubMed](#)]
146. Principe, S.; Hui, A.B.; Bruce, J.; Sinha, A.; Liu, F.F.; Kislinger, T. Tumor-derived exosomes and microvesicles in head and neck cancer: Implications for tumor biology and biomarker discovery. *Proteomics* **2013**, *13*, 1608–1623. [[CrossRef](#)] [[PubMed](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).