

Review

# Surface Water under Growing Anthropogenic Loads: From Global Perspectives to Regional Implications

Tatyana I. Moiseenko 

V.I. Vernadsky Institute of Geochemistry and Analytical Chemistry RAS, Kosygina Street 19, 119991 Moscow, Russia; moiseenko.ti@gmail.com

**Abstract:** This paper reviews currently available evaluations of the effects of continuously increasing anthropogenic loads on water resources. The increase in the fluxes of elements and compounds into the environment, such as the emission of greenhouse gases and dispersion of nutrients (nitrogen and phosphorus), acidifying gases, and toxic elements and compounds that adversely affect water quality, are considered. The significance of fresh waters as a life support factor for human populations is demonstrated. Examples illustrate how key anthropogenic-induced processes develop in land waters under the effect of anthropogenic loads, as exemplified by the Russian Kola regions. Climate warming and the increasing dispersion of elements are demonstrated to result in the eutrophication of surface waters, even in areas remote from anthropogenically impacted regions. Although the emissions of acidifying gases diminish, the waters are still acidified in acid-vulnerable areas, and the chemical compositions of the waters have been significantly modified over the past decades, indicating that the changes in the chemical composition of the waters are irreversible. A new feature of the waters is distinguished: the toxicity of the habitats for aquatic organisms. The novelty of the work was to highlight the bridge between anthropogenic loads on surface water on a planetary scale and their consequences in certain regions, which reflect changes in water quality. Arguments are presented for establishing a theoretical approach for evaluating critical loads.

**Keywords:** surface water; anthropogenic loads; warming climate; eutrophication; acidification; critical loads



**Citation:** Moiseenko, T.I. Surface Water under Growing Anthropogenic Loads: From Global Perspectives to Regional Implications. *Water* **2022**, *14*, 3730. <https://doi.org/10.3390/w14223730>

Academic Editor:  
George Arhonditsis

Received: 28 September 2022

Accepted: 14 November 2022

Published: 17 November 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The explosive growth of the human population, the intense development of mineral resources, and technological progress in the 20th century dramatically affected the environment, as had become obvious by the mid-20th century. V.I. Vernadsky [1] stressed that human activities can fundamentally and rapidly modify both many natural processes and what is referred to as laws of nature.

The paramount significance of fresh waters for the human population of our planet and for preserving the species diversity on Earth demonstrates how important to study water resources affected by ever-growing anthropogenic loads. The transformations of catchment, airborne pollution, the discharge of industrial wastes and domestic sewage, and diffuse runoffs contaminate aquatic systems and change the biogeochemical cycles of elements in the systems of catchments and water bodies, leading to the acidification, salinization, eutrophication, and bring toxic compounds into the water body. These processes eventually downgrade the quality of the waters and diminish the biological diversity of the aquatic systems.


Assessment of anthropogenic impacts on the environment on a planetary scale and its negative consequences are presented in several works [2–6]. Many lakes and rivers on different continents experience multiple stressors. The withdrawal of the runoff and climate warming are also reflected in changes in the hydrogeological cycles and fluxes of elements from catchments to lakes and rivers [7,8]. Livestock and farm waste causes water

eutrophication in local surface water throughout the planet. Salinization in arid and semi-arid surface water appears due to increased evaporation in a warm climate. Dispersion of metals is manifested everywhere, especially in regions with developed industry and mining. Oil and gas production transportation affects land waters in case of accidents on pipelines during their transportation. The combustion of fuel, coal and fuel oil leads to the acidification of waters, mainly in regions where waters are sensitive to acid loads. Understanding all direct and indirect consequences of anthropogenic impacts on water resources is hard.

Numerous stressors are constantly degrading the quality of surface waters worldwide, creating many challenges for their conservation and restoration. The types of combined effects of multiple stressors range from the dominance of a single stressor to complex interactions. Identifying commonly combined impacts is critical to environmental management as it helps prioritize stressors for mitigation.

The work aimed to show the influence of increasing anthropogenic loads on surface water on a planetary scale and their consequences on the example of a number of regions, mainly the North of the European part of Russia or those regions where negative consequences for land waters have manifested themselves, such as salinization in a warm climate. This work focused mainly on the following problems (Figure 1):

- the use of water resources as a life necessity on the planet;
- the effect of climate warming on the eutrophication of waters;
- the consequences of acidic deposition for the water acidification;
- the transfer of toxic elements and compounds to natural waters;
- principal theoretical approaches to the evaluation of critical loads.



**Figure 1.** Flowchart showing anthropogenic loads on surface water and its consequences.

Analysis of currently available estimations of the quality of natural surface waters in the Arctic basin shows that this territory is still studied inadequately. Anthropogenic loads have impacted the northeastern part of the Scandinavia shield (northern Kola Peninsula) for more than eight decades, and the local waters are susceptible to all anthropogenic loads because of the sluggish material and energy exchange in the cold latitudes. Below we employ this region as an illustrative example to discuss, based on the results of our long-term studies in the area [9–15], how anthropogenic-induced processes evolve in the surface waters. In natural environments, the waters of the region are low in total dissolved salt (TDS) and are poor in biogenic elements (nutrients). In their pristine natural state, lakes in the Arctic basin are ultra-fresh and oligotrophic. Note that this water chemistry, as an anthropogenic load, is typical of numerous lakes in the Arctic zone of our planet. When reviewing the climate warming impact on surface water, several climatic regions, including arid ones, were considered.

## 2. The Importance of Freshwater in the Life Support

The worldwide utilization of freshwaters is evaluated at 2600 km<sup>3</sup> per year and was 415 km<sup>3</sup> per year in the preindustrial epoch [2]. Nowadays, approximately 70% of the world's freshwater resources are used for agricultural needs, industries utilize 20%, and 10% is consumed for domestic purposes. According to global evaluations [3], the allowable global water withdrawal threshold has not been exceeded as of yet. The global consumption of freshwater by humankind amounts to 4000 km<sup>3</sup> per year.

It should be mentioned that water resources are unevenly distributed over the planet's surface, and many of these resources occur in northern areas that are sparsely populated. For example, the average long-term total runoff of northern rivers is estimated at 4300 km<sup>3</sup>, which is commensurable with the allowable freshwater consumption by the whole world's human population. The northern areas are populated very sparsely. The total volume of secular freshwater resources stored in Russia's lakes is 26,500 km<sup>3</sup>. (including 23,000 km<sup>3</sup> in Lake Baikal, 903 km<sup>3</sup> in Ladoga Lake, and 295 km<sup>3</sup> in Onega Lake). First, the runoff of large Siberian rivers, Ob, Lena, and Yenisei, amounts to 1600 km<sup>3</sup> per year, i.e., one-third of the world's riverine runoff [16].

The greatest deficit of water resources (not only in quantity but also in the quality of the waters) occurs in densely populated steppe and arid parts of our planet, where more than 40% of the world's human population currently lives. Statistical data indicate that practically one-fifth of the global population suffers from an acute shortage of drinkable water [4]. According to scientific estimates, more than half of countries worldwide will suffer from serious water shortage or insufficient supply by 2025 and three-quarters of the Earth's population will have been critically short in freshwater by the mid-21st century. According to estimates, 47% of the world's population will have been seriously threatened by water shortage by 2030. The population will significantly grow by 2050 in rapidly developing countries, which are already short of water [17].

Along with the water deficit, an important problem is the contamination of freshwaters with industrial emissions and discharges, the runoff of fertilizer-contaminated waters from urbanized territories, atmospheric fallouts, and the penetration of salty waters into fresh aquifers in coastal areas because of groundwater withdrawal. Operating along with such global processes as climate warming, anthropogenic loads result in both quantitative and qualitative depletion of water resources.

## 3. Effect of Climate Warming on Aquatic Systems

Lately, many researchers worldwide have focused on the effects of climate warming. The planet's average air temperature has increased by 1.5 °C since 1980, and the Earth's surface continues to warm at a rate of approximately 0.16 °C per decade, varying from one region to another. It has been proved that the warming of the biosphere brings about weather instability, along with changes in atmospheric precipitation and disturbances in hydrological cycles (longer arid periods and expansion of deserts in a warm climate with a simultaneous increase in precipitation and flooding in humid zones). The principal reason for climate warming is the ever-growing emission of greenhouse gases, first of all, CO<sub>2</sub>. The atmosphere now contains 42% more CO<sub>2</sub> than at the beginning of the industrial era [5]. The latest IPCC report shows that emissions of greenhouse gases continue to rise, and current plans to address climate change are not ambitious enough to limit warming to 1.5 °C above the preindustrial level [6].

Global warming intensifies bio-cycling in the freshwater system and facilitates the establishment of feedback and changes in the environment, landscapes, and human society. An increase in air temperature, particularly in summertime and early autumn, means that the atmosphere can retain more water [8]. Climate changes also brings more moisture from lower latitudes to the pole. This increases the amount of precipitation in the Arctic, with this precipitation falling off in the form of either rain or snowfalls. In many parts of the Arctic, the amount of precipitation in the form of rain (but not snow) has increased, and the snow-cover period has become shorter [18].

Climate warming also affects runoff [19]. An increase in the precipitation, runoff and ice melting in ice covers results in a greater freshwater flow into the Arctic Ocean. For example, the riverine runoff was estimated to increase to  $4200 \text{ km}^3 (\pm 420 \text{ km}^3)$  in 2000–2010 compared to  $3900 \text{ km}^3 (\pm 390 \text{ km}^3)$  in 1980–2000. These changes are predicted to continue, and numerical simulations indicate that the riverine runoff will perhaps increase by 25 to 50% over most of the Arctic [18]. The example of large Siberian rivers, such as Ob (at the gauging section in Salekhard), Yenisei (Igarka), and Lena (Kyusyur), indicates that an increase in the runoff of these rivers was simultaneous with the onset of the modern air warming. It has been demonstrated that the long-lasting phases of runoff changes are synchronous with phases of changes in the air temperature and large-scale atmospheric circulation [20].


The most hazardous phenomenon is that warming impacts permafrost rocks (PFR), widespread in continental West Siberia over an integral area of about 700 thousand  $\text{km}^2$ , i.e., more than one-fifth of the area. The thawing of permafrost peatlands in northern West Siberia may increase the release of methane and other greenhouse gases and augment water volumes in the rivers and lakes. The intensification of the thermokarst process should increase the number of lakes and their surface area [21–24].

The snow cover is the main source of annual runoff water in the summertime in northern territories and controls the export dynamics of nutrients and dissolved organic carbon (DOC). Changes in the hydrological cycles should inevitably modify concentrations and remove chemical elements in the waters, including suspending matter, DOC, and nutrients [25]. Climate warming should change (with regard to the predictions) the time and intensity of snow melting, which in turn, should change the runoff to the lakes and marginal seas. In addition, the frequency of autumn storms and floods may increase [26]. It is still largely uncertain how these factors may correlate with changes in the transfer of dissolved compounds from the catchments, but it is evident that the biogeochemical cycles in the catchment–water body system should thereby change.


The literature presents ample evidence that climate changes result in changes in the chemical composition of the waters [27–29]. D. Houle et al. [30] mention that higher annual average air temperatures correlate with the pH of the lake waters in long time series of monitoring data. The increase in organic matter content may be explained by the increase in the influx of nutrients (nitrates and phosphates) from the catchments as a consequence of climate warming [14,31].

The influence of temperature on water resources is controlled primarily through changes in the hydrological conditions under which the waters are formed and through biogeochemical cycles, i.e., the amount of precipitation, the occurrence of a snow cover, the conditions of the rocks underlying soils at the catchments, the saturation or depletion of the waters with exchange bases and accumulation at the catchment over the historical period, microbiological activity, the acceleration of vegetation growth, and perhaps, also the runoff of nutrients.

It has been demonstrated [11] how an elevated temperature affects the chemical composition of waters in various natural climatic zones (from the Arctic to steppes). These data indicate that temperature most strongly affects water eutrophication. Three-parameter dependencies were derived for parameters of the chemical composition of the waters on climatic parameters at the catchments. These dependencies provided a basis for prognostic models that enabled predicting the probable changes in salt and phosphorus concentrations in surface waters at warming for 0.5, 1.0, 1.5, and 2.0 °C (Figure 2).



(a)




(b)

**Figure 2.** The observed and predicted values of sum ions (a) and total phosphorus (b) for the climatic ecoregions within European Russia along a climatic transect from northern tundra to the southern arid zone [11].

According to the calculated dependencies, the intensity of chemical weathering at a temperature increase of 0.5 and 1 °C should not result in any significant increase in the total dissolved salts (TDS) of waters in northern areas. However, a notable increase (about 25%) in the total salt concentrations should occur in central and southern regions of the European part of Russia at a temperature increase of 2 °C. A significant increase (by approximately 50%) in the total phosphorus concentration in lake waters should occur practically everywhere (except only tundra and forested tundra territories), already at an increase in the average daily temperatures by 0.5 °C. Phosphorus concentrations should particularly increase in arid zones, in which climate warming should result in significant eutrophication of the waters.

Lately acquired data indicate that the temperature in the Arctic has significantly increased, and this resulted in a notable increase in concentrations of nutrients in the waters [14], which will be discussed below. Climate warming thus changes not only hydrological conditions under which the waters are formed but also modifies the chemical composition of the waters: the salt concentrations increase, as also do the concentrations of nutrients, and this should be most clearly seen in southern regions and also be discernible in the Arctic (Figure 3).



**Figure 3.** An example of an increase in mean monthly surface air temperature  $T$  (°C) a 28 year period (1990–2018) on the territory of the Kola North. Significant positive slope of the temperature ( $T$ ) SKT = (+4.71),  $p < 0.001$ .

Climate warming can stimulate the processes of salinization of lakes in arid and semi-arid regions due to an increase in evaporative concentration. This phenomenon has been established for a number of lakes in arid zones in North America, Kazakhstan, South Russia, Iran (Lake Urmia), and other water bodies [32–36]. The Crimean lakes of Russia on the western coast of the peninsula are located in the arid climate zone, they are typical hypersaline reservoirs, and in summer, they dry out at high air and water temperatures. The dynamics of changes in their ecological state during the year season may reflect the process of ecosystem transformation during the period of extremely hot conditions in the summer period, followed by restoration to normal functioning in the autumn-winter period [35]. From the cited works, it is concluded that climate warming consequences are most dramatic in the land waters of the southern regions. At the same time, there are changes in the water of the northern regions in the direction of increasing their trophic status.

#### 4. Eutrophication

Climate warming most significantly affects the biogeochemical cycling of nutrients and organic matter by stimulating eutrophication. Eutrophication is greatly contributed by the global anthropogenic dispersion of phosphorus and nitrogen. The main reason for freshwater eutrophication is the increase in nitrogen and phosphorus anthropogenic loads on the environment has intensified worldwide over the past years [3,37–39].

The overall runoff of phosphorus to the ocean is evaluated to have a nine to tenfold increase compared to the preindustrial period. Natural phosphorus comes to the environment through the chemical weathering of rocks, and approximately 1.1 million tons of natural phosphorus is annually introduced into the natural cycling of this element. Several models suggest that global phosphorus loads in the runoff and leaching into freshwater may vary from 0.16 to 5 TgP per year [40,41].

Anthropogenic activities modify the cycle of another nutrient element: nitrogen, with  $N_2$  converted into reactive nitrogen as a consequence of the following four processes:

industrial chemical binging (80 MegatonsN per year); agricultural chemical binging through the cultivation of leguminous (40 MegatonsN per year); combustion of effluent gases (20 MegatonsN per year); and biomass combustion at fires (10 MegatonsN per year) [42]. Both phosphorus and nitrogen are produced mostly for increasing agricultural productivity. Nowadays, it is not profitable to limit fertilizer treatments of soils because it is necessary to increase agricultural productivity to provide food for the ever-growing planet's population. NO<sub>x</sub> emissions prevail in countries with a high gross national product, whereas NH<sub>3</sub> is emitted mostly in countries oriented on food production [43].

Eutrophication is hazardous for waters utilized for drinking water supply, fish farming, recreational purposes, and industrial needs because this process intensifies the growth of cyanobacteria and aquatic macrophytes and thus decreases the oxygen contents and leads to the death of the aquatic flora and fauna. The state of thousands of lakes, mouths of reservoirs, and wetlands near large population centers worldwide worsens because of the increase in nitrogen and phosphorus concentrations [44,45].

The eutrophication of waters in the Arctic basin is still poorly described in the literature. No such problem is mentioned in AMAP reports [18,46]. At many conferences related to studies of the Arctic basin, eutrophication of water bodies at high latitudes is not discussed at all, even in the context of climate warming. Eutrophication is undoubtedly hampered by such natural features in the areas as a significant contribution of atmospheric precipitation, the intense flowage of the lakes, and low temperatures. Most lakes in the Arctic zone are oligotrophic (their surface areas are >20 km<sup>2</sup>) and dystrophic (bogged lakes highly enriched in humic acids, their surface areas are usually <20 km<sup>2</sup>). The large lakes (>100 km<sup>2</sup>) are usually thought to be oligotrophic because the effect of allochthonous organic matter on the chemical composition of the natural waters is not as significant.

Northern regions of our planet abound in small oligotrophic lakes that are not directly affected by human activities. Cambell et al. [26] have demonstrated the effects of temperature on the biogeochemical cycles of nutrients in some other lakes. With regard to the classification of lakes in the Kola Peninsula based on their phosphorus concentrations, the number of lakes with low phosphorus concentrations has decreased over the past three decades, whereas the number of lakes corresponding to the meso- and eutrophic types in terms of phosphorus concentration has simultaneously increased [14]. According to [47], practically no oligotrophic lakes are now in America, even lakes not impacted by anthropogenic activities.

In the long period of the polar winter, an oxygen deficiency develops near the bottom because accumulated organic matter is oxidized. Despite the low, anoxia develops by the end of the wintertime in the bottom water layers. This phenomenon can lead to the death of the northern benthic fauna, which is demanding oxygen. In areas where basins in the Arctic are affected by eutrophication, the role of green, cryptophyte, and cyanobacteria in the total phytoplankton biomass increases [15].

## 5. Acidifying Compounds

Human activities in the 20th century, first and foremost, the combustion of fuel-oil residue and fossil oil, coal, as well as smelting operations, resulted in vast atmospheric emissions of acidifying gases (SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>), which converted to acids in the atmosphere and ecosystems and thus lead to the anthropogenic acidification of the waters. The emission levels of anthropogenic sulfur in Europe and North America began to grow rapidly in the early 20th century and reached a maximum in the mid-century [48]. Water acidification had been discernible in acid-sensitive regions in many countries worldwide by the 1960s and continued to intensify from the 1970s until the 1990s. Numerous lakes with low pH of their water and devoid of any living organisms were identified in Scandinavia and the eastern United States. Due to the hazardous character of this phenomenon, the International Cooperative Program on Assessment and Monitoring of Acidification of Rivers and Lakes (ICP-Water) was established in the 1980s to join the efforts of all researchers studying the acidification of waters [49].

Since the 1980s, reducing the emissions of acidifying gases has decreased acid loads onto catchments in central Europe and North America. Surface waters in many acid-vulnerable regions began partly to recover from their acidification in response to the reduction in SO<sub>2</sub> and NO<sub>3</sub> emissions [50–55]. Despite the significant reduction of SO<sub>2</sub> emissions over the past fifty years in North America and Europe, as well as in China starting in 2005 [56], acidic precipitations at catchments still exceed their level during the preindustrial period are continuing to increase (Table 1).

**Table 1.** Predictive scenario of anthropogenic emission of acid-forming substances from natural and anthropogenic sources converted to equivalents of produced acids.

| <i>Acid Equivalent</i>         | 1990 | 2000 | 2025 | 2050 | 2100 |
|--------------------------------|------|------|------|------|------|
| <b>Natural</b>                 |      |      |      |      |      |
| SO <sub>x</sub> , million t-eq | 1, 0 | 1, 0 | 1, 0 | 1, 0 | 1, 0 |
| NO <sub>x</sub> , million t-eq | 0, 4 | 0, 4 | 0, 4 | 0, 4 | 0, 4 |
| <b>Anthropogenic</b>           |      |      |      |      |      |
| SO <sub>x</sub> , million t-eq | 1, 8 | 1, 9 | 2, 9 | 3, 8 | 3, 6 |
| NO <sub>x</sub> , million t-eq | 1, 4 | 1, 5 | 2, 3 | 3, 0 | 4, 1 |
| <b>TOTAL</b>                   | 4, 6 | 4, 8 | 6, 6 | 8, 2 | 9, 1 |

While countries in America and Europe had principally reduced the SO<sub>2</sub> emissions in the 1970s through 1980s, the emissions continued to increase in the Russian Federation and China, and it was not until 1995 that a tendency started to be discernible toward a decrease in the emission of acidifying gases, predominates sulfur dioxide [57–60]. Duan et al. [61] have demonstrated that, despite decades of significant loads of acidic fallouts in Eastern Asia, the problem of water acidification is not as severe because the geochemical structures of the catchments are well capable of neutralizing acidic fallouts. These data are consistent with the calculated critical loads at catchments in China [62].

Acidic precipitation modifies the transport fluxes of chemical elements in the Earth's shells, intensifies the chemical weathering and leaching of major cations and anions, and changes the biogeochemical cycles of elements in the atmosphere and freshwater hydro-sphere, and kills living species susceptible to low pH. The intensity of water acidification is controlled by the following two major factors: (i) the level of airborne anthropogenic load of acidifying compounds on the catchments, with regard to the exposure time, and (ii) the natural susceptibility and vulnerability of the territories to geological, landscape-geographical, and climatic conditions. If the structure of a catchment and its soils are acidic, the waters should be acidified fairly quickly because protons (H<sup>+</sup>) and aluminum ions (Al<sup>3+</sup>) should accompany mobile anions (SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup>) and the water currents formed at the catchments. If the buffer capacity is higher, then the progress of acidification should depend on the volumes of acidifying agents accumulated at the catchments [49,51,52,56]. In the European part of Russia and in West Siberia, acidified lakes were found [13], even in areas devoid of acid loads.

The global and regional reduction in emissions has led to the recovery of acidified waters on all continents. Analysis of literature data led us to distinguish the following three scenarios for the development of long-term changes in the chemistry of waters at a reduction in acidic fallouts [12,47,50–52,54]: (i) the acidification of the waters continues, (ii) neither the pH nor the alkalinity of the waters changes, and (iii) the acid-neutralization capacity (ANC) of the waters is recovered. The uneven character of the recovery tendencies in various countries and regions at similar reductions in the contents of sulfates in the waters is explained by features and circumstances at which the waters are formed at the catchments. The influx of basic cations (BC) from the catchment plays a leading role in forming differences in the recovery of the buffer capacity of the waters.

Similar tendencies were identified in the evolution of some other parameters of water chemistry at a reduction in acidic fallouts: an increase in the content of dissolved organic



carbon (DOC), nitrogen, and phosphorus. Using long-term data on the increase in carbon concentrations, Corman et al. [63] have determined that an increase in DOC content in a water body is associated with a significant increase in the contents of nitrogen and phosphorus in the lake, and this process is associated with a simultaneous increase in TOC. This phenomenon was named brownification and was identified in waters in some areas in North America and Europe [50,64,65]. A number of hypotheses were put forth to explain the increase in organic content matter in the aquatic systems. Montañez et al. [50] believe this process is related to the role of humic acids in recovering the chemical composition of lake waters after their acidification over the past two to three decades. Lately, many researchers have explained this phenomenon by the additional effect of climate warming [14,27,53,66].

## 6. Contamination of the Waters with Toxic Compounds

Toxic contamination of water is one of the most hazardous processes. Toxic properties of chemical elements and compounds are understood as their ability to adversely affect living organisms. It is widely known that many newly synthesized compounds are highly toxic to living organisms. It is also known that industrial byproducts and many naturally occurring elements in atypically high concentrations, for example, the essential metals, such as Cu and Zn, are also toxic to living organisms [67–70]. Aquatic systems are collectors of all toxic elements and compounds known to occur in the environment, and the implications and consequences of this phenomenon are still not fully understood because they may manifest their hazardous properties even if occurring in low concentrations. Coming into aquatic systems, toxic elements and compounds may circulate and be accumulated when moving along the food chains.

All ecologically toxic compounds can be grouped into the following classes:

1. Metals and metalloids, which are elements occurring in nature and toxically affecting living organisms when in high concentrations or certain speciation;
2. Persistent Organic Pollutants (POP<sub>s</sub>), are industrially synthesized compounds or byproducts of some technological processes involving naturally occurring compounds. This group comprises a broad class of organic xenobiotics: herbicides, insecticides, dioxins, furans, phthalates, etc.

Some contaminants are brought to the environment by technological disasters, such as oil spills, massive discharges (and/or leaks) of toxic compounds, and protection-dam breaks at tailing dumps. Natural disastrous events, such as volcanic activity and fires, can also release toxic compounds in concentrations harmful to living organisms; one example is mercury [71].

Metals and metalloids. Metals are brought to water arteries with the runoff from various industrial facilities, diffuse sources, and are leached from rocks by acidic precipitation. Mining operations and smelting facilitate the dispersion of chemical elements. Anthropogenic activities result in the volume of metals coming to water bodies due to anthropogenic processes comparable to the naturally occurring fluxes of these metals (Table 2). The number of publications on the levels of metal concentrations in water, sediments, and biota, as well as their toxic properties (experimental data), is high and continues to increase [72,73].

Two regions north of the Arctic Circle, the Kola Peninsula and the Norilsk area, are the most ecologically alarming regarding water contamination with heavy metals (2020). It has been proved that a broad spectrum of metals (Ni, Cu, Mn, Sr, Fe, Al, Co, Cr, Cd, and Pb) migrates in Arctic water bodies dominantly in the form of ions, which are the most toxic. The toxic properties of the metals are enhanced by the associated eutrophication and acidification of the waters. The acidic atmospheric precipitation intensifies the metals leaching from rocks, further enhancing the hazardousness of the metals in acidic waters [78].

**Table 2.** Natural runoff of metals with rivers and anthropogenic input of elements into the environment and lakes, including metals (according to estimates by various authors: (1)—Bryan [74]; (2)—Moore, Ramamurth [75]; (3)—Venitsianov, Lepikhin [76]; (4)—Heath [77].

| Metals | River Runoff by Chemical Weathering | Inflow to Lakes (According to Our Calculations) | Dispersion Into the Environment             |
|--------|-------------------------------------|---|---|
| Cr     | 60.0 <sup>(3)</sup>                 | 63  | 54–130 <sup>(3)</sup>                       |
| Mn     | 440 <sup>(1)</sup>                  | 2903  | -   |
| Fe     | 25,000 <sup>(1)</sup>               | 31,925  | -   |
| Ni     | 300 <sup>(1)</sup>                  | 161   | 47.4 <sup>(2)</sup> ; 43–98 <sup>(3)</sup>  |
| Cu     | 375 <sup>(1)</sup>                  | 229   | 56 <sup>(2)</sup> ; 56–263 <sup>(3)</sup>   |
| Zn     | 370 <sup>(1)</sup>                  | 693   | 314 <sup>(2)</sup> ; 315–840 <sup>(3)</sup> |
| Mo     | 13 <sup>(1)</sup>                   | 92  | -   |
| Ag     | 5 <sup>(1)</sup>                    | 3.9   | -   |
| Cd     | 4.65 <sup>(3)</sup>                 | 43  | 7–11 <sup>(3)</sup>                         |
| Sn     | 1.5 <sup>(1)</sup>                  | 26  | -   |
| Sb     | 1.3 <sup>(1)</sup>                  | 22.2  | -   |
| Pb     | 180 <sup>(1)</sup>                  | 119   | 449 <sup>(2)</sup> ; 360–440 <sup>(3)</sup> |
| Hg     | 3 <sup>(4)</sup>                    | -   | 5–10 <sup>(3)</sup>                         |

At accompanying eutrophication and the development of oxygen deficit in Arctic regions, the desorption of metals from bottom sediments is responsible for high gradients of the toxic effects in the bottom water layers for the fauna during long wintertime in the Arctic. The mechanism of the redox cycle has been thoroughly studied for Mn and Fe. These elements ascend in the form of dissolved reduced species to the oxic boundary, into oxygen-enriched water layers, where these metals are oxidized again, form low-solubility compounds, and precipitate to the bottom, where the aforementioned cycle is restarted under reduced conditions. Data on the layer-by-layer distribution of elements in the water column of an Arctic lake indicate that a concentration gradient is thus produced not only for Fe and Mn but also for a large group of other elements, such as Cd, Hg, Cu, Mo, Ni, Pb, Zn, Cr, Co, Ba, Ga, and U. A generalized scheme of the metal cycle in the Arctic waters is presented in previous work [9].

The world's scientific community currently pays much attention to such hazardous elements as Hg, Cd, and Pb. In water bodies in the studied areas of the Arctic, Hg concentrations were very low, whereas the accumulation of these elements in various organs and tissues of the fish indicates that the regional waters are ubiquitously contaminated with Hg. The lakes mostly contain Pb (>70%) in the form of its ionic species, which can readily penetrate into the fish organism. This element is accumulated in all systems of the fish organism depending on Pb concentration in the water. Cd is accumulated in all systems of the fish organism, and its highest concentrations were found in the kidneys [79].

Nowadays, environment-protection administrating bodies declare that methods for evaluating the biological availability of elements should be incorporated into procedures for assessing risks of water pollution with metals and into ecological management of water quality [15,80–82]. For example, the Water Framework Directive assumed in the European Community stresses that standards for water quality should include bioavailable Ni and Pb species [83]. The United States Environmental Protection Agency (EPA) has introduced a tool for assessing the bioavailability of Cu in standards for water contamination with this metal [84]. The models rely on determining the amounts of metal ions of the greatest penetrating capacity, i.e., bioavailability. While the standards assumed for Hg in Russia are comparable to those in the West, those for Pb and Cd are much more strict in Europe and the United States.

Persistent organic pollutants (POPs). These are industrially produced compounds that can be retained for years in the environment and be accumulated in fatty tissues. POPs include pesticides, such as aldrin, chlordan, DDT, dieldrin, endrin, heptachlor, hexachlorbenzene, mirex,  $\alpha$ - and  $\beta$ -hexachlorocyclohexane, chlordecone, and lindane, as well as industri-

ally manufactured chemical compounds, such as polychlorobiphenyls, hexachlorobenzene (which is also a pesticide), hexabromdiphenyl, hexa- and- heptabromdiphenyl ether (commercial octabromdiphenyl ether), pentachlorobenzene, perfluorooctane sulphonic acid, and its salts, perfluorooctane sulfonyl fluoride, tetra- and pentabromdiphenyl ether (commercial pentabromdiphenyl ether), and such unintentional byproducts as polychlorinated dioxins and furans [85]. These highly toxic compounds can come to the environment due to various anthropogenic activities, can be transported for long distances by air and water, and can make up hazardous concentrations in waters [73,86,87]. In 2021, the European Commission adopted a proposal to protect human health and the environment from some of the most harmful chemicals in waste: POPs. The convention provided for the possibility of appending the current list of compounds with other ones with the accumulation of necessary information if the newly introduced compounds have principal characteristics of POPs: high stability in the environment, resistance to degradation, acute and chronic toxicity, bioaccumulation, and the ability to be transferred across boundaries for great distances in the environment [86,87]. Table 3 lists the time periods needed for the decomposition of these compounds. These data show how long these compounds can be retained at a catchment and/or in a water body.

**Table 3.** Half-life of some xenobiotics in the environment [88].

| Pollutant         | Half-Life in Temperate Climates |            |                  |              |
|-------------------|---------------------------------|------------|------------------|--------------|
|                   | Air                             | Water      | Bottom Sediments | Soil         |
| DDT               | 2 days                          | >1 year    |                  |              |
| TCDD              | 9 days                          | >5 years   | >1 year          | 10 years     |
| Aldrin            | <9 h                            | <590 days  | no data          | 5 years      |
| Dieldrin          | <40.5 h                         | >2 years   | no data          | >2 years     |
| Endrin            | 1.45 h                          | >112 days  | no data          | <12 years    |
| Chlordane         | <52 h                           | >4 years   | no data          | 1 year       |
| Heptachlor        | no data                         | <1 day     | no data          | 120–240 days |
| Hexachlorobenzene | <4.3 years                      | >100 years | no data          | >3 years     |
| Mirex             | no data                         | >10 h      | >600 years       | >600 years   |
| Toxaphene         | <5 days                         | 20 years   | no data          | 10 years     |
| PCB               | 3–21 days                       | >5 days    | no data          | >40 days     |

The behavior of these compounds in aquatic ecosystems is characterized by the fact that the compounds can be strongly accumulated in aquatic systems because of gradual enrichment in the food chains.

The adverse effects of toxic compounds at their direct or diffuse introduction into lakes and rivers commonly manifest themselves through the chronic effects of low concentrations. In these situations, disturbances are slowly (and often imperceptibly) accumulated in the aquatic organisms and may manifest themselves with the passage of time or in some critical situations, for example, when the temperature is anomalously high, during rain flooding, at stormy weather, etc. [68]. Chronic effects are more difficult to identify in spatiotemporal sections, and their identification requires long-term observations of the organisms, populations (during a few iterations), and communities. Despite great progress in studying the consequence of toxic contaminations, it is still not possible to predict all remote aftermaths of small doses of chronic contamination [10,69,89–91].

## 7. Critical Loads: Theoretical Approaches

Along with other safety types, the ecological one is now one of the principal challenges on this planet because of the complexity, multi-aspect nature, and the importance and extent of the emerging ecological problems. It is hard to suggest either an adequate monitoring program or reasonable measures aimed at data acquisition and processing without preliminary reasonably profound studying of how the quality of waters and their characteristics are formed, which are the mechanisms of the response of organisms and

ecosystems to various anthropogenic impacts. the principle of precluding contamination of aquatic systems instead of fighting the consequences of this contamination should be viewed as one of the guiding lines.


A large team of scientists [2] has analyzed the historical epochs of critical states to identify permissible (threshold) loads for principal anthropogenic loads, i.e., to determine the ability of ecosystems and the biosphere as a whole to deal with anthropogenic fluxes of chemical elements and compounds without causing harm for living organisms. Many scientists believe that threshold concentrations are now exceeded for the increase in CO<sub>2</sub> content, phosphorus fluxes, and other factors. It is however, hard to say how much can be warranted to apply global constraints to individual aquatic systems. Many water bodies on our planet are significantly contaminated with various wastes, whereas vast other areas are not affected by anthropogenic activities at all, such as large territories in Western and Eastern Siberia and the Russian Far East, as well as Antarctica.

Over the past decades, the concept of critical loads was developed as a scientifically justified strategy for precluding the degradation of natural complexes and controlling the state of natural objects. Thereby critical loads are understood as the maximal permissible influxes of one or more contaminants into an ecosystem that does not induce adverse changes in the most vulnerable links of the system (as estimated at the current state of knowledge). When determining the certain permissible (critical) loads in real natural systems, it is necessary to solve complicated fundamental problems of the science of Earth and life, such as:

- (1). behavior patterns of anthropogenically introduced elements, interaction with natural factors, involvement in the natural cycles of elements, and the bioavailability of ecotoxicants and
- (2). stability mechanisms and variability limits of biological systems, the long-term consequences of anthropogenic contaminations for living organisms, and the estimation criteria.

Until these problems are solved, it is not possible to determine how long a natural environment can assimilate the influx of contaminants without adverse consequences for the preservation of survival conditions. A flow chart of an algorithm for the calculation of critical load and management of the anthropogenic impacts is presented in Figure 4. This diagram reflects the convergence of various fields of science aimed at studying ecological processes and real aquatic systems. As demonstrated above, multiple and diverse anthropogenically induced phenomena and processes occur in nature, in which organisms and ecosystems are affected by resultants of all direct and indirect effects of numerous contaminants, climatic factors, and changes in landscapes and biogeochemical cycles.

The international practice offers examples of calculations, which were conducted using methods devised for evaluating critical loads of the necessary levels to which the transfer of some hazardous compounds into terrestrial and aquatic ecosystems should be reduced. The acquired scientifically justified data provided a basis for making international decisions on reducing emissions of the most hazardous contaminants (such as the Gothenburg Protocol on the reduction of emissions of acidifying gases, the Stockholm Convention on Persistent Organic Pollutants, the Minamata Convention on Mercury, etc.). Modern science provides means for calculating the permissible fluxes of nutrients into aquatic systems, and the application of these estimates resulted in the oligotrophication of some lakes and rivers in the recovery of water quality after acidification in some water bodies in Scandinavia and North America. In the example of Lake Urmia (Iran), a strategy of control and recovery after severe desiccation and salinization is shown. The decisions were made based on fundamental studies aimed at determining critical loads.



**Figure 4.** Flowchart of the estimation of critical loads and management of anthropogenic impacts.

A number of important issues were not considered in this paper, such as microplastic pollution [92,93], radioactive waste [94], local industrial pollution, and other anthropogenic influences that manifest themselves in specific water bodies. Understanding their significance for the environment and land waters, we provide links to works in which they are considered.

## 8. Conclusions

The anthropogenic load on surface waters continues to increase despite the significant reduction in atmospheric emissions and discharges of industrial wastewater and agricultural sewage in the current century. Climate warming, in combination with the dispersion of nutrients, ubiquitously results in the eutrophication of waters as a global process. The contents of organic matter and nutrients in surface waters increase, which decreases the number of oligotrophic lakes and leads to the deterioration of water quality at cold latitudes as a consequence of the eutrophication of water bodies at climate warming.

The amounts of anthropogenic-produced sulfates decrease due to the reduction of SO<sub>2</sub> emissions, which enhances the acid-neutralizing capacity of the waters of vulnerable lakes (in terms of the geological structure of their catchments). However, biogeochemical cycles do not return to their original natural parameters, and simultaneously the contents of organic matter and nutrients increase, which modifies the cycling of compounds in the catchment–water body systems.

The migration activity of metals is enhanced; they come to be involved in transport flows, usually in the form of organometallic complexes. The most biologically available and ecologically toxic species of metals are their ionic compounds, except only highly lipophile metals (Hg). The contents of organic xenobiotics in surface waters decrease as a consequence of constraints placed on their use in well-developed countries. Organic ecotoxicants involved in biogeochemical cycles continue to migrate in surface waters as a consequence of the global dispersion of these compounds in the past.

Many lakes and rivers are recovering, and the quality of their water has improved. For example, oligotrophication was detected in some large lakes worldwide, the number of acidified small lakes decreased, and the fluxes of toxic compounds into rivers and lakes were reduced. At the same time, the aquatic systems do not return to their pristine nature parameters, and the ecosystems and biogeochemical cycles reach new states different from the natural ones. The comprehensive and profound understanding of interrelated

transformations of inert and living matter under the effect of human activities provides a basis for the evaluation of permissible constraints aimed at maintaining ecological safety.

**Funding:** This study was supported by Ministry of Education of the Russian Federation (N 0137-2019-0008).

**Conflicts of Interest:** The author declare that they have no conflict of Interest.

## References

- Vernadsky, V.I. *Scientific Ideas as Planetary Phenomenon*; Nauka: Moscow, Russia, 1991; p. 270.
- Rockström, J.; Steffen, W.; Noone, K.; Persson, A.; Chapin, F.S., 3rd; Lambin, E.F.; Lenton, T.M.; Scheffer, M.; Folke, C.; Schellnhuber, H.J.; et al. A safe operating space for humanity. *Nature* **2009**, *461*, 472–475. [[CrossRef](#)] [[PubMed](#)]
- Zhang, X.; Davidson, E.A.; Zou, T.; Lassaletta, L.; Quan, Z.; Li, T.; Zhang, W. Quantifying nutrient budgets for sustainable nutrient management. *Glob. Biogeochem. Cycles* **2020**, *34*, e2018GB006060. [[CrossRef](#)]
- Danilov-Danilyan, V.I.; Khranovich, I.L. *Water Resources Management. Coordination of Water Use Strategies*; Nauchny mir: Moscow, Russia, 2010; p. 233.
- AR5 Synthesis Report: Climate Change. 2014. Available online: <https://www.ipcc.ch/report/ar5/syr/Ar5/ar5.html> (accessed on 19 January 2022).
- AR6 Climate Change 2022: Mitigation of Climate Change—IPCC. Sixth Assessment Report. 2022. Available online: <https://www.ipcc.ch/report/sixth-assessment-report> (accessed on 9 July 2021).
- Cesonien, E.L.; Šileikienė, D.; Marozas, V.; Čiteikė, L. Influence of Anthropogenic Loads on Surface Water Status: A Case Study in Lithuania. *Sustainability* **2021**, *13*, 4341. [[CrossRef](#)]
- Biswas, A.K. *Water Security Under Climate Change*; Asit, K., Ed.; Springer Nature Singapore Pte Ltd.: Singapore, 2022; p. 830. [[CrossRef](#)]
- Moiseenko, T.I. A Fate of Metals in Arctic Surface Waters. Method for Defining Critical Levels. *Sci. Total Environ.* **1999**, *236*, 19–39. [[CrossRef](#)]
- Moiseenko, T.I. Aquatic ecotoxicology: Theoretical principles and practical application. *Water Resour.* **2008**, *35*, 530–541. [[CrossRef](#)]
- Moiseenko, T.I.; Skjelkvåle, B.L.; Gashkina, N.A.; Shalabodov, A.D.; Khoroshavin, V.Y. Water chemistry in small lakes along a transect from boreal to arid ecoregions in European Russia: Effects of air pollution and climate change. *Appl. Geochem.* **2013**, *28*, 69–79. [[CrossRef](#)]
- Moiseenko, T.I.; Dinu, M.I.; Bazova, M.M.; de Wit, H.A. Long-term changes in the water chemistry of subarctic lakes as a response to reduction of air pollution: Case study in the Kola North, Russia. *Water Air Soil Pollut.* **2015**, *226*, 1–12. [[CrossRef](#)]
- Moiseenko, T.I.; Dinu, M.I.; Gashkina, N.A.; Jones, V.; Khoroshavin, V.Y.; Kremleva, T.A. Present status of water chemistry and acidification under nonpoint sources of pollution across European Russia and West Siberia. *Environ. Res. Lett.* **2018**, *13*, 105007. [[CrossRef](#)]
- Moiseenko, T.I.; Bazova, M.M.; Dinu, M.I.; Gashkina, N.A.; Kudryavtseva, L.P. Changes in the Geochemistry of Land Waters at Climate Warming and a Decrease in Acid Deposition: Recovery of the Lakes or Their Evolution? *Geochem. Int.* **2022**, *60*, 685–701. [[CrossRef](#)]
- Moiseenko, T.; Sharov, A. Large Russian lakes Ladoga, Onega, and Imandra under strong pollution and in the period of revitalization: A review. *Geosciences* **2019**, *9*, 492. [[CrossRef](#)]
- Edelstein, K.K. *Continental Hydrology*; Publishing Center “Academy”: Moscow, Russia, 2011; p. 304.
- UN World Water Development Report: Valuing Water. 2021. Available online: <https://www.unwater.org/publications/un-world-water-development-report-2021> (accessed on 27 September 2022).
- Arctic Freshwater System in a Changing Climate, WCRP CliC Project. AMAP. IASC CliC/AMAP/IASC. 2016. Available online: [http://www.amap.no/documents/doc/The-Arctic-Freshwater-System-in-a-Changing-Climate/1375.AMA\\$%20Pi\\$](http://www.amap.no/documents/doc/The-Arctic-Freshwater-System-in-a-Changing-Climate/1375.AMA$%20Pi$) (accessed on 1 January 2020).
- Hayhoe, K.; Wake, C.P.; Huntington, T.G.; Luo, L.; Schwartz, M.D.; Sheffield, J.; Wood, E.; Anderson, B.; Bradbury, J.; DeGaetano, A. Past and future changes in climate and hydrological indicators in the US Northeast. *Clim. Dyn.* **2007**, *28*, 381–407. [[CrossRef](#)]
- Georgiadi, A.G.; Kashutina, E.A. Long-term runoff changes of the largest siberian rivers. *Izv. Ross. Akad. Nauk. Seriya Geogr.* **2016**, *5*, 70–81. (In Russian) [[CrossRef](#)]
- Jin, H.; Huang, Y.; Bense, V.F.; Ma, Q.; Marchenko, S.S.; Shepelev, V.V.; Hu, Y.; Liang, S.; Spektor, V.V.; Jin, X.; et al. Permafrost Degradation and Its Hydrogeological Impacts. *Water* **2022**, *14*, 372. [[CrossRef](#)]
- Pavlov, A.V.; Malkova, G.D. Small-scale mapping of trends of the contemporary ground temperature changes in the Russian north. *Earth Cryosphere* **2009**, *103*, 32–39.
- Grosse, G.; Goetz, S.; McGuire, D.; Romanovsky, V.E.; Schuur, E.A. Changing permafrost in a warming world and feedbacks to the Earth system. *Environ. Res. Lett.* **2016**, *11*, 040201. [[CrossRef](#)]
- Gao, X.; Schlosser, C.A.; Sokolov, A.; Anthony, K.W.; Zhuang, Q.; Kicklighter, D. Permafrost degradation and methane: Low risk of biogeochemical climate-warming feedback. *Environ. Res. Lett.* **2013**, *8*, 035014. [[CrossRef](#)]

25. Meingast, K.M.; Kane, E.; Coble, A.A.; Marcarelli, A.M.; Toczydlowski, D. Climate, snowmelt dynamics and atmospheric deposition interact to control dissolved organic carbon export from a northern forest stream over 26 years. *Environ. Res. Lett.* **2020**, *15*, 104034. [[CrossRef](#)]
26. Campbell, J.L.; Rustad, L.E.; Boyer, E.W.; Christopher, S.F.; Driscoll, C.T.; Fernandez, I.J.; Groffman, P.M.; Houle, D.; Kiebusch, J.; Magill, A.H.; et al. Consequences of climate change for biogeochemical cycling in forests of northeastern North America. *Can. J. For. Res.* **2009**, *39*, 264–284.
27. De Wit, H.A.; Stoddard, J.L.; Monteith, D.T.; Sample, J.E.; Austnes, K.; Couture, S.; Fölster, J.; Higgins, S.N.; Houle, D.; Hruška, J.; et al. Cleaner air reveals growing influence of climate on dissolved organic carbon trends in northern headwaters. *Environ. Res. Lett.* **2021**, *16*, 104009. [[CrossRef](#)]
28. Watmough, S.A.; Eimers, C.; Baker, S. Impediments to recovery from acid deposition. *Atmos. Environ.* **2016**, *146*, 15–27.
29. Gavin, A.L.; Nelson, S.J.; Klemmer, A.J.; Fernandez, I.J.; Strock, K.E.; McDowell, W.H. Acidification and climate linkages to increased dissolved organic carbon in high elevation lakes. *Water Resour. Res.* **2018**, *54*, 5187–5877. [[CrossRef](#)]
30. Houle, D.; Couture, S.; Gagnon, C. Relative role of decreasing precipitation sulfate and climate on recent lake recovery. *Glob. Biogeochem. Cycles* **2010**, *24*, 4029. [[CrossRef](#)]
31. Corman, J.R.; Bertolo, B.L.; Casson, N.J.; Sebestyen, S.D.; Kolka, R.K.; Stanley, E.H. Nitrogen and phosphorus loads to temperate seepage lakes associated with allochthonous dissolved organic carbon loads. *Geophys. Res. Lett.* **2018**, *45*, 5481–5490. [[CrossRef](#)]
32. Jellison, R.; Williams, W.D.; Timms, B.; Aladin, N.V. Salt lakes: Values and future. In *Aquatic Ecosystems-Trends and Global Prospects*; Cambridge University Press: Cambridge, UK, 2008; pp. 92–110.
33. Pouladi, P.; Mohammad, A.; Afshar, H.; Molajou, A.; Farahmand, H. Agent-based socio-hydrological modeling for restoration of Urmia Lake: Application of theory of planned behavior. *J. Hydrol.* **2019**, *576*, 736–748. [[CrossRef](#)]
34. Tussupova, K.; Hjorth, A.P.; Morave, M. Drying lakes: A review on the applied restoration strategies and health conditions in contiguous areas. *Water* **2020**, *12*, 749. [[CrossRef](#)]
35. Rudneva, I.I.; Shaidaa, V.G.; Shcherba, A.V.; Zavyalova, A.V. Influence of Climatic Factors on Interannual and Seasonal Dynamics of the Environmental State of the Salt Lake Adzhi-Baichi (Crimea). *Arid. Ecosys-Tems* **2021**, *11*, 434–442. [[CrossRef](#)]
36. Mojtahedi, A.; Dadashzadeh, M.; Azizkhani, M.; Mohammadian, A.; Almasi, R. Assessing climate and human activity effects on lake characteristics using spatio-temporal satellite data and an emotional neural network. *Environ. Earth Sci.* **2022**, *81*, 1–20. [[CrossRef](#)]
37. Mekonnen, M.M.; Hoekstra, A.Y. Global anthropogenic phosphorus loads to freshwater and associated grey water footprints and water pollution levels: A high-resolution global study. *Water Resour. Res.* **2018**, *54*, 345–358. [[CrossRef](#)]
38. Penuelas, J.; Poulter, B.; Sardans, J.; Ciais, P.; Van Der Velde, M.; Bopp, L. Human-induced nitrogen–phosphorus imbalances alter natural and managed ecosystems across the globe. *Nat. Commun.* **2013**, *4*, 2934. [[CrossRef](#)]
39. Davidson, E.A.; Markewitz, D.; de Figueiredo, R.O.; de Camargo, P.B. Nitrogen fixation inputs in pasture and early successional forest in the Brazilian amazon region: Evidence from a claybox mesocosm study. *J. Geophys. Res. Biogeosciences* **2018**, *123*, 712–721. [[CrossRef](#)]
40. Bouwman, A.; Beusen, A.; Griffioen, J.; Van Groenigen, J.; Hefting, M.; Oenema, O.; Van Puijenbroek, P.J.T.M.; Seitzinger, S.; Slomp, C.P.; Stehfest, E. Global trends and uncertainties in terrestrial denitrification and N<sub>2</sub>O emissions. *Philos. Trans. R. Soc. B Biol. Sci.* **2013**, *368*, 20130112. [[CrossRef](#)] [[PubMed](#)]
41. Bouwman, A.F.; Beusen, A.H.; Lassaletta, L.; van Apeldoorn, D.F.; van Grinsven, H.J.; Zhang, J.; van Ittersum, M.K. Lessons from temporal and spatial patterns in global use of N and P fertilizer on cropland. *Sci. Rep.* **2017**, *7*, 40366. [[CrossRef](#)] [[PubMed](#)]
42. Gruber, N.; Galloway, J.N. An Earth-system perspective of the global nitrogen cycle. *Nature* **2008**, *451*, 293–296. [[CrossRef](#)] [[PubMed](#)]
43. Powers, S.M.; Bruulsema, T.W.; Burt, T.P.; Chan, N.I.; Elser, J.J.; Haygarth, P.M. Long-term accumulation and transport of anthropogenic phosphorus in three river basins. *Nat. Geosci.* **2016**, *9*, 353–356. [[CrossRef](#)]
44. Abid, A.A.; Gill, S.S. *Eutrophication: Causes, Consequences and Control*; Springer: Dordrecht, The Netherlands; Heidelberg, Germany; London, UK, 2014; Volume 2, p. 262.
45. Beusen, A.H.; Bouwman, A.F.; Van Beek, L.P.; Mogollón, J.M.; Middelburg, J.J. Global riverine N and P transport to ocean increased during the 20th century despite increased retention along the aquatic continuum. *Biogeosciences* **2016**, *13*, 2441–2451. [[CrossRef](#)]
46. Nilsson, A.E.; Wilson, S.J.; Murray, J.L.; Huntington, H.P. *Arctic Pollution Issues: A State of the Arctic Environment Report*; Publication by AMAP: Oslo, Norway, 1997; p. 188.
47. Stoddard, J.L.; Van Sickle, J.; Herlihy, A.T.; Brahney, J.; Paulsen, S.; Peck, D.V. Continental-scale increase in lake and stream phosphorus: Are oligotrophic systems disappearing in the United States? *Environ. Sci. Technol.* **2016**, *50*, 3409–3415. [[CrossRef](#)]
48. Galloway, J.N. Acid deposition: Perspectives in time and space. *Water Air Soil Pollut.* **1995**, *85*, 15–24. [[CrossRef](#)]
49. Kvaeven, B.; Ulstein, M.J.; Skjelkvåle, B.L. ICP Waters—An international program for surface water monitoring. *Water Air Soil Pollut.* **2001**, *130*, 775–780. [[CrossRef](#)]
50. Monteith, D.T.; Stoddard, J.L.; Evans, C.D.; de Wit, H.A.; Forsius, M.; Hogasen, T.; Wilander, A.; Skjelkvåle, B.L.; Jeffries, D.S.; Vuorenmaa, J.; et al. Dissolved organic carbon trends resulting from changes in atmospheric deposition chemistry. *Nature* **2007**, *450*, 537–539. [[CrossRef](#)]

51. Skelkvale, B.L.; Stoddard, J.L.; Andersen, T. Trends in surface water acidification in Europe and North America (1989–1998). *Water Air Soil Pollut.* **2001**, *130*, 787–792. [[CrossRef](#)]
52. Skjelkvale, B.L.; Stoddard, J.L.; Jeffries, D.S.; Torseth, K.; Hogasen, T.J. Regional scale evidence for improvements in surface water chemistry 1990–2001. *Environ. Pollut.* **2005**, *137*, 165–176. [[CrossRef](#)] [[PubMed](#)]
53. Clair, T.A.; Dennis, I.F.; Vet, R. Water chemistry and dissolved organic carbon trends in lakes from Canada’s Atlantic Provinces: No recovery from acidification measured after 25 years of lake monitoring. *J. Fish. Aquat. Sci.* **2011**, *68*, 663–674. [[CrossRef](#)]
54. Strock, K.E.; Nelson, S.J.; Kahl, J.S.; Saros, J.E.; Mc Dowell, W.H. Decadal trends reveal recent acceleration in the rate of recovery from acidification in the northeastern U.S. *Environ. Sci. Technol.* **2014**, *48*, 4681–4689. [[CrossRef](#)] [[PubMed](#)]
55. Rogora, M.; Colombo, L.; Marchetto, A.; Mosello, R.; Steingruber, S. Temporal and spatial patterns in the chemistry of wet deposition in Southern Alps. *Atmos. Environ.* **2016**, *146*, 44–54. [[CrossRef](#)]
56. Garmo, O.G.; Skjelkvåle, B.L.; de Wit, H.A.; Colombo, L.; Curtis, C.; Folster, J.; Hoffmann, A. Trends in surface water chemistry in acidified areas in Europe and North America from 1990 to 2008. *Water Air Soil Pollut.* **2014**, *225*, 1880. [[CrossRef](#)]
57. Graedel, T.E.; Benkovitz, C.M.; Keene, W.C.; Lee, D.S.; Marland, G. Global emission inventories of acid-related compounds. *Water Air Soil Pollut.* **1995**, *85*, 25–36. [[CrossRef](#)]
58. Zhao, B.; Shuxiao, W.; Xinyi, D.; Jiandong, W.; Lei, D.; Xiao, F.; Jiming, H.; Joshua, F. Environmental effects of the recent emission changes in China: Implications for particulate matter pollution and soil acidification. *Environ. Res. Lett.* **2013**, *8*, 024031. [[CrossRef](#)]
59. Wang, S.X.; Zhao, B.; Cai, S.Y.; Klimont, Z.; Nielsen, C.P.; Morikawa, T.; Woo, J.H.; Kim, Y.; Fu, X.; Xu, J.Y.; et al. Emission trends and mitigation options for air pollutants in East Asia. *Atmos. Chem. Phys.* **2014**, *14*, 6571–6603. [[CrossRef](#)]
60. Meng, C.; Dashuan, T.; Hui, Z.; Zhaolei, L.; Chuixiang, Y.; Shuli, N. Global soil acidification impacts on belowground processes. *Environ. Res. Lett.* **2019**, *14*, 074003. [[CrossRef](#)]
61. Duan, L.; Ma, X.X.; Larssen, T.; Mulder, J.; Hao, J.M. Response of surface water acidification in upper Yangtze River to SO<sub>2</sub> emissions abatement in China. *Environ. Sci. Technol.* **2011**, *45*, 3275–3281. [[CrossRef](#)]
62. Yu, Q.; Zhang, T.; Cheng, Z.; Zhao, B.; Mulder, J.; Larssen, T.; Wang, S.; Duan, L. Is surface water acidification a serious regional issue in China? *Sci. Total Environ.* **2017**, *15*, 584–585. [[CrossRef](#)] [[PubMed](#)]
63. Hanson, P.C.; Pace, M.L.; Carpenter, S.R.; Cole, J.J.; Stanley, E.H. Integrating landscape carbon cycling: Research needs for resolving organic carbon budgets of lakes. *Ecosystems* **2015**, *18*, 363–375. [[CrossRef](#)]
64. Clark, J.M.; Bottrell, S.H.; Evans, C.D.; Monteith, D.T.; Bartlett, R.; Rose, R.; Newton, R.J.; Chapman, P.J. The importance of the relationship between scale and process in understanding long-term DOC dynamics. *Sci. Total Environ.* **2010**, *408*, 2768. [[CrossRef](#)] [[PubMed](#)]
65. De Wit, H.A.; Valinia, S.; Weyhenmeyer, G.A.; Futter, M.N.; Kortelainen, P.; Austnes, K.; Hessen, D.O.; Räike, A.; Laudon, H.; Vuorenmaa, J. Current browning of surface waters will be further promoted by wetter climate. *Environ. Sci. Technol. Lett.* **2016**, *3*, 430–435. [[CrossRef](#)]
66. Strock, K.E.; Theodore, N.; Gawley, W.G.; Ellsworth, A.C.; Saros, J.E. Increasing dissolved organic carbon concentrations in northern boreal lakes: Implications for lake water transparency and thermal structure. *J. Geophys. Res. Biogeosci.* **2017**, *122*, 1022–1035. [[CrossRef](#)]
67. Chapman, P.M. Integrating toxicology and ecology: Putting the “eco” into ecotoxicology. *Mar. Pollut. Bull.* **2002**, *44*, 7–15. [[CrossRef](#)]
68. Newman, M.C.; Clements, W.H. *Aquatic Toxicology: Concepts, Practice, New Directions*. In *General, Applied and Systems Toxicology*; John Wiley & Sons, Ltd.: Hoboken, NJ, USA, 2009; Available online: <https://onlinelibrary.wiley.com/doi/book/10.1002/9780470744307> (accessed on 27 September 2022).
69. Boudou, A. *Aquatic Ecotoxicology. Volume 1: Fundamental Concepts and Methodologies*; Taylor & Francis eBooks: College Park, MD, USA, 2018; p. 328. Available online: <https://www.routledge.com/Aquatic-Ecotoxicology-Volume-1-Fundamental-Concepts-and-Methodologies/Boudou/p/book/9781315890753> (accessed on 27 September 2022).
70. Vasseur, P.; Masfaraud, J.-F.; Blaise, C. Ecotoxicology, revisiting its pioneers. *Environ. Sci. Pollut. Res. Int.* **2021**, *28*, 3852–3857. [[CrossRef](#)]
71. Fitzgerald, L.; Wikoff, D.S. Persistent organic pollutants. In *Encyclopedia of Toxicology*, 3rd ed.; Wexler, P., Ed.; Elsevier Inc.: Amsterdam, The Netherlands; Academic Press: Cambridge, MA, USA, 2014; Volume 3, pp. 820–825. Available online: <https://www.sciencedirect.com/topics/agricultural-and-biological-sciences/persistent-organic-pollutant> (accessed on 27 September 2022).
72. Davidson, T.; Costa, M. Selected Molecular Mechanisms of Metal Toxicity and Carcinogenicity. In *Handbook on the Toxicology of Metals*; Nordberg, G.F., Fowler, B.A., Nordberg, M., Eds.; Academic Press: Cambridge, MA, USA, 2014; pp. 79–100.
73. Bjerregaard, P.; Andersen, O. Ecotoxicology of Metals—Sources, Transport, and Effects in the Ecosystem. In *Handbook on the Toxicology of Metals*; Nordberg, G.F., Fowler, B.A., Nordberg, M., Eds.; Elsevier: Amsterdam, The Netherlands, 2014.
74. Bryan, G.W. *Heavy Metal Contamination in the Sea, Marine Pollution*; Johnston, R., Ed.; Academic Press: Cambridge, MA, USA; Elsevier: London, UK, 1976; pp. 185–302.
75. Moore, J.W.; Ramamurthy, S. *Heavy Metals in Natural Waters: Applied Monitoring and Impact Assessment*; Springer: Berlin/Heidelberg, Germany, 1984; p. 288.
76. Venitsianov, E.V.; Lepikhin, A.P. *Physico-Chemical Bases of Modeling Migration and Transformation of Heavy Metals in Natural Waters*; Publishing House of RosNIIVH: Yekaterinburg, Russia, 2002; p. 236.
77. Heath, A.G. *Water Pollution and Fish Physiology*; Lewis Publishers: Amsterdam, The Netherlands, 2002; p. 506.



78. Moiseenko, T.I.; Gashkina, N.A.; Dinu, M.I.; Kremleva, T.A.; Khoroshavin, V.Y. Water Chemistry of Arctic Lakes under Airborne Contamination of Watersheds. *Water* **2020**, *12*, 1659. [[CrossRef](#)]
79. Moiseenko, T.I.; Gashkina, N.A. Distribution and bioaccumulation of heavy metals (Hg, Cd, and Pb) In fish: Influence of the aquatic environment and climate. *Environ. Res. Lett.* **2020**, *15*, 115013. [[CrossRef](#)]
80. Magalhaes, D.; Marques, M.; Baptista, D.; Forsin, D.; Buss, D. Metal bioavailability and toxicity in freshwaters. *Environ. Chem. Lett.* **2015**, *13*, 69–87. [[CrossRef](#)]
81. Merrington, G.; Peters, A.; Schlegel, C.E. Accounting for metal bioavailability in assessing water quality: A step change? *Environ. Toxicol. Chem.* **2016**, *35*, 257–265. [[CrossRef](#)] [[PubMed](#)]
82. Väänänen, K.; Leppänen, M.T.; Chen, X.; Akkanen, J. Metal bioavailability in ecological risk assessment of freshwater ecosystems: From science to environmental management. *Ecotoxicol. Environ. Saf.* **2018**, *147*, 430–446. [[CrossRef](#)] [[PubMed](#)]
83. European Commission. *Directive 2013/39/EC Amending Directives 2000/60/EC and 2008/105/EC as Regards Priority Substances in the Field of Water Policy*. 2013/39/EC; European Commission: Brussels, Belgium, 2013; Available online: <https://www.eea.europa.eu/policy-documents/2013-39-ec> (accessed on 27 September 2022).
84. USA EPA. *National Recommended Water Quality Criteria. Report 4304T. Office of Water, Office of Science and Technology*. (EPA/600/4-91/002); USA EPA: Springfield, IL, USA, 2007.
85. Stockholm Convention on Persistent Organic Pollutants. UN Convention; 2004. Available online: <https://www.state.gov/key-topics-office-of-environmental-quality-and-transboundary-issues/stockholm-convention-on-persistent-organic-pollutants> (accessed on 2 April 2022).
86. Lewis, K.A.; Tzilivakis, J.; Warner, D.; Green, A. An international database for pesticide risk assessments and management. *Hum. Ecol. Risk Assess. Int. J.* **2016**, *22*, 1050–1064. [[CrossRef](#)]
87. Barrick, A.; Champeau, O.; Butler, J.; Wiles, T.; Boundy, M.; Tremblay, L.A. Hazard/Risk Assessment Investigating the Ecotoxicity of Select Emerging Organic Contaminants Toward the Marine Copepod *Gladioferens pectinatus*. *Environ. Toxicol. Chem.* **2022**, *41*, 792–799. [[CrossRef](#)] [[PubMed](#)]
88. Khudoley, V.V. *Stockholm Convention, and National Plan of Action Against Persistent Organic Pollutants (POPs)*; The International Ecological Forum: St.Petersburg, Russia, 2002; pp. 89–91.
89. Zhulidov, A.V.; Robarts, R.D.; Headley, J.V.; Korotova, L.G.; Pavlov, D.F.; Zhulidov, D.A.; Zhulidova, O.V. A review riverine fluxes of hexachlorocyclohexane and DDT to Azov and Black seas from the former USSR and Russian Federation. *J. Environ. Sci. Health* **2003**, *5*, 38.
90. Blus, L.I. Organochlorine Pesticides. In *Handbook of Ecotoxicology*; Hoffman, D.J., Rattner, B.A., Burton, G.A., Cairnce, J., Jr., Eds.; Lewis Publishers: New York, NY, USA, 2005; pp. 314–329.
91. Rice, C.P.; O’Keefe, P.; Kubiak, T.J. Sources, Partways and effects of PCBs, Dioxins and Dibenzofurans. In *Handbook of Ecotoxicology*; Hoffman, D.J., Rattner, B.A., Burton, G.A., Cairnce, J., Jr., Eds.; Lewis Publishers: New York, NY, USA, 2005; pp. 501–556.
92. Ding, J.; Zhang, S.; Razanajatovo, R.M. Accumulation, tissue distribution, and biochemical effects of polystyrene microplastics in the freshwater fish red tilapia (*Oreochromis niloticus*). *Environ. Pollut.* **2018**, *238*, 1–9. [[CrossRef](#)]
93. Wang, W.; Gao, H.; Jin, S.; Li, R.; Na, G. The ecotoxicological effects of microplastics on aquatic food web, from primary producer to human: A review. *Ecotoxicol. Environ. Saf.* **2019**, *173*, 110–117. [[CrossRef](#)]
94. Gianni, P. *2020 Nuclear Safety*, 2nd ed.; Elsevier Ltd.: Amsterdam, The Netherlands, 2020.