

Supplementary Material

For

Thyroid-disrupting effects of cadmium and mercury in zebrafish embryos/larvae

Liqiao Zhong ^{1,†}, He Zhang ^{2,†}, Luyin Wu ¹, Huijun Ru ¹, Nian Wei ¹, Fan Yao ¹, Zhaohui Ni ¹, Xinbin Duan ^{1,*} and Yunfeng Li ^{1,*}

¹ Fishery Resources and Environmental Science Experimental Station of the Upper-Middle Reaches of Yangtze River (Ministry of Agriculture and Rural Affairs), Yangtze River Fisheries Research Institute, Chinese Academy of Fishery Sciences, Wuhan 430223, China

² Zhejiang Provincial Key Lab for Subtropical Water Environment and Marine Biological Resources Protection, College of Life and Environmental Science, Wenzhou University, Wenzhou 325035, China

* Correspondence: duan@yfi.ac.cn (X.D.); lyf086@yfi.ac.cn (Y.L.);
Tel.: +86-27-81780131 (X.D.); +86-27-81780050 (Y.L.);
Fax: +86-27-81780088 (Y.L.)

† These authors contributed equally to this work.




Figure S1. Morphology of zebrafish larvae exposure to Cd^{2+} and Hg^{2+} . Normal zebrafish larvae (A); the morphological changes caused by Cd^{2+} (B–D), including yolk sac edema (B), tail malformation (C), and yolk sac edema and pericardial edema (D); the morphological changes caused by Hg^{2+} (E–I), including tail malformation (E), yolk sac edema (I), pericardial edema (H) and spinal curvature (I).