

Photocatalytic degradation of Sulfamethazine using decatungstate anions in aqueous solution: Mechanistic approach

Mohammed-Amine Edaala ^{1,*}, Lekbira El Mersly ¹, Abdelaziz Aloui Tahiri ¹; Pascal Wong-Wah-Chung ²; Lahssen El blidi³, Maher M. Alrashed³ and Salah Rafqah¹

¹ Laboratoire de Chimie Analytique et Moléculaire, Faculté polydisciplinaire de Safi, Université Cadi Ayyad, Morocco.;

mohammedamine.edaala@gmail.com, s.rafqah@uca.ma, kabira.elmersly@gmail.com, az_alaoui@yahoo.fr

² Aix Marseille Université, CNRS, LCE, Marseille, France, pascal.wong-wah-chung@univ-amu.fr

³ Chemical Engineering Department, College of Engineering, King Saud University, P.O. Box 800, Riyadh 11421, Saudi Arabia, lelblidi@ksu.edu.sa, mabdulaziz@ksu.edu.sa

* Correspondence: mohammedamine.edaala@gmail.com ; Tel.: +212651-345751

Supplementary material




Figure S1: Molecular structure of SMZ






Figure S2 : Absorption spectrum of the Sulfamethazine ([SMZ] =27.8 mg/L, pH=4 in water)






Figure S5: CID spectrum of P2 (CE = 20 eV)




Figure S6: CID spectrum of P3 (CE = 20 eV)

