

Arable field species (FIE) ¹ ; df: 5, 73				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.388	6.456	7.289	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	-0.099	0.033	-3.037	0.003
Herb [%]	-0.084	0.131	-0.638	0.525
Grass [%]	0.075	0.045	1.682	0.097
Litter [%]	-0.097	0.045	-2.183	0.032
Open ground [%]	0.039	0.044	0.876	0.384
Grassland species (GRL); df: 5, 74				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.422	5.297	16.819	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	-0.094	0.024	-3.960	<0.001
Herb [%]	0.077	0.121	0.636	0.527
Grass [%]	-0.034	0.074	-0.466	0.643
Litter [%]	0.068	0.056	1.217	0.228
Open ground [%]	0.134	0.055	2.423	0.018
Ruderal species (RUD) ² ; df: 5, 73				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.375	3.167	35.332	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	0.091	0.015	6.170	<0.001
Herb [%]	0.076	0.042	1.808	0.075
Grass [%]	-0.010	0.019	-0.530	0.598
Litter [%]	-0.012	0.027	-0.431	0.668
Open ground [%]	-0.051	0.028	-1.815	0.074
Forest species, in general (FOR); df: 5, 74				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.506	19.358	1.967	0.094
Coefficients _{HAC}	B	SE	t	p
Shade [%]	0.061	0.099	0.611	0.543
Herb [%]	0.783	0.284	2.754	0.007
Grass [%]	0.732	0.288	2.544	0.013
Litter [%]	0.724	0.251	2.886	0.005
Open ground [%]	0.835	0.309	2.705	0.009
Group of species with different habitat preferences (MIX); df: 5, 74				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.213	5.224	7.652	0.008
Coefficients _{HAC}	B	SE	t	p
Shade [%]	-0.037	0.039	-0.950	0.345
Herb [%]	-0.150	0.107	-1.404	0.165
Grass [%]	-0.120	0.161	-0.743	0.460
Litter [%]	-0.086	0.083	-1.034	0.305
Open ground [%]	0.063	0.060	1.060	0.293

1) After elimination of the outlier in SRC3 trap4 2011, outlier test +/- 3 SD

2) After elimination of the outlier in SRC1 trap3 2013. outlier test +/- 3 SD