

Article

New Hermite–Hadamard Type Inequalities in Connection with Interval-Valued Generalized Harmonically (h_1, h_2) -Godunova–Levin Functions

Soubhagya Kumar Sahoo ^{1,2}, Pshtiwan Othman Mohammed ³, Donal O' Regan ⁴, Muhammad Tariq ⁵
and Kamsing Nonlaopon ^{6,*}

- ¹ Department of Mathematics, Institute of Technical Education and Research, Siksha 'O' Anusandhan University, Bhubaneswar 751030, India
² Department of Mathematics, Aryan Institute of Engineering and Technology, Bhubaneswar 752050, India
³ Department of Mathematics, College of Education, University of Sulaimani, Sulaimani 46001, Iraq
⁴ School of Mathematical and Statistical Sciences, National University of Ireland, H91 TK33 Galway, Ireland
⁵ Department of Basic Sciences and Related Studies, Mehran University of Engineering and Technology, Jamshoro 76062, Pakistan
⁶ Department of Mathematics, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand
* Correspondence: nkamsi@kku.ac.th; Tel.: +66-0866421582

Abstract: As is known, integral inequalities related to convexity have a close relationship with symmetry. In this paper, we introduce a new notion of interval-valued harmonically (m, h_1, h_2) -Godunova–Levin functions, and we establish some new Hermite–Hadamard inequalities. Moreover, we show how this new notion of interval-valued convexity has a close relationship with many existing definitions in the literature. As a result, our theory generalizes many published results. Several interesting examples are provided to illustrate our results.

Keywords: interval-valued analysis; Hermite–Hadamard type inequalities; Godunova–Levin functions; harmonically convex functions

MSC: 26A51; 26A33; 26D10; 26D15



Citation: Sahoo, S.K.; Mohammed, P.O.; O'Regan, D.; Tariq, M.; Nonlaopon, K. New Hermite–Hadamard Type Inequalities in Connection with Interval-Valued Generalized Harmonically (h_1, h_2) -Godunova–Levin Functions. *Symmetry* **2022**, *14*, 1964. <https://doi.org/10.3390/sym14101964>

Academic Editors: Natanael Karjanto and Angelo Favini

Received: 23 August 2022

Accepted: 14 September 2022

Published: 20 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Interval analysis is a subset of set-valued analysis, which is the study of sets in the context of mathematics and general topology. The Archimedean approach, which includes determining a circle's circumference, is a classic illustration of interval enclosure. This theory addresses the interval uncertainty that exists in many computational and mathematical models of deterministic real-world systems. With this approach, errors that result in incorrect conclusions are avoided by studying interval variables instead of point variables and expressing computation results as intervals. Consideration of the error estimates of the numerical solutions for finite state machines was one of the initial goals of the interval-valued analysis. Interval analysis, which Moore first described in his renowned book [1], is one of the fundamental techniques in numerical analysis. As a result, it has found applications in many fields, including differential equations for intervals [2], neural network output optimization [3], automatic error analysis [4], computer graphics [5], and many more. For results and applications, we refer interested readers to [6–10].

Inequalities have a significant impact on mathematics, particularly those connected to the Jensen, Ostrowski, Hermite–Hadamard, Bullen, Simpson, and Opial inequalities. Many of these inequalities have recently been extended to interval-valued functions by some well-known researchers (see, for example, [11–14]), and many have also researched

the Hermite–Hadamard inequality for convex functions. The traditional H–H inequality is given as:

$$\mathcal{G}\left(\frac{c+d}{2}\right) \leq \frac{1}{d-c} \int_c^d \mathcal{G}(x) dx \leq \frac{\mathcal{G}(c) + \mathcal{G}(d)}{2},$$

where \mathcal{G} is a convex function.

On the other hand, the generalized convexity of mappings is a potent tool for addressing a broad range of issues in applied analysis and nonlinear analysis, including various problems in mathematical physics. Recently, a number of generalizations of convex functions have been thoroughly researched. Mathematical analytic study on the idea of integral inequalities is interesting. The study of differential and integral equations has also been considered to be relevant for inequalities and various extended convex mappings. Electrical networks, symmetry analysis, operations research, finance, decision making, numerical analysis, and equilibrium are just a few areas where they have had a substantial impact. We investigate how the subjective properties of convexity might be encouraged by using a number of fundamental integral inequalities.

The Hermite–Hadamard inequality is related to various classes of convexity; for some examples, see [15–19]. Iscan [20], in 2014, presented the idea of harmonic convexity and established a few related Hermite–Hadamard type inequalities. Harmonic h -convex functions and some associated Hermite–Hadamard inequalities were first described by the authors of [21] in 2015. Numerous researchers have linked integral inequalities with interval-valued functions in recent years, producing many significant findings. The Opial-type inequalities were introduced by Costa [22], the Ostrowski-type inequalities were investigated by Chalco-Cano [23] by using the generalized Hukuhara derivative, the Minkowski-type inequalities and the Beckenbach-type inequalities were established by Roman-Flores [24]. By introducing interval-valued coordinated convex functions and creating related H – H-type inequalities, Zhao et al. [25] recently improved on this idea. It was also utilized to support the H – H- and Fejér-type inequalities for the n -polynomial convex interval-valued function [26] and preinvex function [27,28]. Interval-valued coordinated preinvex functions are a recent extension of the interval-valued preinvex function notion introduced by Lai et al. [29]. Combined with interval analysis, the H – H inequality was extended to interval h -convex functions in [30], to interval harmonic h -convex functions in [31], to interval (h_1, h_2) -convex functions in [32] and to interval harmonically (h_1, h_2) -convex functions in [33]. The definition of the h -Godunova–Levin function was utilized by the authors in [34] to take into account this inequality. Additionally, the author in [35] published a fuzzy Jensen-type integral inequality for fuzzy interval-valued functions, while the authors in [36] created a Jensen-type inequality for (h_1, h_2) interval-nonconvex functions.

Our research is inspired by the strong literature and the specific articles [33,34]. The idea of interval-valued harmonically (m, h_1, h_2) -Godunova–Levin functions is introduced first, and new H – H-type inequalities are then constructed for the aforementioned notion. The structure of the paper is as follows: In Section 2, the introduction and the mathematical background are given. The issue and our key findings are discussed in Section 3. Section 5 contains the conclusion and future scope.

2. Preliminaries

We begin by introducing some of the terms, characteristics, and notations that will be utilized in the article. Let I be represented as the intervals of the collection of real numbers \mathbb{R} .

$[a] = [\underline{a}, \bar{a}] = \{x \in \mathbb{R} \mid \underline{a} \leq x \leq \bar{a}\}$, $\underline{a}, \bar{a} \in \mathbb{R}$, where the real interval $[a]$ is a closed and bounded subset of \mathbb{R} . We call $[a]$ positive when $\underline{a} > 0$, and $[a]$ is negative when $\bar{a} < 0$. Let us denote all intervals of the set of real numbers by R_I of \mathbb{R} , all positive intervals by R_I^+ , and all negative intervals by R_I^- . The inclusion " \subseteq " is defined as:

$$[a] \subseteq [b] \iff [\underline{a}, \bar{a}] \subseteq [\underline{b}, \bar{b}] \iff \underline{b} \leq \underline{a}, \bar{a} \leq \bar{b}.$$

Suppose λ is any real number, and $[a]$ is an interval; then, the $v[a]$ is given as:

$$\lambda \cdot [a, \bar{a}] = \begin{cases} [\lambda a, \lambda \bar{a}], & \text{if } \lambda > 0 \\ \{0\}, & \text{if } \lambda = 0 \\ [\lambda \bar{a}, \lambda a], & \text{if } \lambda < 0. \end{cases}$$

For $[a] = [a, \bar{a}]$, and $[b] = [b, \bar{b}]$, the following algebraic operations hold true:

$$\begin{aligned} [a] + [b] &= [a + b, \bar{a} + \bar{b}], \\ [a] - [b] &= [a - b, \bar{a} - \bar{b}], \\ [a] \cdot [b] &= [\min\{a\bar{b}, a\bar{b}, \bar{a}b, \bar{a}b\}, \max\{a\bar{b}, a\bar{b}, \bar{a}b, \bar{a}b\}], \\ [a] / [b] &= [\min\{a/b, a/\bar{b}, \bar{a}/b, \bar{a}/\bar{b}\}, \max\{a/b, a/\bar{b}, \bar{a}/b, \bar{a}/\bar{b}\}], \end{aligned}$$

where $0 \notin [a, \bar{a}]$.

For the intervals $[a, \bar{a}]$ and $[b, \bar{b}]$ the Hausdorff–Pompeiu distance is defined as:

$$d([a, \bar{a}], [b, \bar{b}]) = \max\{|a - b|, |\bar{a} - \bar{b}|\}.$$

Definition 1 (see [37]). Let $\mathcal{G} : [c, d] \rightarrow R_I$ be an interval valued function such that $\mathcal{G}(u) = [\underline{\mathcal{G}}(u), \overline{\mathcal{G}}(u)]$ for each $u \in [c, d]$. Then, the function \mathcal{G} is Riemann integrable over the interval $[c, d]$, and

$$(IR) \int_c^d \mathcal{G}(u) du = \left[(R) \int_c^d \underline{\mathcal{G}}(u) du, (R) \int_c^d \overline{\mathcal{G}}(u) du \right],$$

where $\underline{\mathcal{G}}, \overline{\mathcal{G}}$ are Riemann integrable over the interval $[c, d]$.

The set of all Riemann integrable interval-valued functions and real-valued functions are represented by the symbols $IR_{[c,d]}$ and $R_{[c,d]}$, respectively.

Definition 2 (see [20]). A set, $S \subset R - \{0\}$, is said to be a harmonic convex set if

$$\frac{cd}{uc + (1 - u)d} \in S,$$

where $\forall c, d \in S$ and $u \in [0, 1]$.

Definition 3 (see [38]). A nonnegative function $\mathcal{G} : S \rightarrow R$ is said to be a Godunova–Levin function, if

$$\mathcal{G}(uc + (1 - u)d) \leq \frac{\mathcal{G}(c)}{u} + \frac{\mathcal{G}(d)}{(1 - u)},$$

where $\forall c, d \in S$ and $u \in (0, 1)$.

Definition 4 (see [20]). A function $\mathcal{G} : S \rightarrow R$ is said to be a harmonically convex function, if

$$\mathcal{G}\left(\frac{cd}{uc + (1 - u)d}\right) \leq u\mathcal{G}(d) + (1 - u)\mathcal{G}(c),$$

where $\forall c, d \in S$ and $u \in [0, 1]$.

Definition 5 (see [21]). The function $\mathcal{G} : S \rightarrow R$ is said to be a harmonically h -convex function, if $\forall c, d \in S$ and $u \in [0, 1]$, we have

$$\mathcal{G}\left(\frac{cd}{uc + (1 - u)d}\right) \leq h(u)\mathcal{G}(d) + h(1 - u)\mathcal{G}(c),$$

where $h : [0, 1] \subseteq S \rightarrow R$ is a nonnegative function with $h \neq 0$.

Definition 6 (see [39]). The function $\mathcal{G} : S \rightarrow R$ is said to be a harmonically (m, h) -convex function, if $\forall c, d \in S, m \in (0, 1]$ and $u \in [0, 1]$, we have

$$\mathcal{G}\left(\frac{mcd}{muc + (1 - u)d}\right) \leq h(u)\mathcal{G}(d) + mh(1 - u)\mathcal{G}(c),$$

where $h : [0, 1] \subseteq S \rightarrow R$ is a nonnegative function with $h \neq 0$.

Definition 7 (see [33]). The function $\mathcal{G} : S \rightarrow R$ is said to be a harmonically (h_1, h_2) -convex function, if $\forall c, d \in S$ and $u \in [0, 1]$, we have

$$\mathcal{G}\left(\frac{cd}{uc + (1 - u)d}\right) \leq h_1(u)h_2(1 - u)\mathcal{G}(c) + h_1(1 - u)h_2(u)\mathcal{G}(d),$$

where $h : [0, 1] \subseteq S \rightarrow R$ is a nonnegative function with $h \neq 0$.

Definition 8 (see [34]). The function $\mathcal{G} : S \rightarrow R$ is said to be a h -Godunova–Levin function, if for all $c, d \in S$ and $u \in (0, 1)$, we have

$$\mathcal{G}(uc + (1 - u)d) \leq \frac{\mathcal{G}(c)}{h(u)} + \frac{\mathcal{G}(d)}{h(1 - u)},$$

where $h : (0, 1) \subseteq S \rightarrow R$ is a nonnegative function.

3. Main Results

We will now introduce harmonically interval-valued (m, h_1, h_2) Godunova–Levin functions (this idea was motivated by [32]).

Definition 9. A function $\mathcal{G} : S \rightarrow R$ is said to be a harmonically (m, h) -Godunova–Levin function, if for all $c, d \in S, u \in (0, 1)$ and $m \in (0, 1]$, we have

$$\mathcal{G}\left(\frac{mcd}{muc + (1 - u)d}\right) \leq \frac{\mathcal{G}(d)}{h(u)} + \frac{m\mathcal{G}(c)}{h(1 - u)},$$

where $h : (0, 1) \subseteq S \rightarrow R$ is a nonnegative function.

Definition 10. A function $\mathcal{G} : S \rightarrow R$ is said to be a harmonically (m, h_1, h_2) -Godunova–Levin function, if for all $c, d \in S, u \in (0, 1)$ and $m \in (0, 1]$, we have

$$\mathcal{G}\left(\frac{mcd}{muc + (1 - u)d}\right) \leq \frac{m\mathcal{G}(c)}{h_1(u)h_2(1 - u)} + \frac{\mathcal{G}(d)}{h_1(1 - u)h_2(u)},$$

where $h_1, h_2 : (0, 1) \subseteq S \rightarrow R$ is a nonnegative functions.

Remark 1. • If we choose $m = 1$ and $h_1(u) = h_2(u) = 1$ in Definition 10, then the notion of a harmonically P -convex function [21] is recovered.

• If we choose $m = 1, h_1(u) = 1,$ and $h_2(u) = \frac{1}{u}$ in Definition 10, then the notion of harmonically convex function [20] is recovered.

• If we choose $m = 1, h_1(u) = 1,$ and $h_2(u) = \frac{1}{h(u)}$ in Definition 10, then the notion of harmonic h -convex function [21] is recovered.

• If we choose $m = 1, h_1(u) = 1,$ and $h_2(u) = (u)^s$ in Definition 10, then the notion of harmonically s -Godunova–Levin function [40] is recovered.

Definition 11. The function $\mathcal{G} : S \rightarrow R_I^+$ is said to be a harmonically (m, h_1, h_2) -Godunova–Levin interval-valued convex function, if $\forall c, d \in S, u \in (0, 1)$ and $m \in (0, 1]$, we have

$$\frac{m\mathcal{G}(c)}{h_1(u)h_2(1-u)} + \frac{\mathcal{G}(d)}{h_1(1-u)h_2(u)} \subseteq \mathcal{G}\left(\frac{mcd}{muc + (1-u)d}\right), \tag{1}$$

where $h_1, h_2 : (0, 1) \subseteq S \rightarrow R$ are non-negative functions such that $h_1, h_2 \neq 0$.

If the above inequality is in reverse order, then the function \mathcal{G} is known as a harmonically (m, h_1, h_2) -Godunova–Levin concave interval-valued function. The space of all harmonically (m, h_1, h_2) -Godunova–Levin convex and (m, h_1, h_2) -Godunova–Levin concave interval-valued functions are denoted by $SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), S, R_I^+\right)$ and $SGHV\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), S, R_I^+\right)$, respectively.

Proposition 1. Suppose $\mathcal{G} : [x, y] \rightarrow R_I^+$ is a harmonically (m, h_1, h_2) -Godunova–Levin interval-valued function such that $\mathcal{G}(u) = [\underline{\mathcal{G}}(u), \overline{\mathcal{G}}(u)]$, $m \in (0, 1]$. Then, $\mathcal{G} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right)$ if and only if

$$\underline{\mathcal{G}} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right) \quad \text{and} \quad \overline{\mathcal{G}} \in SGHV\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right).$$

Proof. Let \mathcal{G} be a harmonically (m, h_1, h_2) -Godunova–Levin interval valued convex function. Assume $c, d \in [x, y], u \in (0, 1)$ and $m \in (0, 1]$. Then, we have

$$\frac{m\mathcal{G}(c)}{h_1(u)h_2(1-u)} + \frac{\mathcal{G}(d)}{h_1(1-u)h_2(u)} \subseteq \mathcal{G}\left(\frac{mcd}{muc + (1-u)d}\right).$$

That is,

$$\left[\frac{m\underline{\mathcal{G}}(c)}{h_1(u)h_2(1-u)} + \frac{\underline{\mathcal{G}}(d)}{h_1(1-u)h_2(u)}, \frac{m\overline{\mathcal{G}}(c)}{h_1(u)h_2(1-u)} + \frac{\overline{\mathcal{G}}(d)}{h_1(1-u)h_2(u)} \right] \subseteq \mathcal{G}\left(\frac{mcd}{muc + (1-u)d}\right).$$

It follows that

$$\frac{m\underline{\mathcal{G}}(c)}{h_1(u)h_2(1-u)} + \frac{\underline{\mathcal{G}}(d)}{h_1(1-u)h_2(u)} \geq \underline{\mathcal{G}}\left(\frac{mcd}{muc + (1-u)d}\right),$$

and

$$\frac{m\overline{\mathcal{G}}(c)}{h_1(u)h_2(1-u)} + \frac{\overline{\mathcal{G}}(d)}{h_1(1-u)h_2(u)} \leq \overline{\mathcal{G}}\left(\frac{mcd}{muc + (1-u)d}\right).$$

This shows that $\underline{\mathcal{G}} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right)$, and $\overline{\mathcal{G}} \in SGHV\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right)$. Conversely, let $\underline{\mathcal{G}} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right)$, and $\overline{\mathcal{G}} \in SGHV\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right)$. Based on the above definition and set inclusion, we obtain $\mathcal{G} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [x, y], R_I^+\right)$. This completes the proof. \square

Proposition 2. Suppose $\mathcal{G} : [c, d] \rightarrow R_I^+$ is a harmonically interval-valued (m, h_1, h_2) -Godunova–Levin function such that $\mathcal{G}(u) = [\underline{\mathcal{G}}(u), \overline{\mathcal{G}}(u)]$, $m \in (0, 1]$.

Then, $\mathcal{G} \in SGHV\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right)$ if and only if

$$\underline{\mathcal{G}} \in SGHV\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right) \quad \text{and} \quad \overline{\mathcal{G}} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right).$$

The proof can be conducted in a similar manner as in Proposition 1; hence, it is omitted for the readers.

Hermite–Hadamard Inequalities

Throughout, $H(x, y) = h_1(x)h_2(y) \forall x, y \in (0, 1)$.

Theorem 1. Let $h_1, h_2 : (0, 1) \rightarrow R^+$ and $\mathcal{G} : [c, d] \rightarrow R_I^+$ be an interval-valued function defined with $\underline{\mathcal{G}}, \overline{\mathcal{G}}$. If $\mathcal{G} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right)$, $m \in (0, 1]$, and $\mathcal{G} \in IR_{[c, d]}$, then

$$\frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]}{2} \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \supseteq \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)}. \tag{2}$$

Proof. Assume that $\mathcal{G} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right)$. Then,

$$\frac{\mathcal{G}(p)}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} + \frac{\mathcal{G}(q)}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \subseteq \mathcal{G}\left(\frac{2pq}{p+q}\right),$$

where

$$p = \frac{mcd}{m\sigma c + (1-\sigma)d},$$

and

$$q = \frac{mcd}{m(1-\sigma)c + \sigma d}; \text{ here } \sigma \in [0, 1].$$

Then,

$$\frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \left[\mathcal{G}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) + \mathcal{G}\left(\frac{mcd}{m(1-\sigma)c + \sigma d}\right) \right] \subseteq \mathcal{G}\left(\frac{2mcd}{mc+d}\right). \tag{3}$$

Multiplying both sides by

$$H\left(\frac{1}{2}, \frac{1}{2}\right),$$

we obtain

$$\left[\mathcal{G}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) + \mathcal{G}\left(\frac{mcd}{m(1-\sigma)c + \sigma d}\right) \right] \subseteq H\left(\frac{1}{2}, \frac{1}{2}\right) \mathcal{G}\left(\frac{2mcd}{mc+d}\right). \tag{4}$$

If we integrate the above inequality over $(0, 1)$ with respect to "σ", we have

$$\int_0^1 \left[\mathcal{G}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) + \mathcal{G}\left(\frac{mcd}{m(1-\sigma)c + \sigma d}\right) \right] d\sigma \subseteq H\left(\frac{1}{2}, \frac{1}{2}\right) \int_0^1 \mathcal{G}\left(\frac{2mcd}{mc+d}\right) d\sigma.$$

Thus,

$$\int_0^1 \underline{\mathcal{G}}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) d\sigma + \int_0^1 \underline{\mathcal{G}}\left(\frac{mcd}{m(1-\sigma)c + \sigma d}\right) d\sigma \geq H\left(\frac{1}{2}, \frac{1}{2}\right) \int_0^1 \underline{\mathcal{G}}\left(\frac{2mcd}{mc+d}\right) d\sigma,$$

and

$$\int_0^1 \overline{\mathcal{G}}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) d\sigma + \int_0^1 \overline{\mathcal{G}}\left(\frac{mcd}{m(1-\sigma)c + \sigma d}\right) d\sigma \leq H\left(\frac{1}{2}, \frac{1}{2}\right) \int_0^1 \overline{\mathcal{G}}\left(\frac{2mcd}{mc+d}\right) d\sigma.$$

It follows (change of variables) that

$$\frac{2mcd}{d - mc} \int_c^d \frac{\underline{\mathcal{G}}(u)}{u^2} du \geq H\left(\frac{1}{2}, \frac{1}{2}\right) \int_0^1 \underline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right) d\sigma = H\left(\frac{1}{2}, \frac{1}{2}\right) \underline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right)$$

and

$$\frac{2mcd}{d - mc} \int_c^d \frac{\overline{\mathcal{G}}(u)}{u^2} du \leq H\left(\frac{1}{2}, \frac{1}{2}\right) \int_0^1 \overline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right) d\sigma = H\left(\frac{1}{2}, \frac{1}{2}\right) \overline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right).$$

As a result, we have

$$\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right] \left[\underline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right), \overline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right) \right] \supseteq \frac{2mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du.$$

Upon dividing both sides by $\frac{1}{2}$, we obtain the desired first inclusion of Theorem 1.

$$\left[\frac{H\left(\frac{1}{2}, \frac{1}{2}\right)}{2} \right] \left[\underline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right), \overline{\mathcal{G}}\left(\frac{2mcd}{mc + d}\right) \right] \supseteq \frac{mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du. \tag{5}$$

From our hypothesis, we have

$$\frac{m\mathcal{G}(c)}{h_1(\sigma)h_2(1 - \sigma)} + \frac{\mathcal{G}(d)}{h_1(1 - \sigma)h_2(\sigma)} \subseteq \mathcal{G}\left(\frac{mcd}{m\sigma c + (1 - \sigma)d}\right).$$

$$\frac{m\mathcal{G}(c)}{h_1(1 - \sigma)h_2(\sigma)} + \frac{\mathcal{G}(d)}{h_1(\sigma)h_2(1 - \sigma)} \subseteq \mathcal{G}\left(\frac{mcd}{m(1 - \sigma)c + \sigma d}\right).$$

Adding the above two inclusions and integrating over $(0, 1)$ with respect to " σ " gives

$$[m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{1}{h_1(\sigma)h_2(1 - \sigma)} d\sigma + [m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{1}{h_1(1 - \sigma)h_2(\sigma)} d\sigma$$

$$\subseteq \int_0^1 \left[\mathcal{G}\left(\frac{mcd}{m\sigma c + (1 - \sigma)d}\right) + \mathcal{G}\left(\frac{mcd}{m(1 - \sigma)c + \sigma d}\right) \right] d\sigma.$$

It is easily seen that $\int_0^1 \frac{1}{h_1(\sigma)h_2(1 - \sigma)} d\sigma = \int_0^1 \frac{1}{h_1(1 - \sigma)h_2(\sigma)} d\sigma$. This implies

$$2[m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{1}{H(\sigma, 1 - \sigma)} d\sigma \subseteq \frac{2mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(\kappa)}{\kappa^2} d\kappa.$$

From the above developments, it follows

$$[m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{1}{H(\sigma, 1 - \sigma)} d\sigma \subseteq \frac{mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(\kappa)}{\kappa^2} d\kappa. \tag{6}$$

Now, combining (5) and (6), we obtain the required result

$$\left[\frac{H\left(\frac{1}{2}, \frac{1}{2}\right)}{2} \right] \mathcal{G}\left(\frac{2mcd}{mc + d}\right) \supseteq \frac{mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(\kappa)}{\kappa^2} d\kappa \supseteq [m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{d\sigma}{H(\sigma, 1 - \sigma)}.$$

□

Remark 2. (1) If we put $h_1(\sigma) = h_2(\sigma) = 1$ into Theorem 1, then it gives results for interval-valued harmonically (m, P) -functions.

$$\frac{1}{2} \mathcal{G}\left(\frac{2mcd}{mc + d}\right) \supseteq \frac{mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [m\mathcal{G}(c) + \mathcal{G}(d)]$$

(2) If we put $m = 1$ $h_1(\sigma) = h_2(\sigma) = 1$ into Theorem 1, then it gives results for interval-valued harmonically (m, P) -functions.

$$\frac{1}{2} \mathcal{G}\left(\frac{2cd}{c+d}\right) \supseteq \frac{cd}{d-c} \int_c^d \frac{\mathcal{G}(u)du}{u^2} \supseteq [\mathcal{G}(c) + \mathcal{G}(d)].$$

(3) Choosing $H(\sigma, y) = h(\sigma)$, Theorem 1 gives results for harmonically (m, h) -Godunova–Levin interval-valued functions.

$$\frac{\left[h\left(\frac{1}{2}\right)\right]}{2} \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \supseteq \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{d\sigma}{h(\sigma)}.$$

(4) Choosing $m = 1$ and $H(\sigma, y) = h(\sigma)$, Theorem 1 gives results for harmonically h -Godunova–Levin interval-valued functions.

$$\frac{\left[h\left(\frac{1}{2}\right)\right]}{2} \mathcal{G}\left(\frac{2cd}{c+d}\right) \supseteq \frac{cd}{d-c} \int_c^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 \frac{d\sigma}{h(\sigma)}.$$

(5) If we choose $H(\sigma, y) = \frac{1}{h(\sigma)}$, then Theorem 1 gives results for harmonically m -interval-valued h -convex functions.

$$\frac{1}{2\left[h\left(\frac{1}{2}\right)\right]} \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \supseteq \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 h(\sigma)d\sigma.$$

(6) If we choose $m = 1$ and $H(\sigma, y) = \frac{1}{h(\sigma)}$, then Theorem 1 gives results for harmonically interval-valued h -convex functions.

$$\frac{1}{2\left[h\left(\frac{1}{2}\right)\right]} \mathcal{G}\left(\frac{2cd}{c+d}\right) \supseteq \frac{cd}{d-c} \int_c^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 h(\sigma)d\sigma.$$

(7) Again, if we choose $H(\sigma, y) = \frac{1}{H(\sigma, y)}$ in Theorem 1, it recovers a result for harmonic (h_1, h_2) -convex interval-valued functions.

$$\frac{1}{2\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \supseteq \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [m\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 H(\sigma, 1-\sigma)d\sigma.$$

(8) Again, if we choose $m = 1$ and $H(\sigma, y) = \frac{1}{H(\sigma, y)}$ in Theorem 1, it recovers a result for harmonic (h_1, h_2) -convex interval-valued functions.

$$\frac{1}{2\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \mathcal{G}\left(\frac{2cd}{c+d}\right) \supseteq \frac{cd}{d-c} \int_c^d \frac{\mathcal{G}(u)}{u^2} du \supseteq [\mathcal{G}(c) + \mathcal{G}(d)] \int_0^1 H(\sigma, 1-\sigma)d\sigma.$$

Theorem 2. Let $\mathcal{G} : [c, d] \rightarrow R_1^+$ be an interval-valued function and $h_1, h_2 : (0, 1) \rightarrow R^+$ such that $H(\frac{1}{2}, \frac{1}{2}) \neq 0$. If $\mathcal{G} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_1^+\right)$, $m \in (0, 1)$ and $\mathcal{G} \in IR_{[c, d]}$; then, the following inequality holds true.

$$\begin{aligned} & \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2}{4} \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \supseteq \Delta_1 \supseteq \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \supseteq \Delta_2 \\ & \supseteq \left\{ m[\mathcal{G}(c) + \mathcal{G}(d)] \left[\frac{1}{2} + \frac{1}{H\left(\frac{1}{2}, \frac{1}{2}\right)} \right] \right\} \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)}, \end{aligned}$$

where

$$\begin{aligned} \Delta_1 &= \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]}{4} \left[\mathcal{G}\left(\frac{4mcd}{mc+3d}\right) + \mathcal{G}\left(\frac{4mcd}{md+3c}\right) \right], \\ \Delta_2 &= \left[\mathcal{G}\left(\frac{2mcd}{mc+d}\right) + m\left(\frac{\mathcal{G}(c) + \mathcal{G}(d)}{2}\right) \right] \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)}. \end{aligned}$$

Proof. Assume that $\mathcal{G} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_1^+\right)$ and $\mathcal{G} \in IR_{[c, d]}$. For $\left[mc, \frac{2cd}{c+d}\right]$, we have

$$\frac{\mathcal{G}\left(\frac{mc \frac{2cd}{c+d}}{m\sigma c + (1-\sigma) \frac{2cd}{c+d}}\right)}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} + \frac{\mathcal{G}\left(\frac{mc \frac{2cd}{c+d}}{m(1-\sigma)c + \sigma \frac{2cd}{c+d}}\right)}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \subseteq \mathcal{G}\left(\frac{4mcd}{mc+3d}\right),$$

and we obtain

$$\frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \left[\mathcal{G}\left(\frac{mc \frac{2cd}{c+d}}{m\sigma c + (1-\sigma) \frac{2cd}{c+d}}\right) + \mathcal{G}\left(\frac{mc \frac{2cd}{c+d}}{m(1-\sigma)c + \sigma \frac{2cd}{c+d}}\right) \right] \subseteq \mathcal{G}\left(\frac{4mcd}{mc+3d}\right).$$

Integrating over $(0, 1)$ with respect to " σ ", we have

$$\begin{aligned} & \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \left[\int_0^1 \underline{\mathcal{G}}\left(\frac{mc \frac{2cd}{c+d}}{m\sigma c + (1-\sigma) \frac{2cd}{c+d}}\right) d\sigma + \int_0^1 \underline{\mathcal{G}}\left(\frac{mc \frac{2cd}{c+d}}{m(1-\sigma)c + \sigma \frac{2cd}{c+d}}\right) d\sigma, \right. \\ & \left. \int_0^1 \overline{\mathcal{G}}\left(\frac{mc \frac{2cd}{c+d}}{m\sigma c + (1-\sigma) \frac{2cd}{c+d}}\right) d\sigma + \int_0^1 \overline{\mathcal{G}}\left(\frac{mc \frac{2cd}{c+d}}{m(1-\sigma)c + \sigma \frac{2cd}{c+d}}\right) d\sigma \right] \subseteq \mathcal{G}\left(\frac{4mcd}{mc+3d}\right). \end{aligned}$$

$$\begin{aligned} & \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]} \left[\frac{2mcd}{d-mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\mathcal{G}(u)}{u^2} du + \frac{2mcd}{d-mc} \int_{\frac{2cd}{c+d}}^d \frac{\mathcal{G}(u)}{u^2} du, \frac{2mcd}{d-mc} \int_{mc}^{\frac{2cd}{c+d}} \overline{\mathcal{G}}(u) du \right. \\ & \left. + \frac{2mcd}{d-mc} \int_{\frac{2cd}{c+d}}^d \overline{\mathcal{G}}(u) du \right] \subseteq \mathcal{G}\left(\frac{4mcd}{mc+3d}\right), \quad (7) \end{aligned}$$

i.e.,

$$\frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]} \left[\frac{4mcd}{d - mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\mathcal{G}(u)}{u^2} du, \frac{4mcd}{d - mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\overline{\mathcal{G}}(u)}{u^2} du \right] \subseteq \mathcal{G}\left(\frac{4mcd}{mc + 3d}\right).$$

Thus,

$$\frac{4}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]} \left[\frac{mcd}{d - mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\mathcal{G}(u)}{u^2} du, \frac{mcd}{d - mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\overline{\mathcal{G}}(u)}{u^2} du \right] \subseteq \mathcal{G}\left(\frac{4mcd}{mc + 3d}\right);$$

so,

$$\frac{4}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]} \left[\frac{mcd}{d - mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\mathcal{G}(u)}{u^2} du \right] \subseteq \mathcal{G}\left(\frac{4mcd}{mc + 3d}\right),$$

and consequently,

$$\frac{mcd}{d - mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\mathcal{G}(u)}{u^2} du \subseteq \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]}{4} \mathcal{G}\left(\frac{4mcd}{mc + 3d}\right). \tag{8}$$

Similarly, for the interval $\left[\frac{2cd}{c+d}, md\right]$, we can have

$$\frac{mcd}{d - mc} \int_{\frac{2cd}{c+d}}^{md} \frac{\mathcal{G}(u)}{u^2} du \subseteq \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]}{4} \mathcal{G}\left(\frac{4mcd}{md + 3c}\right). \tag{9}$$

Adding (8) and (9), we obtain

$$\Delta_1 = \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]}{4} \left[\mathcal{G}\left(\frac{4mcd}{3c + md}\right) + \mathcal{G}\left(\frac{4mcd}{mc + 3d}\right) \right] \supseteq \left[\frac{mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \right].$$

Consequently, we obtain

$$\begin{aligned} \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]^2}{4} \mathcal{G}\left(\frac{2mcd}{mc + d}\right) &= \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]^2}{4} \mathcal{G}\left(\frac{2 \frac{4mcd}{mc+3d} \frac{4mcd}{md+3c}}{\frac{4mcd}{mc+3d} + \frac{4mcd}{md+3c}}\right) \\ &\supseteq \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]^2}{4} \left[\frac{\mathcal{G}\left(\frac{4mcd}{mc+3d}\right)}{H\left(\frac{1}{2}, \frac{1}{2}\right)} + \frac{\mathcal{G}\left(\frac{4mcd}{md+3c}\right)}{H\left(\frac{1}{2}, \frac{1}{2}\right)} \right] \\ &= \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]^2}{4H\left(\frac{1}{2}, \frac{1}{2}\right)} \left[\mathcal{G}\left(\frac{4mcd}{mc + 3d}\right) + \mathcal{G}\left(\frac{4mcd}{md + 3c}\right) \right] \\ &= \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right) \right]}{4} \left[\mathcal{G}\left(\frac{4mcd}{mc + 3d}\right) + \mathcal{G}\left(\frac{4mcd}{md + 3c}\right) \right] \\ &= \Delta_1 \\ &\supseteq \frac{mcd}{d - mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du. \end{aligned}$$

A minor modification of the argument in Theorem 1 will give

$$\begin{aligned}
 & \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)}{u^2} du \\
 &= \frac{1}{2} \left[\frac{2mcd}{d-mc} \int_{mc}^{\frac{2cd}{c+d}} \frac{\mathcal{G}(u)}{u^2} du + \frac{2mcd}{d-mc} \int_{\frac{2cd}{c+d}}^{md} \frac{\mathcal{G}(u)}{u^2} du \right] \\
 &\supseteq \frac{1}{2} \left[\left[m\mathcal{G}(c) + \mathcal{G}\left(\frac{2cd}{c+d}\right) \right] \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} + \left[m\mathcal{G}(d) + \mathcal{G}\left(\frac{2cd}{c+d}\right) \right] \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} \right] \\
 &= \frac{1}{2} \left[\left\{ m\mathcal{G}(c) + m\mathcal{G}(d) + 2\mathcal{G}\left(\frac{2cd}{c+d}\right) \right\} \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} \right] \\
 &= \left[\frac{m\mathcal{G}(c) + m\mathcal{G}(d)}{2} + \mathcal{G}\left(\frac{2cd}{c+d}\right) \right] \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} \\
 &= \Delta_2 \\
 &\supseteq \left[\frac{m\mathcal{G}(c) + m\mathcal{G}(d)}{2} + \frac{m\mathcal{G}(c)}{H\left(\frac{1}{2}, \frac{1}{2}\right)} + \frac{m\mathcal{G}(d)}{H\left(\frac{1}{2}, \frac{1}{2}\right)} \right] \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} \\
 &= \left[\frac{m\mathcal{G}(c) + m\mathcal{G}(d)}{2} + \frac{1}{H\left(\frac{1}{2}, \frac{1}{2}\right)} [m\mathcal{G}(c) + m\mathcal{G}(d)] \right] \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} \\
 &= \left\{ m[\mathcal{G}(c) + \mathcal{G}(d)] \left[\frac{1}{2} + \frac{1}{H\left(\frac{1}{2}, \frac{1}{2}\right)} \right] \right\} \int_0^1 \frac{d\sigma}{H(\sigma, 1-\sigma)}.
 \end{aligned}$$

This completes the proof. \square

Theorem 3. Let $h_1, h_2 : (0, 1) \rightarrow R^+$ and $\mathcal{G}, \mathcal{V} : [c, d] \rightarrow R_I^+$ be interval-valued functions. If $\mathcal{G}, \mathcal{V} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right)$, $m \in (0, 1]$ and $\mathcal{G}\mathcal{V} \in IR_{[c,d]}$, then, we have

$$\begin{aligned}
 & \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du \\
 & \supseteq \mathcal{M}(c, d) \int_0^1 \frac{1}{H^2(\sigma, 1-\sigma)} d\sigma + \mathcal{N}(c, d) \int_0^1 \frac{1}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} d\sigma,
 \end{aligned}$$

where

$$\begin{aligned}
 \mathcal{M}(c, d) &= m^2\mathcal{G}(c)\mathcal{V}(c) + \mathcal{G}(d)\mathcal{V}(d). \\
 \mathcal{N}(c, d) &= m[\mathcal{G}(c)\mathcal{V}(d) + \mathcal{G}(d)\mathcal{V}(c)].
 \end{aligned}$$

Proof. Assume that $\mathcal{G}, \mathcal{V} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right)$. Then, we have

$$\begin{aligned}
 \mathcal{G}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) &\supseteq \frac{m\mathcal{G}(c)}{h_1(\sigma)h_2(1-\sigma)} + \frac{\mathcal{G}(d)}{h_1(1-\sigma)h_2(\sigma)}. \\
 \mathcal{V}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) &\supseteq \frac{m\mathcal{V}(c)}{h_1(\sigma)h_2(1-\sigma)} + \frac{\mathcal{V}(d)}{h_1(1-\sigma)h_2(\sigma)}.
 \end{aligned}$$

From the above hypotheses, we obtain

$$\begin{aligned} \mathcal{G}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right)\mathcal{V}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) & \supseteq \frac{m^2\mathcal{G}(c)\mathcal{V}(c)}{H^2(\sigma, 1-\sigma)} + \frac{m[\mathcal{G}(c)\mathcal{V}(d) + \mathcal{G}(d)\mathcal{V}(c)]}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} + \frac{\mathcal{G}(d)\mathcal{V}(d)}{H^2(1-\sigma, \sigma)}. \end{aligned}$$

Integrating both sides over (0, 1) with respect to "σ", we have

$$\begin{aligned} & \int_0^1 \mathcal{G}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right)\mathcal{V}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) d\sigma \\ &= \left[\int_0^1 \underline{\mathcal{G}}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right)\underline{\mathcal{V}}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) d\sigma, \right. \\ & \quad \left. \int_0^1 \overline{\mathcal{G}}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right)\overline{\mathcal{V}}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right) d\sigma \right] \\ &= \left[\frac{mcd}{d-mc} \int_{mc}^d \frac{\underline{\mathcal{G}}(u)\underline{\mathcal{V}}(u)}{u^2} du, \frac{mcd}{d-mc} \int_{mc}^d \frac{\overline{\mathcal{G}}(u)\overline{\mathcal{V}}(u)}{u^2} du \right] \\ &= \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du \\ &\supseteq \int_0^1 \frac{[m^2\mathcal{G}(c)\mathcal{V}(c) + \mathcal{G}(d)\mathcal{V}(d)]}{H^2(\sigma, 1-\sigma)} d\sigma + \int_0^1 \frac{m[\mathcal{G}(c)\mathcal{V}(d) + \mathcal{G}(d)\mathcal{V}(c)]}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} d\sigma. \end{aligned}$$

It follows that

$$\frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du \supseteq \mathcal{M}(c, d) \int_0^1 \frac{d\sigma}{H^2(\sigma, 1-\sigma)} + \mathcal{N}(c, d) \int_0^1 \frac{d\sigma}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)}.$$

The Theorem is proved. □

Theorem 4. Let $h_1, h_2 : (0, 1) \rightarrow R^+$ and $\mathcal{G}, \mathcal{V} : [c, d] \rightarrow R_I^+$ be interval-valued functions. If $\mathcal{G}, \mathcal{V} \in SGHX\left(\left(\frac{1}{h_1}, \frac{1}{h_2}\right), [c, d], R_I^+\right)$, $m \in (0, 1]$ and $\mathcal{G}\mathcal{V} \in IR_{[c, d]}$, then, we have

$$\begin{aligned} \left[\frac{H\left(\frac{1}{2}, \frac{1}{2}\right)}{1+m^2}\right]^2 \mathcal{G}\left(\frac{2mcd}{mc+d}\right)\mathcal{V}\left(\frac{2mcd}{mc+d}\right) & \supseteq \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du \\ & + \mathcal{M}(c, d) \int_0^1 \frac{1+m^2}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} d\sigma + \mathcal{N}(c, d) \int_0^1 \frac{m}{H^2(\sigma, 1-\sigma)} d\sigma. \end{aligned}$$

Proof. By hypothesis, one has

$$\begin{aligned} \mathcal{G}\left(\frac{2mcd}{mc+d}\right) & \supseteq \frac{\mathcal{G}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right)}{H\left(\frac{1}{2}, \frac{1}{2}\right)} + \frac{\mathcal{G}\left(\frac{mcd}{m\sigma d + (1-\sigma)c}\right)}{H\left(\frac{1}{2}, \frac{1}{2}\right)}. \\ \mathcal{V}\left(\frac{2mcd}{mc+d}\right) & \supseteq \frac{\mathcal{V}\left(\frac{mcd}{m\sigma c + (1-\sigma)d}\right)}{H\left(\frac{1}{2}, \frac{1}{2}\right)} + \frac{\mathcal{V}\left(\frac{mcd}{m\sigma d + (1-\sigma)c}\right)}{H\left(\frac{1}{2}, \frac{1}{2}\right)}. \end{aligned}$$

Then,

$$\begin{aligned}
 \mathcal{G}\left(\frac{2mcd}{mc+d}\right)\mathcal{V}\left(\frac{2mcd}{mc+d}\right) &\supseteq \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\mathcal{G}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right)\mathcal{V}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right) \right. \\
 &\quad \left. + \mathcal{G}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right)\mathcal{V}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right) \right] \\
 &\quad + \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\mathcal{G}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right)\mathcal{V}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right) \right. \\
 &\quad \left. + \mathcal{G}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right)\mathcal{V}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right) \right] \\
 &\supseteq \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\mathcal{G}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right)\mathcal{V}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right) \right. \\
 &\quad \left. + \mathcal{G}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right)\mathcal{V}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right) \right] \\
 &\quad + \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\left(\frac{m\mathcal{G}(c)}{H(\sigma, 1-\sigma)} + \frac{\mathcal{G}(d)}{H(1-\sigma, \sigma)}\right) \left(\frac{m\mathcal{V}(c)}{H(1-\sigma, \sigma)} + \frac{\mathcal{V}(d)}{H(\sigma, 1-\sigma)}\right) \right. \\
 &\quad \left. + \left(\frac{\mathcal{G}(c)}{H(1-\sigma, \sigma)} + \frac{m\mathcal{G}(d)}{H(\sigma, 1-\sigma)}\right) \left(\frac{\mathcal{V}(c)}{H(\sigma, 1-\sigma)} + \frac{m\mathcal{V}(d)}{H(1-\sigma, \sigma)}\right) \right] \\
 &= \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\mathcal{G}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right)\mathcal{V}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right) \right. \\
 &\quad \left. + \mathcal{G}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right)\mathcal{V}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right) \right] \\
 &\quad + \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\left(\frac{1+m^2}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)}\right) [\mathcal{G}(c)\mathcal{V}(c) + \mathcal{G}(d)\mathcal{V}(d)] \right. \\
 &\quad \left. + m\left(\frac{1}{H^2(\sigma, 1-\sigma)} + \frac{1}{H^2(1-\sigma, \sigma)}\right) [\mathcal{G}(c)\mathcal{V}(d) + \mathcal{G}(d)\mathcal{V}(c)] \right] \\
 &= \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\mathcal{G}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right)\mathcal{V}\left(\frac{mcd}{m\sigma c+(1-\sigma)d}\right) \right. \\
 &\quad \left. + \mathcal{G}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right)\mathcal{V}\left(\frac{mcd}{m\sigma d+(1-\sigma)c}\right) \right] \\
 &\quad + \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\left(\frac{1+m^2}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)}\right) \mathcal{M}(c, d) \right. \\
 &\quad \left. + m\left(\frac{1}{H^2(\sigma, 1-\sigma)} + \frac{1}{H^2(1-\sigma, \sigma)}\right) \mathcal{N}(c, d) \right].
 \end{aligned}$$

Integrating with respect to "σ" over (0, 1), we have

$$\begin{aligned} & \int_0^1 \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \mathcal{V}\left(\frac{2mcd}{mc+d}\right) d\sigma \\ &= \left[\int_0^1 \underline{\mathcal{G}}\left(\frac{2mcd}{mc+d}\right) \mathcal{V}\left(\frac{2mcd}{mc+d}\right) d\sigma, \int_0^1 \overline{\mathcal{G}}\left(\frac{2mcd}{mc+d}\right) \mathcal{V}\left(\frac{2mcd}{mc+d}\right) d\sigma \right] = \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \mathcal{V}\left(\frac{2mcd}{mc+d}\right) \\ &\supseteq \frac{1+m^2}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du \right] \\ &+ \frac{1}{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2} \left[\mathcal{M}(c, d) \int_0^1 \frac{1+m^2 d\sigma}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} + \mathcal{N}(c, d) \int_0^1 \frac{md\sigma}{H^2(\sigma, 1-\sigma)} \right]. \end{aligned}$$

Multiplying both sides of the above equation by $\frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2}{2}$, we obtain

$$\begin{aligned} & \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2}{1+m^2} \mathcal{G}\left(\frac{2mcd}{mc+d}\right) \mathcal{V}\left(\frac{2mcd}{mc+d}\right) \\ &\supseteq \frac{mcd}{d-mc} \int_{mc}^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du + \mathcal{M}(c, d) \int_0^1 \frac{1+m^2}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} d\sigma \\ &+ \mathcal{N}(c, d) \int_0^1 \frac{md\sigma}{H^2(\sigma, 1-\sigma)}. \end{aligned}$$

This completes the proof. □

4. Examples

Example 1. Let $m = 1$, $h_1(\sigma) = \frac{1}{\sigma}$, and $h_2(\sigma) = 1$ where $\sigma \in (0, 1)$, $[c, d] = \left[\frac{1}{2}, 1\right]$ and $\mathcal{G} : [c, d] \rightarrow R_1^+$ is defined by $\mathcal{G}(u) = [u^2, 5 - e^u]$. Then,

$$\begin{aligned} & \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]}{2} f\left(\frac{2cd}{c+d}\right) = \mathcal{G}\left(\frac{2}{3}\right) \approx \left[\frac{4}{9}, 5 - e^{\frac{2}{3}}\right]. \\ & \frac{cd}{d-c} \int_c^d \frac{\mathcal{G}(u)}{u^2} du = \left[\int_{\frac{1}{2}}^1 du, \int_{\frac{1}{2}}^1 \frac{(5 - e^u)}{u^2} du \right] \approx \left[\frac{1}{2}, 2.98\right]. \\ & [\mathcal{G}(c) + \mathcal{G}(d)] \int_{\frac{1}{2}}^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} = [\mathcal{G}(c) + \mathcal{G}(d)] \int_{\frac{1}{2}}^1 \sigma d\sigma \approx \left[\frac{15}{32}, \frac{3(10 - \sqrt{e} - e)}{8}\right]. \end{aligned}$$

Thus, we obtain

$$\left[\frac{4}{9}, 5 - e^{\frac{2}{3}}\right] \supseteq \left[\frac{1}{2}, 2.98\right] \supseteq \left[\frac{15}{32}, \frac{3(10 - \sqrt{e} - e)}{8}\right].$$

This verifies Theorem 1.

Example 2. For $m = 1$, $h_1(\sigma) = \frac{1}{\sigma}$, and $h_2(\sigma) = 1$, where $\sigma \in (0, 1)$, $[c, d] = \left[\frac{1}{2}, 1\right]$ and $\mathcal{G} : [c, d] \rightarrow R_1^+$ is defined as $\mathcal{G}(u) = [u^2, 4 - e^u]$. Then,

$$\begin{aligned} & \left[\frac{H\left(\frac{1}{2}, \frac{1}{2}\right)}{4} \right]^2 \mathcal{G}\left(\frac{2cd}{c+d}\right) = \mathcal{G}\left(\frac{2}{3}\right) \approx \left[\frac{4}{9}, 4 - e^{\frac{2}{3}} \right]. \\ \Delta_1 &= \frac{H\left(\frac{1}{2}, \frac{1}{2}\right)}{4} \left[\mathcal{G}\left(\frac{4cd}{c+3d}\right) + \mathcal{G}\left(\frac{4cd}{d+3c}\right) \right] \\ &= \frac{1}{2} \left[\left[\frac{16}{49}, 4 - e^{\frac{4}{7}} \right] + \left[\frac{16}{25}, 4 - e^{\frac{4}{5}} \right] \right] \approx \left[\frac{592}{1225}, \frac{8 - e^{\frac{4}{7}} - e^{\frac{4}{5}}}{2} \right]. \\ \Delta_2 &= \left[f\left(\frac{2cd}{c+d}\right) + \left(\frac{\mathcal{G}(c) + \mathcal{G}(d)}{2}\right) \right] \int_{\frac{1}{2}}^1 \frac{d\sigma}{H(\sigma, 1-\sigma)} \\ &= \frac{3}{8} \left[\mathcal{G}\left(\frac{2}{3}\right) + \left(\frac{\mathcal{G}\left(\frac{1}{2}\right) + \mathcal{G}(1)}{2}\right) \right] \approx \left[\frac{77}{192}, \frac{-3e - 3\sqrt{e} - 6e^{\frac{2}{3}} + 48}{16} \right]. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} \left[\frac{4}{9}, 4 - e^{\frac{2}{3}} \right] &\supseteq \left[\frac{592}{1225}, \frac{8 - e^{\frac{4}{7}} - e^{\frac{4}{5}}}{2} \right] \supseteq \left[\frac{1}{2}, 2.98 \right] \\ &\supseteq \left[\frac{77}{192}, \frac{-3e - 3\sqrt{e} - 6e^{\frac{2}{3}} + 48}{16} \right]. \end{aligned}$$

This verifies Theorem 2.

Example 3. Let $m = 1$, $h_1(\sigma) = \frac{1}{\sigma}$, and $h_2(\sigma) = 1$, where $\sigma \in (0, 1)$, $[c, d] = \left[\frac{1}{2}, 1\right]$, and $\mathcal{G}, \mathcal{V} : [c, d] \rightarrow R_1^+$ are defined as $\mathcal{G}(u) = [u^2, 4 - e^u]$ and $\mathcal{V}(u) = [u, 3 - u^2]$. Then,

$$\begin{aligned} \frac{cd}{d-c} \int_c^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du &= \left[\int_{\frac{1}{2}}^1 u du, \int_{\frac{1}{2}}^1 \frac{(4 - e^u)(3 - u^2)}{u^2} du \right] \approx \left[\frac{3}{8}, 5.0094 \right]. \\ \mathcal{M}(c, d) \int_0^1 \frac{1}{H^2(\sigma, 1-\sigma)} d\sigma &= \mathcal{M}\left(\frac{1}{2}, 1\right) \int_0^1 \sigma^2 d\sigma \approx \left[\frac{9}{24}, \frac{19}{3} - \frac{2\sqrt{e}}{3} - \frac{11e}{12} \right]. \\ \mathcal{N}(c, d) \int_0^1 \frac{1}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} d\sigma &= \mathcal{N}\left(\frac{1}{2}, 1\right) \int_0^1 (\sigma - \sigma^2) d\sigma \approx \left[\frac{3}{24}, \frac{19}{6} - \frac{11\sqrt{e}}{24} - \frac{e}{3} \right]. \end{aligned}$$

It follows that

$$\begin{aligned} \left[\frac{3}{8}, 5.0094 \right] &\supseteq \left[\frac{9}{24}, \frac{19}{3} - \frac{2\sqrt{e}}{3} - \frac{11e}{12} \right] + \left[\frac{3}{24}, \frac{19}{6} - \frac{11\sqrt{e}}{24} - \frac{e}{3} \right] \\ &= \left[\frac{1}{2}, \frac{19}{6} + \frac{-30e - 27\sqrt{e} + 152}{24} \right]. \end{aligned}$$

This verifies Theorem 3.

Example 4. Let $m = 1$, $h_1(\sigma) = \frac{1}{\sigma}$, and $h_2(\sigma) = 1$, where $\sigma \in (0, 1)$, $[c, d] = \left[\frac{1}{2}, 1\right]$, and $\mathcal{G}, \mathcal{V} : [c, d] \rightarrow R_1^+$ are defined as $\mathcal{G}(u) = [u^2, 4 - e^u]$ and $\mathcal{V}(u) = [u, 3 - u^2]$. Then,

$$\begin{aligned} \frac{\left[H\left(\frac{1}{2}, \frac{1}{2}\right)\right]^2}{2} \mathcal{G}\left(\frac{2cd}{c+d}\right) \mathcal{V}\left(\frac{2cd}{c+d}\right) &= 2\mathcal{G}\left(\frac{2}{3}\right) \mathcal{V}\left(\frac{2}{3}\right) \approx \left[\frac{16}{27}, \frac{46(4-e^{\frac{2}{3}})}{9}\right]. \\ \frac{cd}{d-c} \int_c^d \frac{\mathcal{G}(u)\mathcal{V}(u)}{u^2} du &= \left[\int_{\frac{1}{2}}^1 u du, \int_{\frac{1}{2}}^1 \frac{(4-e^u)(3-u^2)}{u^2} du\right] \approx \left[\frac{3}{8}, 5.0094\right]. \\ \mathcal{M}(c, d) \int_0^1 \frac{1}{H(\sigma, \sigma)H(1-\sigma, 1-\sigma)} d\sigma &= \mathcal{M}\left(\frac{1}{2}, 1\right) \int_0^1 (\sigma - \sigma^2) d\sigma = \left[\frac{9}{48}, \frac{19}{6} - \frac{\sqrt{e}}{3} - \frac{11e}{24}\right]. \\ \mathcal{N}(c, d) \int_0^1 \frac{1}{H^2(\sigma, 1-\sigma)} d\sigma &= \mathcal{N}\left(\frac{1}{2}, 1\right) \int_0^1 \sigma^2 d\sigma \approx \left[\frac{3}{12}, \frac{19}{3} - \frac{11\sqrt{e}}{12} - \frac{2e}{3}\right]. \end{aligned}$$

It follows that

$$\begin{aligned} \left[\frac{16}{27}, \frac{184-46e^{\frac{2}{3}}}{9}\right] &\supseteq \left[\frac{3}{8}, 5.0094\right] + \left[\frac{9}{48}, \frac{19}{6} - \frac{\sqrt{e}}{3} - \frac{11e}{24}\right] \\ &+ \left[\frac{3}{12}, \frac{19}{3} - \frac{11\sqrt{e}}{12} - \frac{2e}{3}\right] \\ &= \left[\frac{13}{16}, \frac{19}{6} + \frac{-27e - 30\sqrt{e} + 152}{24} + 5.0094\right]. \end{aligned}$$

This verifies Theorem 4.

5. Conclusions

Interval-valued functions may be a valuable alternative for incorporating uncertainty into the prediction processes. We first introduced a new notion of interval-valued harmonic convexity, i.e., a harmonically interval-valued (h_1, h_2) -Godunova–Levin function and established the Hermite–Hadamard and Pachpatt- type inequalities employing this new notion. Our new definition generalized many existing definitions present in the literature. We thereby added to the extension of many classical integral inequalities in the set-valued setting. Some numerical examples were provided to further explain the results.

Future presentations of various inequalities, including those of the Hermite–Hadamard, Ostrowski, Hadamard–Mercer, Simpson, Fejér, and Bullen types, could make use of this novel idea. The inequalities indicated above can be demonstrated for a variety of interval-valued LR convexities, fuzzy interval convexities, and CR convexities. These findings will also be generalized in relation to quantum calculus, coordinated interval-valued functions, fractional calculus, etc. Due to the fact that these are the most active areas of study in the subject of integral inequalities, many mathematicians will be interested in investigating how different types of interval-valued analysis might be applied to the integral inequalities.

Author Contributions: Conceptualization, S.K.S., P.O.M., D.O.R., and K.N.; Data curation, S.K.S., P.O.M., and M.T.; Formal analysis, S.K.S.; Funding acquisition, K.N.; Investigation, S.K.S., P.O.M., D.O.R., M.T., and K.N.; Methodology, M.T.; Project administration, D.O.R. and K.N.; Resources, D.O.R.; Software, P.O.M. and M.T.; Supervision, D.O.R. and K.N.; Validation, K.N.; Visualization, K.N.; Writing—original draft, S.K.S. and P.O.M.; Writing—review and editing, D.O.R. and M.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Acknowledgments: This research was supported by the National Science, Research and Innovation Fund (NSRF), Thailand.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Moore, R.E. *Interval Analysis*; Prentice-Hall: Hoboken, NJ, USA, 1966.
2. Gasilov, N.A.; Emrah Amrahov, S. Solving a nonhomogeneous linear system of interval differential equations. *Soft Comput.* **2018**, *22*, 3817–3828. [[CrossRef](#)]
3. De Weerd, E.; Chu, Q.P.; Mulder, J.A. Neural network output optimization using interval analysis. *IEEE Trans. Neural Netw.* **2009**, *20*, 638–653. [[CrossRef](#)] [[PubMed](#)]
4. Rothwell, E.J.; Cloud, M.J. Automatic error analysis using intervals. *IEEE Trans. Edu.* **2011**, *55*, 9–15. [[CrossRef](#)]
5. Snyder, J. Interval analysis for computer graphics. *SIGGRAPH Comput. Graph.* **1992**, *26*, 121–130. [[CrossRef](#)]
6. Chalco-Cano, Y.; Rufián-Lizana, A.; Román-Flores, H.; Jiménez-Gamero, M.D. Calculus for interval-valued functions using generalized Hukuhara derivative and applications. *Fuzzy Sets Syst.* **2013**, *219*, 49–67. [[CrossRef](#)]
7. Chalco-Cano, Y.; Silva, G.N.; Rufián-Lizana, A. On the Newton method for solving fuzzy optimization problems. *Fuzzy Sets Syst.* **2015**, *272*, 60–69. [[CrossRef](#)]
8. Entani, T.; Inuiguchi, M. Pairwise comparison based interval analysis for group decision aiding with multiple criteria. *Fuzzy Sets Syst.* **2015**, *274*, 79–96. [[CrossRef](#)]
9. Osuna-Gómez, R.; Chalco-Cano, Y.; Hernández-Jiménez, B.; Ruiz-Garzón, G. Optimality conditions for generalized differentiable interval-valued functions. *Information* **2015**, *321*, 136–146. [[CrossRef](#)]
10. Moore, R.E.; Kearfott, R.B.; Cloud, M.J. *Introduction to Interval Analysis*; Society for Industrial and Applied Mathematics (SIAM): Philadelphia, PA, USA, 2009.
11. Chalco-Cano, Y.; Flores-Franulić, A.; Román-Flores, H. Ostrowski type inequalities for interval-valued functions using generalized Hukuhara derivative. *Comput. Appl. Math.* **2012**, *31*, 457–472.
12. Costa, T.M.; Román-Flores, H. Some integral inequalities for fuzzy-interval-valued functions. *Information* **2017**, *420*, 110–125. [[CrossRef](#)]
13. Srivastava, H.M.; Sahoo, S.K.; Mohammed, P.O.; Kodamasingh, B.; Nonlaopon, K.; Abualnaja, K.M. Interval valued Hadamard-Fejér and Pachpatte Type inequalities pertaining to a new fractional integral operator with exponential kernel. *AIMS Math.* **2022**, *7*, 15041–15063. [[CrossRef](#)]
14. Kalsoom, H.; Latif, M.A.; Khan, Z.A.; Vivas-Cortez, M. Some New Hermite-Hadamard-Fejér fractional type inequalities for h -convex and harmonically h -Convex interval-valued Functions. *Mathematics*, **2021**, *10*, 74. [[CrossRef](#)]
15. Dragomir, S.S.; Pečarić, J.; Persson, L.E. Some inequalities of Hadamard type. *Soochow J. Math.* **1995**, *21*, 335–341.
16. Dragomir, S.S. Inequalities of Hermite-Hadamard type for functions of selfadjoint operators and matrices. *J. Math. Inequal.* **2017**, *11*, 241–259. [[CrossRef](#)]
17. Latif, M. On Some New Inequalities of Hermite-Hadamard Type for Functions Whose Derivatives are s -convex in the Second Sense in the Absolute Value. *Ukr. Math. J.* **2016**, *67*, 1552–1571. [[CrossRef](#)]
18. Noor, M.A.; Cristescu, G.; Awan, M.U. Generalized Fractional Hermite-Hadamard Inequalities for Twice Differentiable s -convex Functions. *Filomat* **2015**, *29*, 807–815. [[CrossRef](#)]
19. Noor, M.A.; Noor, K.I.; Awan, M.U.; Li, J. On Hermite-Hadamard Inequalities for h -Preinvex Functions. *Filomat* **2014**, *28*, 1463–1474. [[CrossRef](#)]
20. İşcan, İ. Hermite-Hadamard type inequalities for harmonically convex functions. *Hacet. J. Math. Stat.* **2014**, *43*, 935–942. [[CrossRef](#)]
21. Noor, M.A.; Noor, K.I.; Awan, M.U.; Costache, S. Some integral inequalities for harmonically h -convex functions. *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.* **2015**, *77*, 5–16.
22. Costa, T.M.; Román-Flores, H.; Chalco-Cano, Y. Opial-type inequalities for interval-valued functions. *Fuzzy Set. Syst.* **2019**, *358*, 48–63. <https://doi.org/10.1016/j.fss.2018.04.012>
23. Chalco-Cano, Y.; Lodwick, W.A. Condori-Equice. Ostrowski type inequalities and applications in numerical integration for interval-valued functions. *Soft Comput.* **2015**, *19*, 3293–3300. [[CrossRef](#)]
24. Román-Flores, H.; Chalco-Cano, Y.; Lodwick, W.A. Some integral inequalities for interval-valued functions. *Comput. Appl. Math.* **2018**, *37*, 1306–1318. [[CrossRef](#)]
25. Zhao, D.; Ali, M.A.; Murtaza, G.; Zhang, Z. On the Hermite-Hadamard inequalities for interval-valued coordinated convex functions. *Adv. Differ. Equ.* **2020**, *2020*, 1–14. [[CrossRef](#)]
26. Nwaeze, E.R.; Khan, M.A.; Chu, Y.M. Fractional inclusions of the Hermite-Hadamard type for m -polynomial convex interval-valued functions. *Adv. Differ. Equ.* **2020**, *2020*, 1–17. [[CrossRef](#)]
27. Sharma, N.; Singh, S.K.; Mishra, S.K.; Hamdi, A. Hermite-Hadamard type inequalities for interval-valued preinvex functions via Riemann-Liouville fractional integrals. *J. Inequal. Appl.* **2021**, *2021*, 98. [[CrossRef](#)]
28. Srivastava, H.M.; Sahoo, S.K.; Mohammed, P.O.; Baleanu, D.; Kodamasingh, B. Hermite-Hadamard type inequalities for interval-valued preinvex functions via fractional integral operators. *Int. J. Comput. Intel. Syst.* **2022**, *15*, 1–12. [[CrossRef](#)]
29. Lai, K.K.; Bisht, J.; Sharma, N.; Mishra, S.K. Hermite-Hadamard-Type Fractional Inclusions for Interval-Valued Preinvex Functions. *Mathematics* **2022**, *10*, 264. [[CrossRef](#)]
30. Zhao, D.; An, T.; Ye, G.; Liu, W. New Jensen and Hermite-Hadamard type inequalities for h -convex interval-valued functions. *J. Inequal. Appl.* **2018**, *302*, 1–14. [[CrossRef](#)]
31. Zhao, D.; An, T.; Ye, G.; Torres, D.F. On Hermite-Hadamard type inequalities for harmonical h -convex interval-valued functions. *arXiv* **2019**, arXiv:1911.06900. [[CrossRef](#)]

32. An, Y.; Ye, G.; Zhao, D.; Liu, W. Hermite-Hadamard type inequalities for interval (h_1, h_2) -convex functions. *Mathematics* **2019**, *7*, 436.
33. Liu, R.; Xu, R. Hermite-Hadamard type inequalities for harmonical (h_1, h_2) convex interval-valued functions. *Math. Found. Comput.* **2021**, *4*, 89. [[CrossRef](#)]
34. Almutairi, O.; Kiliçman, A.A.A. Some integral inequalities for h -Godunova-Levin preinvexity. *Symmetry* **2019**, *11*, 1500. [[CrossRef](#)]
35. Costa, T. Jensen's inequality type integral for fuzzy-interval-valued functions. *Fuzzy Sets Syst.* **2017**, *327*, 31–47. [[CrossRef](#)]
36. Bai, H.; Saleem, M.S.; Nazeer, W.; Zahoor, M.S.; Zhao, T. Hermite-Hadamard and Jensen type inequalities for interval nonconvex function. *J. Math.* **2020**, *2020*, 3945384. [[CrossRef](#)]
37. Dinghas, A. Zum Minkowskischen Integralbegriff abgeschlossener Mengen. *Math. Z.* **1956**, *66*, 173–188. [[CrossRef](#)]
38. Godunova, E.K.; Levin, V.I. Neravenstva dlja funkci sirokogo klassa, soderzascego vypuklye, monotonnnye i nekotorye drugie vidy funkii. *Vycislitel. Mat. i. Fiz. Mezvuzov. Sb. Nauc. Tr. MGPI Moskva.* **1985**, *9*, 138–142. [[CrossRef](#)]
39. Jung, C.Y.; Yussouf, M.; Chu, Y.M.; Farid, G.; Kang, S.M. Generalized fractional Hadamard and Fejér–Hadamard inequalities for generalized harmonically convex functions. *J. Math.* **2020**, *2020*, 8245324. [[CrossRef](#)]
40. Awan, M.U. Integral inequalities for harmonically s -Godunova-Levin functions. *FU Math Inform.* **2015**, *29*, 415–424. [[CrossRef](#)]