

Article

On Implicit Time–Fractal–Fractional Differential Equation

McSylvester Ejighikeme Omaba ¹ , Soh Edwin Mukiawa ¹  and Eze R. Nwaeze ^{2,*} 

¹ Department of Mathematics, College of Science, University of Hafr Al Batin, P.O. Box 1803, Hafr Al Batin 31991, Saudi Arabia; mcomaba@uhb.edu.sa (M.E.O.); mukiawa@uhb.edu.sa (S.E.M.)

² Department of Mathematics and Computer Science, Alabama State University, Montgomery, AL 36101, USA

* Correspondence: enwaeze@alasu.edu

Abstract: An implicit time–fractal–fractional differential equation involving the Atangana’s fractal–fractional derivative in the sense of Caputo with the Mittag–Leffler law type kernel is studied. Using the Banach fixed point theorem, the well-posedness of the solution is proved. We show that the solution exhibits an exponential growth bound, and, consequently, the long-time (asymptotic) property of the solution. We also give examples to illustrate our problem.

Keywords: well-posedness; exponential growth bound; fractal–fractional operators; Mittag–Leffler type kernel

MSC: 26A33; 28A80; 34A08; 34A09

1. Introduction

The fractal–fractional differential equation is a link between the fractal and fractional differential equations. Fractal and fractional differential equations are known for modeling complex physical processes and phenomena, particularly irregular systems with memory. Although fractional equations are renowned for representing systems with long-term memory and long-range interactivity, fractal calculus, conversely, is immensely effective in working with occurrence in stratified or porous media. That is, fractal–fractional differential operator models physical phenomena and real-world activities that exhibit or display fractional behaviours (sponge-like media, aquifer, turbulence, etc.) namely finance, viscoelasticity, control theory, electrical networks, groundwater flow and geo-hydrology, wave propagation, plasma physics and fusion, rheology, chaotic processes, fluid mechanics and biological activities [1–6]. For more applications of fractal-fractional differential equations, see [7–10] and for recent results on fractional differential equations and their applications, see [11–13]. To explore more results on implicit fractional differential equations and their applications, see [14–17]. There are many results relating to implicit fractional differential equations in literature involving Caputo fractional derivatives both for initial value problems (IVP) and boundary value problems (BVP) [15,18–21].

In 2015, Benchohra and Soudi in [18] studied the existence of integrable solutions for IVP for some given implicit fractional order functional differential equations with infinite delay

$$\begin{cases} {}^C\mathcal{D}^\mu\phi(\tau) = \sigma(\tau, \phi_\tau, {}^C\mathcal{D}^\mu\phi_\tau), \tau \in I = [0, b], \\ \phi(\tau) = \omega(\tau), \tau \in (-\infty, 0], \end{cases}$$

where ${}^C\mathcal{D}^\mu$ is the Caputo fractional differential operator, $\sigma : I \times \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{R}$ is a given function and \mathbb{B} is a phase space with its element $\phi_\tau(\epsilon) = \phi(\tau + \epsilon)$, $\epsilon \in [-\infty, 0]$.

In 2016, Kucche et al., in [20], considered the following equation:

$${}^C\mathcal{D}^\mu\phi(\tau) = \sigma(\tau, \phi(\tau), {}^C\mathcal{D}^\mu\phi(\tau)), \phi(0) = \phi_0 \in \mathbb{R}, \tau \in [0, T], \quad (1)$$



Citation: Omaba, M.E.; Mukiawa, S.E.; Nwaeze, E.R. On Implicit Time–Fractal–Fractional Differential Equation. *Axioms* **2022**, *11*, 348. <https://doi.org/10.3390/axioms11070348>

Academic Editor: Adrian Petrusel

Received: 28 June 2022

Accepted: 18 July 2022

Published: 20 July 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

where ${}^C\mathcal{D}^\mu$ ($0 < \mu < 1$) stands for the Caputo fractional derivative and $\sigma : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a known continuous function fulfilling some conditions. The authors investigated the well-posedness, interval of existence, and continuous dependence on the initial condition of solutions to Equation (1). Recently, in 2021, Shabbir et al. [17] worked on an implicit boundary value problem (BVP) involving an Atangana–Baleanu–Caputo (ABC) derivative of the form

$$\begin{cases} {}_0^{ABC}\mathcal{D}_\tau^\mu \phi(\tau) = \sigma(\tau, \phi(\tau), {}_0^{ABC}\mathcal{D}_\tau^\mu \phi(\tau)), & 1 < \mu \leq 2, \tau \in I = [0, b], \\ \phi(0) = \phi_0, \phi(b) = \phi_1, \end{cases} \tag{2}$$

where ${}_0^{ABC}\mathcal{D}_\tau^\mu$ denotes the ABC derivative of order $1 < \mu \leq 2$ and $\sigma : I \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function. Here, the authors established the existence of solution, uniqueness of solution and stability of solution to the class of implicit BVPs (2) with an ABC type derivative and integral.

Motivated by some applications of the implicit fractal–fractional differential equation in modeling complex phenomena and systems in porous media with memory, and the result in [17], where the authors used the ABC derivative operator to study Equation (2); therefore, we generalize (2) for a class of fractal–fractional derivative operator known as the Mittag–Leffler kernel law Fractal–Fractional (FFM), to study the well-posedness, exponential growth bound, and long-time behaviour of a solution to the class of implicit time–fractal–fractional differential equation:

$$\begin{cases} {}_a^{FFM}\mathcal{D}_t^{\mu, \nu} \psi(t) = \varsigma \left(t, \psi(t), {}_a^{FFM}\mathcal{D}_t^{\mu, \nu} \psi(t) \right), & 0 < a < t \leq T < \infty, \\ \psi(a) = \psi_a, \end{cases} \tag{3}$$

with ψ_a taken to be a bounded and non-negative function, ${}_a^{FFM}\mathcal{D}_t^{\mu, \nu}$ represents Atangana’s fractal–fractional derivative of orders $\mu, \nu \in (0, 1]$ in the sense of Caputo with generalized Mittag–Leffler law type kernel, $\varsigma : [a, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz continuous. Information within our disposal, suggests that we are the first to study this class of implicit fractal–fractional differential equation. Using similar ideas in [2,3], we give the formulation of the solution to Equation (3) as follows:

Definition 1. Let $\varsigma : [a, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function. Then, the IVP (3) is equivalent to

$$\begin{aligned} \psi(t) = & \psi_a + \frac{(1 - \mu)\nu}{\mathbb{A}\mathbb{B}(\mu)} t^{\nu-1} \varsigma \left(t, \psi(t), {}_a^{FFM}\mathcal{D}_t^{\mu, \nu} \psi(t) \right) \\ & + \frac{\mu\nu}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \int_a^t \tau^{\nu-1} (t - \tau)^{\mu-1} \varsigma \left(\tau, \psi(\tau), {}_a^{FFM}\mathcal{D}_\tau^{\mu, \nu} \psi(\tau) \right) d\tau, \end{aligned} \tag{4}$$

which follows by the definition of the operator ${}_a^{FFM}\mathcal{I}_t^{\mu, \nu}$.

Next, we define the norm of the solution ψ by

$$\|\psi\| := \sup_{a \leq t \leq T} |\psi(t)|.$$

The organization of the paper is as follows. In Section 2, we present the preliminaries; and in Section 3, we give the statements and proofs of the main results of the paper. Section 5 contains a brief summary of the paper.

2. Preliminaries

In this section, one gives some concepts that will be useful for the main result.

Definition 2 ([22]). Suppose $\phi : (a, b) \rightarrow \mathbb{R}$ is differentiable and $0 < \mu < 1$. Then the ABC fractional derivative for function ϕ of order μ is defined as

$${}^{\text{ABC}}\mathcal{D}_t^\mu \phi(t) = \frac{\mathbb{M}(\mu)}{1-\mu} \int_a^t \phi'(\tau) E_\mu \left[-\frac{\mu}{1-\mu} (t-\tau)^\mu \right] d\tau, \tag{5}$$

with $\mathbb{M}(\mu) > 0$ a normalization function satisfying $\mathbb{M}(0) = \mathbb{M}(1) = 1$.

Definition 3 ([1–3]). Let $\phi : (a, b) \rightarrow \mathbb{R}$ be a differentiable function, suppose ϕ is fractal differentiable in (a, b) with order $v > 0$. Then, the fractal–fractional derivative of $\phi(t)$ of order μ in Caputo sense with the Mittag–Leffler kernel is defined by

$${}^{\text{FFM}}\mathcal{D}_t^{\mu,v} \phi(t) = \frac{\mathbb{A}\mathbb{B}(\mu)}{1-\mu} \int_a^t \frac{d\phi(\tau)}{d\tau^v} E_\mu \left[-\frac{\mu}{1-\mu} (t-\tau)^\mu \right] d\tau, \tag{6}$$

with $0 < \mu, v \leq 1$ and $\mathbb{A}\mathbb{B}(\mu) = 1 - \mu + \frac{\mu}{\Gamma(\mu)}$. The generalized form is given by

$${}^{\text{FFM}}\mathcal{D}_t^{\mu,v,\epsilon} \phi(t) = \frac{\mathbb{A}\mathbb{B}(\mu)}{1-\mu} \int_a^t \frac{d^\epsilon \phi(\tau)}{d\tau^v} E_\mu \left[-\frac{\mu}{1-\mu} (t-\tau)^\mu \right] d\tau, \quad 0 < \mu, v, \theta \leq 1,$$

where

$$\frac{d^\epsilon \phi(\tau)}{d\tau^v} = \lim_{t \rightarrow \tau} \frac{\phi^\epsilon(t) - \phi^\epsilon(\tau)}{t^v - \tau^v}.$$

Remark 1. When $v = 1$ in Equation (6), one obtains Equation (5).

Definition 4 ([1–3]). Let $\phi : (a, b) \rightarrow \mathbb{R}$ be a continuous function. Then, the fractal–fractional integral of ϕ with order μ possessing Mittag–Leffler type kernel is defined as

$${}^{\text{FFM}}\mathcal{I}_t^{\mu,v} \phi(t) = \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} \phi(\tau) d\tau + \frac{(1-\mu)v t^{v-1}}{\mathbb{A}\mathbb{B}(\mu)} \phi(t).$$

Definition 5 ([23]). One defines the incomplete beta function by

$$B_\tau(\mu, v) = \int_0^\tau t^{\mu-1} (1-t)^{v-1} dt, \quad \tau \in [0, 1].$$

It also has a representation in terms of a hypergeometric function given by

$$B_\tau(\mu, v) = \frac{\tau^\mu}{\mu} {}_2F_1(\mu, 1-v; \mu+1; \tau).$$

Definition 6 ([24]). The regularized incomplete beta function is given by

$$I_\tau(\mu, v) = \frac{B_\tau(\mu, v)}{B(\mu, v)} = \frac{1}{B(\mu, v)} \int_0^\tau \tau^{\mu-1} (1-\tau)^{v-1} d\tau,$$

satisfying the following properties:

1. $I_\tau(\mu, v) = I_\tau(\mu + 1, v - 1) + \frac{\tau^\mu (1-\tau)^{v-1}}{\mu B(\mu, v)}$;
2. $I_\tau(\mu, v) = I_\tau(\mu + 1, v + 1) - \frac{\tau^\mu (1-\tau)^{v-1}}{v B(\mu, v)}$;
3. $I_\tau(\mu, v) = I_\tau(\mu + 1, v) + \frac{\tau^\mu (1-\tau)^v}{\mu B(\mu, v)}$;
4. $I_\tau(\mu, v) = I_\tau(\mu, v + 1) - \frac{\tau^\mu (1-\tau)^v}{v B(\mu, v)}$;
5. $I_\tau(\mu, v) + I_{1-\tau}(v, \mu) = 1$;
6. $I_1(\mu, v) = 1$ and $I_\tau(\mu, v) \in [0, 1]$.

Lemma 1 ([25]). For all $\mu > 0, v \geq 1, 0 \leq \tau \leq 1$, we have

1. $\frac{\tau^\mu(1-\tau)^{v-1}}{\mu B(\mu, v)} \leq I_\tau(\mu, v)$;
2. $I_\tau(\mu, v) \leq \frac{\tau^\mu}{\mu B(\mu, v)}$.

3. Main Results

This section starts with a Lipschitz condition on $\zeta(\cdot)$:

Condition 1. Let $0 < \text{Lip}_\zeta < \infty$. Given that $\xi_1, \xi_2, \zeta_1, \zeta_2 \in \mathbb{R}$, one has

$$|\zeta(\tau, \xi_1, \zeta_1) - \zeta(\tau, \xi_2, \zeta_2)| \leq \text{Lip}_\zeta (|\xi_1 - \xi_2| + |\zeta_1 - \zeta_2|),$$

with $\zeta(\tau, 0, 0) = 0$ and

$$|\zeta(\tau, \xi_1, \zeta_1)| \leq \text{Lip}_\zeta (|\xi_1| + |\zeta_1|). \tag{7}$$

Lemma 2. Let $0 < \text{Lip}_\zeta < 1$ and Condition 1 holds. Then ${}^{FFM}\mathcal{D}_t^{\mu, v}$ is a global Lipschitz continuous operator.

Proof. From Equations (3) and (7), we have

$$\left| {}^{FFM}\mathcal{D}_\tau^{\mu, v} \psi(\tau) \right| = \left| \zeta\left(\tau, \psi(\tau), {}^{FFM}\mathcal{D}_\tau^{\mu, v} \psi(\tau)\right) \right| \leq \text{Lip}_\zeta \left(|\psi(\tau)| + \left| {}^{FFM}\mathcal{D}_\tau^{\mu, v} \psi(\tau) \right| \right),$$

and, therefore,

$$\left| {}^{FFM}\mathcal{D}_\tau^{\mu, v} \psi(\tau) \right| \leq \frac{\text{Lip}_\zeta}{1 - \text{Lip}_\zeta} |\psi(\tau)|.$$

Furthermore, one obtains

$$\left| {}^{FFM}\mathcal{D}_\tau^{\mu, v} \psi(\tau) - {}^{FFM}\mathcal{D}_\tau^{\mu, v} \varphi(\tau) \right| \leq \frac{\text{Lip}_\zeta}{1 - \text{Lip}_\zeta} |\psi(\tau) - \varphi(\tau)|.$$

□

3.1. Existence and Uniqueness Result

Here, we establish the well-posedness of solution to Equation (3). Now, define

$$\begin{aligned} \mathcal{A}\psi(t) &= \psi_a + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \zeta\left(t, \psi(t), {}^{FFM}\mathcal{D}_t^{\mu, v} \psi(t)\right) \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} \zeta\left(\tau, \psi(\tau), {}^{FFM}\mathcal{D}_\tau^{\mu, v} \psi(\tau)\right) d\tau, \end{aligned}$$

and obtain the following auxiliary results:

Lemma 3. Let ψ be a solution satisfying Equation (4) and let Condition 1 be satisfied. Then, it follows that for all $\mu, v \in (0, 1]$ such that $\mu + v \geq 1$, we have

$$\|\mathcal{A}\psi\| \leq c_1 + c_2 \frac{\text{Lip}_\zeta}{1 - \text{Lip}_\zeta} \|\psi\|,$$

where c_1 and $c_2 := \frac{v}{\mathbb{A}\mathbb{B}(\mu)} \left[(1-\mu)a^{v-1} + \frac{\mu\Gamma(v)}{\Gamma(\mu+v)} T^{\mu+v-1} \right]$ are positive constants with $|\psi_a| \leq c_1$.

Proof. By taking absolute value on the operator \mathcal{A} , we have

$$\begin{aligned} |\mathcal{A}\psi(t)| &\leq |\psi_a| + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \left| \zeta\left(t, \psi(t), {}^{FFM}\mathcal{D}_t^{\mu,v} \psi(t)\right) \right| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \left| \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} \zeta\left(\tau, \psi(\tau), {}^{FFM}\mathcal{D}_\tau^{\mu,v} \psi(\tau)\right) d\tau \right| \\ &\leq |\psi_a| + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \left| \zeta\left(t, \psi(t), {}^{FFM}\mathcal{D}_t^{\mu,v} \psi(t)\right) \right| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} \left| \zeta\left(\tau, \psi(\tau), {}^{FFM}\mathcal{D}_\tau^{\mu,v} \psi(\tau)\right) \right| d\tau. \end{aligned}$$

Applying Condition 1 and $|\varphi_a| \leq c_1$, to obtain

$$\begin{aligned} |\mathcal{A}\psi(t)| &\leq c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \text{Lip}_\zeta \left[|\psi(t)| + \left| {}^{FFM}\mathcal{D}_t^{\mu,v} \psi(t) \right| \right] \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} \text{Lip}_\zeta \left[|\varphi(\tau)| + \left| {}^{FFM}\mathcal{D}_\tau^{\mu,v} \psi(\tau) \right| \right] d\tau. \end{aligned}$$

From Lemma 2, we arrive at

$$\begin{aligned} |\mathcal{A}\psi(t)| &\leq c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \text{Lip}_\zeta \left[|\psi(t)| + \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \right] \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} \text{Lip}_\zeta \left[|\psi(\tau)| + \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(\tau)| \right] d\tau \\ &= c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} |\psi(\tau)| d\tau \\ &\leq c_1 + \frac{(1-\mu)\beta}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\psi\| \int_a^t \tau^{v-1} (t-\tau)^{\mu-1} d\tau. \end{aligned}$$

Evaluating the integral above, we have

$$\begin{aligned} |\mathcal{A}\psi(t)| &\leq c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\psi\| t^{\mu+v-1} [B(\mu, v) - B_{\frac{t}{\Gamma}}(\mu, v)] \\ &= c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\psi\| B(\mu, v) t^{\mu+v-1} [1 - I_{\frac{t}{\Gamma}}(\mu, v)] \\ &\leq c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{v-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\psi\| B(\mu, v) t^{\mu+v-1}, \end{aligned} \tag{8}$$

since $1 - I_a^\mu(\mu, v) \in [0, 1]$. We observe that $a^{v-1} > t^{v-1}$ since $v - 1 < 0$. Thus, taking supremum over $t \in [a, T]$ in (8) and recalling that $\mu + v \geq 1$, we obtain

$$\begin{aligned} \|\mathcal{A}\psi\| &\leq c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} a^{v-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\psi\| + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\varphi\| B(\mu, v) T^{\mu+v-1} \\ &= c_1 + \frac{v}{\mathbb{A}\mathbb{B}(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \left[(1-\mu)a^{v-1} + \frac{\mu\Gamma(v)}{\Gamma(\mu+v)} T^{\mu+v-1} \right] \|\psi\|, \end{aligned}$$

and the proof is complete. \square

Lemma 4. Suppose ψ and φ are solutions satisfying Equation (4) and let Condition 1 be satisfied. Then, if it follows that for all $\mu, v \in (0, 1]$ such that $\mu + v \geq 1$, we have

$$\|\mathcal{A}\psi - \mathcal{A}\varphi\| \leq c_2 \|\psi - \varphi\|.$$

Proof. The proof is skipped since it follows similar steps as the proof of Lemma 3. \square

Next, we state the existence and uniqueness theorem for Equation (3).

Theorem 1. Let $\alpha + \beta \geq 1$ and suppose Condition 1 is satisfied. Let $c_2 > 0$, such that $c_2 < \frac{1-\text{Lip}_\zeta}{\text{Lip}_\zeta}$, where $c_2 := \frac{v}{\mathbb{A}\mathbb{B}(\mu)} \left[(1-\mu)a^{v-1} + \frac{\mu\Gamma(v)}{\Gamma(\mu+v)} T^{\mu+v-1} \right]$. Then, there exists a unique solution to Equation (3).

Proof. We proceed by applying the Banach fixed point theorem. Let $\psi(t) = \mathcal{A}\psi(t)$, then using Lemma 3, we have

$$\|\psi\| = \|\mathcal{A}\psi\| \leq c_1 + c_2 \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\psi\|.$$

Collecting similar terms, we have $\|\psi\| \left[1 - c_2 \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \right] \leq c_1$. This shows that $\|\psi\| < \infty$ since $c_2 < \frac{1-\text{Lip}_\zeta}{\text{Lip}_\zeta}$. Furthermore, if $\psi \neq \varphi$ are solutions to Equation (3), then from Lemma 4, we have

$$\|\psi - \varphi\| = \|\mathcal{A}\psi - \mathcal{A}\varphi\| \leq c_2 \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \|\psi - \varphi\|.$$

It follows that $\|\psi - \varphi\| \left[1 - c_2 \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \right] \leq 0$. Since $1 - c_2 \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} > 0$, that is, $c_2 < \frac{1-\text{Lip}_\zeta}{\text{Lip}_\zeta}$, then $\|\psi - \varphi\| < 0$. This is a contradiction and, therefore, $\|\psi - \varphi\| = 0$. \square

Exponential Growth

We present an inequality needed in proving the upper growth bound:

Theorem 2 ([26]). Given that $f, g, h : I \rightarrow \mathbb{R}^+$ are continuous functions. If $\phi : I \rightarrow \mathbb{R}^+$ is continuous and

$$\phi(t) \leq f(t) + \int_{t_0}^t (t-\tau)^{v-1} g(\tau) \phi(\tau) d\tau + \int_{t_0}^t (t-\tau)^{v-1} h(\tau) \phi^\gamma(\tau) d\tau, t \in I,$$

with constants $v > 0$ and $0 < \gamma < 1$, then the following statements are true.

(i) Given that $\nu > \frac{1}{2}$. It follows that

$$\begin{aligned} \phi(t) &\leq \left[A_1^{1-\gamma}(t) \right. \\ &\quad \left. + (1-\gamma)K_1 \int_{t_0}^t \exp\left((\gamma-1)K_1 \int_{t_0}^s g^2(\tau)d\tau\right) h^2(s)R_1(s)ds \right]^{\frac{1}{2(1-\gamma)}} \\ &\quad \times \exp\left(t + \frac{K_1}{2} \int_{t_0}^t g^2(s)ds\right), \quad t \in I, \end{aligned}$$

where $A_1(t) = \max_{t_0 \leq s \leq t} \{2e^{-2s} f^2(s)\}$, $K_1 = \frac{2\Gamma(2\nu-1)}{4^{\nu-1}}$, and $R_1(t) = e^{2(\rho-1)t}$.

(ii) Given that $\rho \in (0, \frac{1}{2}]$, $\varrho = \frac{1+\nu}{\nu}$ and $p = 1 + \nu$. Then

$$\begin{aligned} \phi(t) &\leq \left[A_2^{1-\gamma}(t) \right. \\ &\quad \left. + (1-\gamma)K_2 \int_{t_0}^t \exp\left((\gamma-1)K_2 \int_{t_0}^s g^\varrho(\tau)d\tau\right) h^\varrho(s)R_2(s)ds \right]^{\frac{1}{\varrho(1-\gamma)}} \\ &\quad \times \exp\left(t + \frac{K_2}{\varrho} \int_{t_0}^t g^\varrho(s)ds\right), \quad t \in I, \end{aligned}$$

where $A_2(t) = \max_{t_0 \leq s \leq t} \{2^{\varrho-1} e^{-s\varrho} f^\varrho(s)\}$, $K_2 = 2^{2\varrho-2} \left(\frac{\Gamma(1-(1-\rho)p}{p^{1-(1-\nu)p}}\right)^{\frac{\varrho}{p}}$, and $R_2(t) = e^{\varrho(\nu-1)t}$.

Theorem 3. Given that ψ satisfies Equation (4) and the initial function ψ_a is bounded above. Suppose Condition 1 is satisfied, then it follows that for all $\mu \in (\frac{1}{2}, 1]$ and $\nu \in [0, 1]$, one gets

$$|\psi(t)| \leq c_5 \exp(t - c_6 t^{\nu-1}), t \in [a, T],$$

where $c_5 = (2e^{-2a} c_3^2)^{1/2} \exp\left(\frac{a^{\nu-1}}{1-\nu}\right)$ and $c_6 = (c_4 \text{Lip}_\zeta)^2 \frac{\Gamma(2\mu-1)}{4^{\mu-1}} \frac{1}{1-\nu}$ are some positive numbers.

Proof. Following from the line of proof of Theorem 3, one obtains

$$\begin{aligned} |\psi(t)| &\leq c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} t^{\nu-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \int_a^t \tau^{\nu-1} (t-\tau)^{\mu-1} |\psi(\tau)| d\tau \\ &\leq c_1 + \frac{(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} a^{\nu-1} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} |\psi(t)| \\ &\quad + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \int_a^t \tau^{\nu-1} (t-\tau)^{\mu-1} |\psi(\tau)| d\tau. \end{aligned} \tag{9}$$

Let $Y(t) := |\psi(t)|$ and $\tilde{c}_2 := 1 - \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \frac{a^{\nu-1}(1-\mu)v}{\mathbb{A}\mathbb{B}(\mu)} > 0$. Thus, for $\text{Lip}_\zeta < \frac{\mathbb{A}\mathbb{B}(\mu)}{\mathbb{A}\mathbb{B}(\mu) + v(1-\mu)a^{\nu-1}}$, it follows from (9) that

$$\tilde{c}_2 Y(t) \leq c_1 + \frac{\mu v}{\mathbb{A}\mathbb{B}(\mu)\Gamma(\mu)} \frac{\text{Lip}_\zeta}{1-\text{Lip}_\zeta} \int_a^t \tau^{\nu-1} (t-\tau)^{\mu-1} Y(\tau) d\tau.$$

Now, dividing by $\tilde{c}_2 = \frac{\mathbb{A}\mathbb{B}(\mu) - \text{Lip}_\zeta [\mathbb{A}\mathbb{B}(\mu) + v(1-\mu)a^{v-1}]}{(1 - \text{Lip}_\zeta)\mathbb{A}\mathbb{B}(\mu)}$, to get

$$Y(t) \leq c_3 + c_4 \text{Lip}_\zeta \int_a^t \tau^{v-1} (t - \tau)^{\mu-1} Y(\tau) d\tau,$$

with $c_3 := \frac{c_1}{\tilde{c}_2}$ and $c_4 := \frac{\mu v}{\Gamma(\mu)(\mathbb{A}\mathbb{B}(\mu) - \text{Lip}_\zeta [\mathbb{A}\mathbb{B}(\mu) + (1-\mu)va^{v-1}])}$. Next, using Theorem 2 (i) with $h(\tau) = 0, g(\tau) = c_4 \text{Lip}_\zeta \tau^{v-1}$ and $v = \mu > \frac{1}{2}$, to obtain

$$\begin{aligned} Y(t) &\leq A_1^{1/2}(t) \exp\left(t + \frac{K_1}{2} (c_4 \text{Lip}_\zeta)^2 \int_a^t \tau^{2v-2} d\tau\right) \\ &= A_1^{1/2}(t) \exp\left(t + (c_4 \text{Lip}_\zeta)^2 \frac{\Gamma(2\mu - 1)}{4^{\mu-1}} \frac{t^{v-1} - a^{v-1}}{v - 1}\right) \\ &= (2e^{-2a} c_3^2)^{1/2} \exp\left(\frac{a^{v-1}}{1 - v}\right) \exp\left(t - (c_4 \text{Lip}_\zeta)^2 \frac{\Gamma(2\mu - 1)}{4^{\mu-1}} \frac{t^{v-1}}{1 - v}\right), \end{aligned}$$

where $K_1 = \frac{2\Gamma(2\mu-1)}{4^{\mu-1}}$ and $A_1(t) = \max_{a \leq s \leq t} \{2e^{-2s} c_3^2\} = 2e^{-2a} c_3^2$, since e^{-2t} is decreasing. \square

3.2. Asymptotic Property of the Solution

Here, we show the long term (limiting) property of our solution. The corollary indicates that the rate of energy growth of the solution is finite when time becomes large.

Corollary 1. *Under the hypotheses of Theorem 3 and for all $0 < \beta < 1$, we have*

$$\limsup_{t \rightarrow \infty} \frac{\log |\varphi(t)|}{t} \leq 1.$$

Proof. We obtained from Theorem 3 that

$$|\psi(t)| \leq c_5 \exp(t - c_6 t^{v-1}), t \in [a, T].$$

If we take log of both sides of the above equation, it will yield

$$\log |\psi(t)| \leq \log(c_5) + t - c_6 t^{v-1}.$$

Next, divide through by t to obtain

$$\frac{\log |\psi(t)|}{t} \leq \frac{\log(c_5)}{t} + 1 - c_6 \frac{t^{v-1}}{t} = \frac{\log(c_5)}{t} + 1 - \frac{c_6}{t^{2-v}}.$$

Since $0 < v < 1$, it follows that $2 - v \geq 1$. Now, take limit supremum over t in both sides to get

$$\limsup_{t \rightarrow \infty} \frac{\log |\psi(t)|}{t} \leq \limsup_{t \rightarrow \infty} \frac{\log(c_5)}{t} + 1 - \limsup_{t \rightarrow \infty} \frac{c_6}{t^{2-v}} = 1.$$

\square

4. Examples

Now, we give examples to illustrate the result in Theorem 3. The following are some plots (graphs) for the upper bound growth of our energy solution $|\psi(t)| \leq \exp(t - \frac{1}{1-v} t^{v-1}), t \in [a, T]$. For convenience, we set $c_5 = 1$ and choose $\mu \in (\frac{1}{2}, 1]$, such that $c_6 = \frac{1}{1-v}$ with $c_4 = 1, \text{Lip}_\zeta = 1$. We plotted graphs of the growth bound for $v = \frac{1}{10}, \frac{1}{5}, \frac{1}{4}, \frac{1}{2}, \frac{2}{3}, \frac{8}{10}, \frac{9}{10}$, and for various time intervals. It is observed that as closer the pa-

parameter ν is to zero, the faster the rate of growth is to the bound. However, as time grows large, the growth rate is at most at $t = 600$ irrespective of the values of ν , as shown in the Figure 1 below.

Some graphs of the upper growth bound: $\exp(t - \frac{1}{1-\beta} t^{\beta-1})$ for $\beta = \frac{1}{10}, \frac{1}{5}, \frac{1}{4}, \frac{1}{2}, \frac{2}{3}, \frac{8}{10}, \frac{9}{10}$, and for various time intervals.




Figure 1. Graphical illustration of the growth bounds.

5. Conclusions

Fractional order derivatives are used to represent memory formalism in modeling phenomena or processes in porous media in order to diminish the size of the pores and the permeability of the porous matrix [27]. Hence, implicit fractal–fractional differential equations are very important because they model many technical processes and systems in porous environment exhibiting long time memory property. In this paper, we estimated the higher growth bound of our solution and it is shown that the solution exhibits an exponential growth in t at a specific rate. Furthermore, the result shows a long time behaviour of the mild solution. Banach fixed point theorem was applied to prove the well-posedness of mild solution to the class of implicit time–fractal–fractional differential

equation with Mittag–Leffler law. For future work, one can investigate the lower growth estimate of the solution, the stability of the solution, and the continuous dependence on the initial condition, as shown in [20].

Author Contributions: Conceptualization, M.E.O.; methodology, M.E.O. and S.E.M.; software, M.E.O.; validation, M.E.O., S.E.M. and E.R.N.; formal analysis, M.E.O., S.E.M. and E.R.N.; writing—original draft preparation, M.E.O.; writing—review and editing, M.E.O., S.E.M. and E.R.N.; funding acquisition, M.E.O. All authors have read and agreed to the published version of the manuscript.

Funding: This research work was funded by Institutional Funding Program under project number IFP-A-2022-2-1-09.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors gratefully acknowledge technical and financial support from the Agency for Research and Innovation, Ministry of Education and University of Hafr Al Batin, Saudi Arabia. We also thank the anonymous referees for their comments and suggestions.

Conflicts of Interest: The authors declare that there is no conflict of interest.

References

- Atangana, A. Fractal–fractional differentiation and integration: Connecting fractal calculus and fractional calculus to predict complex system. *Chaos Solit. Fract.* **2017**, *102*, 396–406. [\[CrossRef\]](#)
- Atangana, A.; Akgül, A.; Owolabi, K.M. Analysis of Frac-tal–fractional differential equations. *Alex. Eng. J.* **2020**, *59*, 1117–1134. [\[CrossRef\]](#)
- Araz, S.I. Numerical analysis of a new Volterra integro-differential equation involving fractal–fractional operators. *Chaos Solit. Fract.* **2020**, *130*, 109396. [\[CrossRef\]](#)
- Owolabi, K.M.; Atangana, A.; Akgül, A. Modelling and Analysis of Fractal-fractional partial differential equations: Application to reaction-diffusion model. *Alex. Eng. J.* **2020**, *59*, 2477–2490. [\[CrossRef\]](#)
- He, J.-H. Fractal calculus and its geometrical explanation. *Results Phys.* **2018**, *10*, 272–276. [\[CrossRef\]](#)
- Hosseininia, M.; Heydari, M.H.; Avazzadeh, Z. The Numerical Treatment of Non-linear Fractal–fractional 2D EMDEN-FOWLER equation utilizing 2D Chelyshkov polynomials. *Fractals* **2020**, *28*, 2040042. [\[CrossRef\]](#)
- Gómez-Aguilar, J.F.; Atangana, A. New chaotic attractors: Applications of fractal-fractional differentiation and integration. *Math. Methods Appl. Sci.* **2020**, *44*, 3036–3065. [\[CrossRef\]](#)
- Saad, K.M.; Alqhtani, M.; Gómez-Aguilar, J.F. Fractal-fractional study of the hepatitis C virus infection model. *Results Phys.* **2020**, *19*, 103555. [\[CrossRef\]](#)
- Zuniga-Aguilar, C.J.; Gómez-Aguilar, J.F.; Romero-Ugalde, H.M.; Ja-hanshahi, H.; Alsaadi, F.E. Fractal-fractional neuro-adaptive method for system identification. *Eng. Comput.* **2020**, *240*, 1–24. [\[CrossRef\]](#)
- Abro, K.A.; Atangana, A.; Gómez-Aguilar, J.F. Ferromagnetic chaos in thermal convection of fluid through fractal-fractional differentiation. *J. Therm. Anal. Calorim.* **2022**, *147*, 8461–84733. [\[CrossRef\]](#)
- Nane, E.; Nwaeze, E.R.; Omaba, M.E. Asymptotic behavior and non-existence of global solution to a class of conformable time-fractional stochastic differential equation. *Stat. Probab. Lett.* **2020**, *163*, 108792. [\[CrossRef\]](#)
- Omaba, M.E. On Space-Fractional Heat Equation with Non-homogeneous Fractional Time Poisson Process. *Progr. Fract. Differ. Appl.* **2020**, *6*, 67–79. [\[CrossRef\]](#)
- Omaba, M.E.; Nwaeze, E.R. Moment Bound of Solution to a Class of Conformable Time-Fractional Stochastic Equation. *Fractal Fract.* **2019**, *3*, 18. [\[CrossRef\]](#)
- Asma; Gómez-Aguilar, J.F.; ur Rahman, G.; Javed, M. Stability for fractional order implicit ψ -Hilfer differential equations. *Math. Methods Appl. Sci.* **2021**, *45*, 2701–2712. [\[CrossRef\]](#)
- Benchohra, M.; Lazreg, J.E. Existence results for nonlinear implicit fractional dif-ferential equations. *Surv. Math. Its Appl.* **2014**, *9*, 79–92.
- Borisut, P.; Bantaojai, T. Implicit fractional differential equations with nonlocal frac-tional integral conditions. *Thai J. Math.* **2021**, *19*, 993–1003.
- Shabbir, A.S.; Shah, K.; Abdeljawad, T. Stability analysis for a class of implicit frac-tional differential equations involving Atangana–Baleanu fractional derivative. *Adv. Differ. Equ.* **2021**, *2021*, 395. [\[CrossRef\]](#)
- Benchohra, M.; Souid, M.S. Integrale solutions for implicit fractional order dif-ferential equations. *Arch. Math.* **2015**, *51*, 67–76.
- Benchohra, M.; Bouriaha, S. Existence and stability results for nonlinear boundary value problem for implicit differential equations of fractional order. *Moroc. J. Pure Appl. Anal.* **2017**, *1*, 22–37. [\[CrossRef\]](#)

20. Kucche, K.D.; Nieto, J.J.; Venkatesh, V. Theory of nonlinear implicit fractional differential equations. *Diff. Equ. Dyn. Syst.* **2016**, *38*, 1–17.
21. Nieto, J.J.; Ouahab, A.; Venkatesh, V. Implicit fractional differential equations via Liouville–Caputo derivative. *Mathematics* **2015**, *3*, 398–411. [[CrossRef](#)]
22. Atangana, A.; Baleanu, D. New fractional derivatives with nonlocal and nonsingular kernel: theory and application to heat transfer model. *Therm. Sci.* **2016**, *20*, 763–769. [[CrossRef](#)]
23. Srivastava, H.M.; Choi, J. *Zeta and q-Zeta Functions and Associated Series and Integrals*; Elsevier: Amsterdam, The Netherlands, 2012; pp. 1–140.
24. Helton, J.W.; Klep, I.; McCullough, S.; Schweighofer, M. Dilations, linear matrix inequalities, the matrix cube problem and beta distribution. *Memirs Am. Math. Soc.* **2019**, *257*, 1232.
25. Aono, Y.; Nguyen, P.Q.; Seito, T.; Shiketa, J. Lower Bounds on Lattice Enumeration with Extreme Pruning. In Proceedings of the 38th Annual International Cryptology Conference, IACR, Santa-Barbara, CA, USA, 19–23 August 2018. [[CrossRef](#)]
26. Shao, J.; Meng, F. Gronwall-Bellman Type Inequalities and Their Applications to Fractional Differential equations. *Abstr. Appl. Anal.* **2013**, *2013*, 217641. [[CrossRef](#)]
27. Di Giuseppe, E.; Moroni, M.; Caputo, M. Flux in porous Media with Memory: Models and Experiment. *Transp. Porous. Med.* **2010**, *83*, 479–500. [[CrossRef](#)]