

Article

Fixed Point Theorems via Orthogonal Convex Contraction in Orthogonal b -Metric Spaces and Applications

Gunasekaran Nallaselli ¹, Amani S. Baazeem ², Arul Joseph Gnanaprakasam ¹ , Gunaseelan Mani ³ , Khalil Javed ⁴, Eskandar Ameer ⁵ and Nabil Mlaiki ^{6,*} 

¹ Department of Mathematics, College of Engineering and Technology, Faculty of Engineering and Technology, SRM Institute of Science and Technology, SRM Nagar, Kanchipuram 603203, India

² Department of Mathematics and Statistics, College of Science, IMSIU (Imam Mohammed Ibn Saud Islamic University), P.O. Box 90950, Riyadh 11623, Saudi Arabia

³ Department of Mathematics, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Chennai 602105, India

⁴ Department of Mathematics and Statistics, International Islamic University Islamabad, Islamabad 04436, Pakistan

⁵ Department of Mathematics, Taiz University, Taiz P.O. Box 6803, Yemen

⁶ Department of Mathematics and Sciences, Prince Sultan University, P.O. Box 66833, Riyadh 11586, Saudi Arabia

* Correspondence: nmlaiki2012@gmail.com or nmlaiki@psu.edu.sa

Abstract: In this paper, we introduce the concept of orthogonal convex structure contraction mapping and prove some fixed point theorems on orthogonal b -metric spaces. We adopt an example to highlight the utility of our main result. Finally, we apply our result to examine the existence and uniqueness of the solution for the spring-mass system via an integral equation with a numerical example.

Keywords: b -metric space; O-set; O-sequence; b_{\perp} -metric space; fixed point; orthogonal convex



Citation: Nallaselli, G.; Baazeem, A.S.; Gnanaprakasam, A.J.; Mani, G.; Javed, K.; Ameer, E.; Mlaiki, N. Fixed Point Theorems via Orthogonal Convex Contraction in Orthogonal b -Metric Spaces and Applications. *Axioms* **2023**, *12*, 143. <https://doi.org/10.3390/axioms12020143>

Academic Editors: Boško Damjanović and Pradip Debnath

Received: 10 January 2023

Revised: 26 January 2023

Accepted: 28 January 2023

Published: 30 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Stefan Banach published his first known result in 1922, which is also possibly the most useful. It is referred to as the Banach contraction mapping principle. According to this theory, each contraction in a complete metric space has a distinct fixed point. It is helpful to note that this fixed point is also a singular fixed point for all iterations of the specified contractive mapping. Many writers generalised Banach's well-known discovery after 1922. On the subject, a lot of papers have been written. Two crucial generalizations were made:

(1) New relations (Kannan, Chatterje, Reich, Hardy-Rogers, Ćirić, ...) were used to bring new circumstances into the existing contractive relation.

(2) The axioms of metric space have been modified. As a result, numerous classes of new spaces are obtained. Visit papers [1–13] for additional information. Takahashi [14] initiated the notions of a convex structure and metric space in 1970 in addition to developing some of the fixed point theorems via his finding convex metric space. Goebel and Kirk [15] also looked at the iterative processes for nonexpansive mappings in the hyperbolic metric space, and in 1988, Xie [16] used Ishikawa's iteration approach to find fixed points for quasi-contraction mappings in convex metric spaces. Nonexpansive iterations in hyperbolic spaces were introduced in 1990 by Reich and Shafrir [17]. Mureşan et al. [18] presented the theory of some fixed point theorems for convex contraction mappings, the limit shadowing property, and Ulam-Hyers stability for the fixed point theorem in 2015. Latif et al. [19] established some approximate fixed point theorems via partial generalized convex contractions and partial generalized convex contractions of order 2 in the setting of α -complete metric spaces. Georgescu [20] studied iterated function systems consisting of generalized convex contractions on the framework of b -metric spaces. They proved the

generalization of Istratescu’s convex contraction fixed point theorem in the setting of complete strong b -metric spaces in 2017. Karaca et al. [21] proved fixed point theorems for the Reich contraction mapping in a convex b -metric space using the Mann iteration sequence in 2021. Also, they have the weak T-stability of the Mann iteration for this mapping in complete convex b -metric spaces. In 2021, Chen et al. [22] first introduced the concept of the convex graphical rectangular b -metric space (GR_bCMS) and obtained strong convergence theorems for these mappings in GR_bCMS under some suitable conditions. Following that, some works on the generalization of such classes of mappings in the setting of various spaces [23–29] appeared.

On the other hand, Gordji et al. [30] presented a new notion of orthogonality in metric spaces and illustrated the fixed point solution for contraction mappings in metric spaces using this new kind of orthogonality. They also showed how these results could be used to talk about the existence and uniqueness of a first-order ODE solution, even though the Banach contraction mapping principle does not work in this case. The fixed point in generalized orthogonal metric spaces was then demonstrated by Eshaghi Gordji and Habibi [31]. The idea of orthogonal F-contraction mappings was recently presented by Sawangsup et al. [32], who also demonstrated the fixed point theorems on orthogonal-complete metric spaces. The investigation of orthogonal contractive type mappings continued, with substantial findings made by numerous other researchers [33,34]. The goal of this study is to carry on these investigations. First, we discussed the novel notions of mappings of a orthogonal convex structural contraction on a orthogonal b -metric space. Then, we show the fixed point theorems on a orthogonal complete b -metric space and examples. We also present an application to resolve a spring-mass system and some examples for nonlinear integral equation of first kind with numerical solution to support of the obtained results.

2. Preliminaries

Throughout this paper, \mathcal{N} represents the set of positive integers, \mathfrak{R} denotes the set of all real numbers and \mathfrak{R}_0^+ is the set of non-negative reals.

Definition 1 ([3]). Let $Y \neq \emptyset$ and $\varrho \geq 1$ be a real number. A function $\delta_b : Y \times Y \rightarrow [0, \infty]$ is said to be a δ_b -metric on Y if the following conditions are satisfied:

- (1) $\delta_b(c, \sigma) = 0$ iff $c = \sigma$;
- (2) $\delta_b(c, \sigma) = \delta_b(\sigma, c)$, for all $c, \sigma \in Y$;
- (3) $\delta_b(c, \sigma) \leq \varrho[\delta_b(c, \mathfrak{d}) + \delta_b(\mathfrak{d}, \sigma)]$, for all $c, \sigma, \mathfrak{d} \in Y$.

The pair $(Y, \delta_b, \varrho \geq 1)$ is called b -metric space (shortly, b -MS).

The following are some examples and properties of a orthogonal set (or \mathfrak{D} -set) as initiated by Gordji et al. [30].

Definition 2 ([30]). Let $Y \neq \emptyset$. If a binary relation $\perp \subseteq Y \times Y$ satisfies the following stipulation:

$$\exists c_0 \in Y : (\forall c \in Y, c \perp c_0) \quad \text{or} \quad (\forall c \in Y, c_0 \perp c),$$

then it is called a orthogonal set (briefly \mathfrak{D} -set) and it is denoted by (Y, \perp) .

Example 1 ([30]). Let $Y = \mathfrak{R}_0^+$ and define $c \perp \sigma$ if $c\sigma \in \{c, \sigma\}$. Then, by letting $c_0 = 0$ or $c_0 = 1$, (Y, \perp) is an \mathfrak{D} -set.

Example 2. Let $Y = [0, \infty)$ and δ_b be a usual metric. Let $\top : Y \rightarrow Y$ be defined by $\top(c) = \frac{c}{2}$ if $c \neq 1$ else $\top(c) = 1$. Define now $c \perp \sigma$ if $c\sigma \leq \min\{c, \sigma\}$. Note that $0 \perp c$ for all $c \in Y$. Hence (Y, \perp) is an \mathfrak{O} -set.

At this point, it is important to remember some basic like, orthogonal sequence, orthogonal continuous, orthogonal complete, orthogonal metric space, orthogonal preserving, and weakly orthogonal preserving.

Definition 3 ([30]). A sequence $\{c_\vartheta\}$ of an \mathfrak{D} -set (Y, \perp) is called a orthogonal sequence (briefly, \mathfrak{D} -sequence) if

$$(\forall \vartheta \in \mathcal{N}, c_\vartheta \perp c_{\vartheta+1}) \quad \text{or} \quad (\forall \vartheta \in \mathcal{N}, c_{\vartheta+1} \perp c_\vartheta).$$

Definition 4 ([30]). We say that (Y, \perp, δ_b) is a orthogonal b -metric space (shortly, b_\perp -MS) if it contains an Definitions 1 and 2.

Definition 5 ([9]). Let $\{c_\vartheta\}$ be an \mathfrak{D} -sequence in (Y, \perp, δ_b) . Then:

1. We say that an \mathfrak{D} -sequence $\{c_\vartheta\}$ in b_\perp -MS (Y, \perp, δ_b) is convergent if $\exists c^* \in Y$ such that $\lim_{\vartheta \rightarrow \infty} \delta_b(c_\vartheta, c^*) = 0$.
2. We say that an \mathfrak{D} -sequence $\{c_\vartheta\}$ in (Y, \perp, δ_b) is a Cauchy \mathfrak{D} -sequence if for every $\varepsilon > 0$, \exists a $\vartheta_0 (> 0) \in \mathcal{N}$ such that $\delta_b(c_\vartheta, c_j) < \varepsilon \forall \vartheta, j > \vartheta_0$. i.e., $\lim_{\vartheta, j \rightarrow \infty} \delta_b(c_\vartheta, c_j) = 0$.
3. We say that (Y, \perp, δ_b) is \mathfrak{D} -complete b_\perp -metric space if every Cauchy \mathfrak{D} -sequence in Y is convergent.

Definition 6 ([30]). Let (Y, \perp, δ_b) be an b_\perp -MS. Then, we say that a function $\top : Y \rightarrow Y$ is a orthogonal continuous (or \perp -continuous) in $c \in Y$ if for each \mathfrak{D} -sequence $\{c_\vartheta\}$ of Y with $c_\vartheta \rightarrow c$ as $\vartheta \rightarrow \infty$, i.e., $\lim_{\vartheta \rightarrow \infty} \top(c_\vartheta) = \top(c)$. Also, we say that \top is \perp -continuous on Y if \top is \perp -continuous in each $c \in Y$.

Remark 1 ([30]). Every continuous mapping is \perp -continuous and the converse is not true.

Definition 7 ([30]). Let (Y, \perp) be an \mathfrak{D} -set. A mapping $\top : Y \rightarrow Y$ is said to be \perp -preserving if $\top c \perp \top \sigma$ whenever $c \perp \sigma$. Also $\top : Y \rightarrow Y$ is said to be weakly \perp -preserving if $\top(c) \perp \top(\sigma)$ or $\top(\sigma) \perp \top(c)$ whenever $c \perp \sigma$.

Definition 8 ([16]). Let $Y \neq \emptyset$ and $I = [0, 1]$. Let $\delta_b : Y \times Y \rightarrow \mathbb{R}_0^+$ be a function and let $\mathcal{V} : Y \times Y \times I \rightarrow Y$ be an \perp -continuous function. Then \mathcal{V} is called a orthogonal convex structure on Y if the conditions are met:

$$\delta_b(\vartheta, \mathcal{V}(c, \sigma; \xi)) \leq \xi \delta_b(\vartheta, c) + (1 - \xi) \delta_b(\vartheta, \sigma), \tag{1}$$

for each $\vartheta \in Y$ and $(c, \sigma; \xi) \in Y \times Y \times I$ with $\vartheta \perp c, \vartheta \perp \sigma$.

In the following section, we inspired and motivated the concepts of convex contraction and orthogonality. First, we define and illustrate orthogonal convex b_\perp -MS. We generalize and prove the fixed point theorem in the context of orthogonal convex b_\perp -MS using orthogonal convex contraction.

3. Main Results

Now, we define the notion of a orthogonal convex b_\perp -MS.

Definition 9. Let $\mathcal{V} : Y \times Y \times I \rightarrow Y$ be a orthogonal convex mapping structure defined on b_\perp -MS (Y, \perp, δ_b) with $\varrho \geq 1$ and $I = [0, 1]$. Then $(Y, \perp, \delta_b, \mathcal{V})$ is called a orthogonal convex b_\perp -MS.

Let $(Y, \perp, \delta_b, \mathcal{V})$ be a orthogonal convex b_\perp -MS and \top be a self-map on Y . Given below extension of iteration of Mann’s method into orthogonal convex b_\perp -MS.

$$\mu_{\vartheta+1} = \mathcal{V}(\mu_\vartheta, \top \mu_\vartheta; \alpha_\vartheta), \vartheta \in \mathcal{N},$$

where $\mu_\theta \in Y$ and $\alpha_\theta \in [0, 1]$. The \mathcal{D} -sequence $\{\mu_\theta\}$ is called the Mann's iteration \mathcal{D} -sequence for \mathbb{T} .

Now we'll look at some specific orthogonal convex b_\perp -MS example.

Example 3. Suppose $Y = [0, \infty)$ and $\epsilon = c_0 = (\frac{1}{2})$, then $\mathcal{S}_\epsilon[\frac{1}{2}] = \mathcal{I}$. Let $\mathbb{T} : Y \rightarrow Y$ be defined by

$$\mathbb{T}c = \begin{cases} c^{\frac{3}{4}}, & \text{if } c = \frac{1}{2}, \\ \frac{c}{6}, & \text{otherwise.} \end{cases}$$

Define now $c \perp \sigma$ if $c\sigma \leq \min\{c, \sigma\}$. Note that $0 \perp c$ for all $c \in Y$ and choosing a mapping $\delta_b : Y \times Y \rightarrow [0, +\infty)$ defined as

$$\delta_b(c, \sigma) = \begin{cases} (c - \sigma)^2, & \text{if both } c, \sigma \in [0, 1], \\ |c - \sigma|, & \text{otherwise.} \end{cases}$$

Demonstrate $\mathcal{V} : Y \times Y \times [0, 1] \rightarrow Y$ as

$$\mathcal{V}(c, \sigma; \alpha) = \alpha c + (1 - \alpha)\sigma$$

for all $c, \sigma \in Y$. Choose $c_\alpha = \mathcal{V}(c_{\alpha-1}, \mathbb{T}c_{\alpha-1}, \alpha_{\alpha-1})$, $b = 2$, and fix $\alpha_{\alpha-1} = \frac{1}{16} = (\frac{1}{4^2})$. Observe that $\alpha c + (1 - \alpha)\sigma \in [0, 1]$ whenever $c, \sigma \in [0, 1]$. Now, consider $\mathfrak{d}, c, \sigma \in Y$. Then we have two cases:

(i) $\mathfrak{d} \notin [0, 1]$, we get

$$\begin{aligned} \delta_b(\mathfrak{d}, \mathcal{V}(c, \sigma; \alpha)) &= |\alpha(\mathfrak{d} - c) + (1 - \alpha)(\mathfrak{d} - \sigma)| \\ &\leq |\alpha(\mathfrak{d} - c)| + |(1 - \alpha)(\mathfrak{d} - \sigma)| \\ &= \alpha\delta_b(\mathfrak{d}, c) + (1 - \alpha)\delta_b(\mathfrak{d}, \sigma). \end{aligned} \tag{2}$$

(ii) When $\mathfrak{d} \in [0, 1]$. We have the following sub cases:

(a) If both $c, \sigma \in [0, 1]$, then obviously $\mathcal{V}(c, \sigma; \alpha) = \alpha c + (1 - \alpha)\sigma \in [0, 1]$, and hence

$$\begin{aligned} \delta_b(\mathfrak{d}, \mathcal{V}(c, \sigma; \alpha)) &= [b(\mathfrak{d} - c) + (1 - b)(\mathfrak{d} - \sigma)]^2 \\ &\leq [b|\mathfrak{d} - c| + (1 - b)|\mathfrak{d} - \sigma|]^2 \\ &= (b|\mathfrak{d} - c|)^2 + ((1 - b)|\mathfrak{d} - \sigma|)^2 + 2b(1 - b)|\mathfrak{d} - c||\mathfrak{d} - \sigma| \\ &\leq (b|\mathfrak{d} - c|)^2 + ((1 - b)|\mathfrak{d} - \sigma|)^2 + b(1 - b)((\mathfrak{d} - c)^2 + (\mathfrak{d} - \sigma)^2) \\ &= b(\mathfrak{d} - c)^2 + (1 - b)(\mathfrak{d} - \sigma)^2 \\ &= b\delta_b(\mathfrak{d}, c) + (1 - b)\delta_b(\mathfrak{d}, \sigma). \end{aligned} \tag{3}$$

(b) If only one of c and σ is in $[0, 1]$, say c is in $[0, 1]$, then obviously $\mathcal{V}(c, \sigma; \alpha) = \alpha c + (1 - \alpha)\sigma \notin [0, 1]$, and hence,

$$\begin{aligned} \delta_b(\mathfrak{d}, \mathcal{V}(c, \sigma; \alpha)) &= |b(\mathfrak{d} - c) + (1 - b)(\mathfrak{d} - \sigma)| \\ &\leq |b(\mathfrak{d} - c)| + |(1 - b)(\mathfrak{d} - \sigma)| \\ &= b(\mathfrak{d} - c)^2 + |(1 - b)(\mathfrak{d} - \sigma)| \\ &= b\delta_b(\mathfrak{d}, c) + (1 - b)\delta_b(\mathfrak{d}, \sigma). \end{aligned} \tag{4}$$

The same can be done for $\sigma \in [0, 1]$ and c not in $[0, 1]$.

(c) If both $c, \sigma \notin [0, 1]$, then obviously $\mathcal{V}(c, \sigma; \alpha) = \alpha c + (1 - \alpha)\sigma \notin [0, 1]$

$$\begin{aligned} \delta_b(\mathfrak{d}, \mathcal{V}(c, \sigma; \alpha)) &= |b(\mathfrak{d} - c) + (1 - b)(\mathfrak{d} - \sigma)| \\ &\leq |b(\mathfrak{d} - c)| + |(1 - b)(\mathfrak{d} - \sigma)| \\ &= b\delta_b(\mathfrak{d}, c) + (1 - b)\delta_b(\mathfrak{d}, \sigma). \end{aligned} \tag{5}$$

From all the possible cases, it is clear that $(Y, \perp, \delta_b, \mathcal{V})$ is a orthogonal convex b_\perp -MS with $b = 2$.

Example 4. Let $Y = \mathfrak{R}$ and $\delta_b : Y \times Y \rightarrow [0, +\infty)$ be a function defined by

$$\delta_b(c, \sigma) = |c - \sigma|^\alpha, \alpha > 1,$$

for all $c, \sigma \in Y$. Define the binary relation \perp on Y by $c \perp \sigma$ if $c\sigma \leq (c \vee \sigma)$, where $c \vee \sigma = c$ or σ . Then, (Y, δ_b) is an O-complete b_\perp -MS. Let $\mathcal{V} : Y \times Y \times \{\frac{1}{2}\} \rightarrow Y$ be a function defined by

$$\mathcal{V}(c, \sigma; a) = \frac{c + \sigma}{2}.$$

Then, $(Y, \perp, \delta_b, \mathcal{V})$ is a orthogonal convex b_\perp -MS with $\rho = 2^{\alpha-1}$. Now, in the usual sense, $(Y, \perp, \delta_b, \mathcal{V})$ is not a orthogonal metric space.

Indeed, given any $\vartheta, j \in [0, +\infty)$, and $\alpha \geq 1$, inequality

$$(\vartheta + j)^\alpha \leq 2^{\alpha-1}(\vartheta^\alpha + j^\alpha)$$

exists, we conclude that (Y, \perp, δ_b) is an b_\perp -MS with $\rho = 2^\alpha - 1$. Now, clear that \mathcal{V} satisfies Equation (1). For each $\vartheta, c, \sigma \in Y$ with $c \perp \sigma \implies \vartheta \perp c, \vartheta \perp \sigma$, we get

$$\begin{aligned} \delta_b(\vartheta, \mathcal{V}(c, \sigma; a)) &= \left| \vartheta - \left[\frac{c + \sigma}{2} \right] \right|^\alpha \\ &\leq 2^{\alpha-1} \left[2^{-\alpha} |\vartheta - c|^\alpha + 2^{-\alpha} |\vartheta - \sigma|^\alpha \right] \\ &= 2^{-1} \left[|\vartheta - c|^\alpha + |\vartheta - \sigma|^\alpha \right] \\ &= a\delta_b(\vartheta, c) + (1 - a)\delta_b(\vartheta, \sigma), \end{aligned}$$

so $(Y, \perp, \delta_b, \mathcal{V})$ be a orthogonal convex b_\perp -MS with $\rho = 2^{\alpha-1}$. However, because δ_b does not satisfy triangle inequality, $(Y, \perp, \delta_b, \mathcal{V})$ is not a orthogonal metric space in the usual sense. Now, take $\alpha = 2$, we get

$$\delta_b(3, 5) = 4 > \delta_b(3, 4) + \delta_b(4, 5) = 2.$$

Using Mann’s iteration algorithm, we will now demonstrate Banach’s contraction principle for O-complete convex b_\perp -MSs.

Theorem 1. Let $(Y, \perp, \delta_b, \mathcal{V}, \rho > 1)$ be an O-complete convex b_\perp -MS and $T : Y \rightarrow Y$ be a contractive self-map on Y . Suppose that there exists $b \in [0, 1)$ such that the following assertions hold:

1. T is \perp -preserving,
2. For all $c, \sigma \in Y$ with $c \perp \sigma$, $[\delta_b(Tc, T\sigma) > 0, \delta_b(Tc, T\sigma) \leq k\delta_b(c, \sigma)]$.

Take $c_0 \in Y$ such that $\delta_b(c_0, Tc_0) = \mathcal{K} < \infty$ and $c_\vartheta = \mathcal{V}(c_{\vartheta-1}, Tc_{\vartheta-1}; a_{\vartheta-1})$, here $0 \leq a_{\vartheta-1} < 1$ with $\vartheta \in \mathcal{N}$. If $k\rho^4 < 1$ and $0 < a_{\vartheta-1} < \frac{\rho^{\frac{1}{4}-k}}{1-k}$ for each $\vartheta \in \mathcal{N}$; then, T has a unique fixed point in Y .

Proof. Since (Y, \perp) is an \mathfrak{D} -set,

$$\exists c_0 \in Y : (\forall c \in Y, c \perp c_0) \quad \text{or} \quad (\forall c \in Y, c_0 \perp c).$$

It follows that $c_0 \perp Tc_0$ or $Tc_0 \perp c_0$. Let $c_1 := Tc_0, c_2 := Tc_1, \dots, c_{\vartheta+1} := Tc_\vartheta, \forall \vartheta \in \mathcal{N} \cup \{0\}$.

If $c_\vartheta = c_{\vartheta+1}$ for each $\vartheta \in \mathcal{N} \cup \{0\}$, it follows that c_ϑ is a fixed point of \top . Postulate that $c_\vartheta \neq c_{\vartheta+1} \forall \vartheta \in \mathcal{N} \cup \{0\}$. Thus, we have $\delta_b(c_\vartheta, c_{\vartheta+1}) > 0$ for all $\vartheta \in \mathcal{N} \cup \{0\}$. By condition (1), we get

$$c_\vartheta \perp c_{\vartheta+1} \quad \text{or} \quad c_{\vartheta+1} \perp c_\vartheta$$

$\forall \vartheta \in \mathcal{N} \cup \{0\}$. Hence $\{c_\vartheta\}$ is an \mathcal{O} -sequence.

For any $\vartheta \in \mathcal{N}$, there exists

$$\delta_b(c_\vartheta, c_{\vartheta+1}) = \delta_b(c_\vartheta, \mathcal{V}(c_\vartheta, \top c_\vartheta; \alpha_\vartheta)) \leq (1 - \alpha_\vartheta) \delta_b(c_\vartheta, \top c_\vartheta)$$

and

$$\begin{aligned} \delta_b(c_\vartheta, \top c_\vartheta) &\leq \varrho \delta_b(c_\vartheta, \top c_{\vartheta-1}) + \varrho \delta_b(\top c_{\vartheta-1}, \top c_\vartheta) \\ &\leq \varrho \delta_b(\mathcal{V}(c_{\vartheta-1}, \top c_{\vartheta-1}; \alpha_{\vartheta-1}), \top c_{\vartheta-1}) + \varrho \mathbb{k} \delta_b(c_{\vartheta-1}, c_\vartheta) \\ &\leq \varrho [\alpha_{\vartheta-1} \delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}) + \mathbb{k}(1 - \alpha_{\vartheta-1}) \delta_b(c_{\vartheta-1}, \top c_{\vartheta-1})] \\ &= \varrho [\alpha_{\vartheta-1} + \mathbb{k}(1 - \alpha_{\vartheta-1})] \delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}). \end{aligned}$$

Let $\xi_{\vartheta-1} = \varrho [\alpha_{\vartheta-1} + \mathbb{k}(1 - \alpha_{\vartheta-1})]$. Combining from the above with $\mathbb{k} \varrho^4 < 1$ and $0 < \alpha_{\vartheta-1} < \frac{\varrho^{\frac{1}{4}-\mathbb{k}}}{1-\mathbb{k}}$ holding for each $\vartheta \in \mathcal{N}$, we get

$$\delta_b(c_\vartheta, \top c_\vartheta) \leq \xi_{\vartheta-1} \delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}) < \frac{1}{\varrho^3} \delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}), \tag{6}$$

which shows that $\{\delta_b(c_\vartheta, \top c_\vartheta)\}$ is decreasing \mathcal{O} -sequence of non-negative reals. Hence, $\exists \lambda \geq 0$ such that

$$\lim_{\vartheta \rightarrow \infty} \delta_b(c_\vartheta, \top c_\vartheta) = \lambda.$$

We prove $\lambda = 0$. Assume that $\lambda > 0$. Taking $\vartheta \rightarrow \infty$ in Equation (6), we get

$$\lambda \leq \frac{1}{\varrho^3} \lambda < \lambda,$$

a contradiction. Hence, we obtain $\lambda = 0$. Next, we get

$$\delta_b(c_\vartheta, c_{\vartheta+1}) \leq (1 - \alpha_\vartheta) \delta_b(c_\vartheta, \top c_\vartheta),$$

which implies that $\lim_{\vartheta \rightarrow \infty} \delta_b(c_\vartheta, c_{\vartheta+1}) = 0$. Next, prove that $\{c_\vartheta\}$ is a Cauchy \mathcal{O} -sequence. Contrary, we assume an \mathcal{O} -sequence $\{c_\vartheta\}$ is not a Cauchy, then $\exists \varepsilon_0 > 0$ and the subsequences $\{c_{s(\omega)}\}$ and $\{c_{t(\omega)}\}$ of $\{c_\vartheta\}$, such that $s(\omega)$ is the smallest number with $s(\omega) > t(\omega) > \omega$,

$$\delta_b(c_{s(\omega)}, c_{t(\omega)}) \geq \varepsilon_0$$

and

$$\delta_b(c_{s(\omega-1)}, c_{t(\omega)}) < \varepsilon_0.$$

Then, we conclude

$$\varepsilon_0 \leq \delta_b(c_{s(\omega)}, c_{t(\omega)}) \leq \varrho [\delta_b(c_{s(\omega)}, c_{t(\omega)+1}) + \delta_b(c_{t(\omega)+1}, c_{t(\omega)})],$$

which implies that $\frac{\varepsilon_0}{\varrho} \leq \limsup_{\omega \rightarrow \infty} \delta_b(c_{s(\omega)}, c_{t(\omega)+1})$.

Note that

$$\begin{aligned}
 \delta_b(\mathbf{c}_{\mathfrak{s}(\omega)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) &= \delta_b\left(\mathcal{V}(\mathbf{c}_{\mathfrak{s}(\omega-1)}, \top \mathbf{c}_{\mathfrak{s}(\omega-1)}; \mathbf{a}_{\mathfrak{s}(\omega-1)}), \mathbf{c}_{\mathfrak{t}(\omega)+1}\right) \\
 &\leq \mathbf{a}_{\mathfrak{s}(\omega-1)} \delta_b(\mathbf{c}_{\mathfrak{s}(\omega-1)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) + (1 - \mathbf{a}_{\mathfrak{s}(\omega-1)}) \delta_b(\top \mathbf{c}_{\mathfrak{s}(\omega-1)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \\
 &\leq \mathbf{a}_{\mathfrak{s}(\omega-1)} \delta_b(\mathbf{c}_{\mathfrak{s}(\omega-1)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) + (1 - \mathbf{a}_{\mathfrak{s}(\omega-1)}) \varrho \left[\delta_b(\top \mathbf{c}_{\mathfrak{s}(\omega-1)}, \top \mathbf{c}_{\mathfrak{t}(\omega)+1}) \right. \\
 &\quad \left. + \delta_b(\top \mathbf{c}_{\mathfrak{t}(\omega)+1}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \right] \\
 &\leq \mathbf{a}_{\mathfrak{s}(\omega-1)} \delta_b(\mathbf{c}_{\mathfrak{s}(\omega-1)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) + (1 - \mathbf{a}_{\mathfrak{s}(\omega-1)}) \varrho \left[\mathbb{K} \delta_b(\mathbf{c}_{\mathfrak{s}(\omega-1)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \right. \\
 &\quad \left. + \delta_b(\top \mathbf{c}_{\mathfrak{t}(\omega)+1}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \right] \\
 &= [\mathbf{a}_{\mathfrak{s}(\omega-1)} + (1 - \mathbf{a}_{\mathfrak{s}(\omega-1)}) \varrho \mathbb{K}] \delta_b(\mathbf{c}_{\mathfrak{s}(\omega-1)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \\
 &\quad + (1 - \mathbf{a}_{\mathfrak{s}(\omega-1)}) \varrho \delta_b(\top \mathbf{c}_{\mathfrak{t}(\omega)+1}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \\
 &< \varrho [\mathbf{a}_{\mathfrak{s}(\omega-1)} \varrho \mathbf{a}_{\mathfrak{s}(\omega-1)} + (1 - \mathbf{a}_{\mathfrak{s}(\omega-1)}) \varrho \mathbb{K}] \left(\delta_b(\mathbf{c}_{\mathfrak{s}(\omega-1)}, \mathbf{c}_{\mathfrak{t}(\omega)}) \right. \\
 &\quad \left. + \delta_b(\mathbf{c}_{\mathfrak{t}(\omega)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \right) + (1 - \mathbf{a}_{\mathfrak{s}(\omega-1)}) \varrho \delta_b(\top \mathbf{c}_{\mathfrak{t}(\omega)+1}, \mathbf{c}_{\mathfrak{t}(\omega)+1}),
 \end{aligned}$$

we obtain $\frac{1}{\varrho} \varepsilon_0 \leq \lim_{\omega \rightarrow \infty} \sup \delta_b(\mathbf{c}_{\mathfrak{s}(\omega)}, \mathbf{c}_{\mathfrak{t}(\omega)+1}) \leq \varrho^2 \frac{1}{\varrho^4} \varepsilon_0$, a contradiction. Thus $\{\mathbf{c}_\vartheta\}$ is a Cauchy O-sequence in Y . By the O-completeness of Y , $\exists \mathbf{c}^* \in Y$ such that $\lim_{\vartheta \rightarrow \infty} \delta_b(\mathbf{c}_\vartheta, \mathbf{c}^*) = 0$. Next, prove that \mathbf{c}^* is a fixed point of \top . Consider

$$\begin{aligned}
 \delta_b(\mathbf{c}^*, \top \mathbf{c}^*) &\leq \varrho [\delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + \delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}^*)] \\
 &\leq \varrho \delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + \varrho^2 [\delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}_\vartheta) + \delta_b(\top \mathbf{c}_\vartheta, \top \mathbf{c}^*)] \\
 &= \varrho \delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + \varrho^2 \delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}_\vartheta) + \varrho^2 \mathbb{K} \delta_b(\mathbf{c}_\vartheta, \mathbf{c}^*).
 \end{aligned}$$

Letting $\vartheta \rightarrow \infty$, we conclude that $\delta_b(\mathbf{c}^*, \top \mathbf{c}^*) = 0$ which proves that $\top \mathbf{c}^* = \mathbf{c}^*$. Hence, \mathbf{c}^* is a fixed point of \top .

Now, prove the uniqueness part. Let $\mathbf{c}^*, \mathfrak{z}$ be two distinct fixed points of \top and postulate that $\top^\vartheta \mathbf{c}^* = \mathbf{c}^* \neq \mathfrak{z} = \top^\vartheta \mathfrak{z} \forall \vartheta \in \mathcal{N}$. From definition 2.2, we get

$$(\mathbf{c}_0 \perp \mathbf{c}^* \text{ and } \mathbf{c}_0 \perp \mathfrak{z}) \text{ or } (\mathbf{c}^* \perp \mathbf{c}_0 \text{ and } \mathfrak{z} \perp \mathbf{c}_0).$$

By condition (1), we have

$$(\top^\vartheta \mathbf{c}_0 \perp \top^\vartheta \mathbf{c}^* \text{ and } \top^\vartheta \mathbf{c}_0 \perp \mathfrak{z}) \text{ or } (\top^\vartheta \mathbf{c}^* \perp \mathbf{c}_0 \text{ and } \top^\vartheta \mathfrak{z} \perp \mathbf{c}_0)$$

for all $\vartheta \in \mathcal{N}$. Now

$$\delta_b(\mathbf{c}^*, \mathfrak{z}) = \delta_b(\top^\vartheta \mathbf{c}^*, \top^\vartheta \mathfrak{z}) \leq \varrho [\delta_b(\top^\vartheta \mathbf{c}^*, \top^\vartheta \mathbf{c}_0) + \delta_b(\top^\vartheta \mathbf{c}_0, \top^\vartheta \mathfrak{z})].$$

As $\vartheta \rightarrow \infty$, we obtain $\delta_b(\mathbf{c}^*, \mathfrak{z}) \leq 0$. Thus, $\mathbf{c}^* = \mathfrak{z}$. Hence, \top has a unique fixed point in Y . \square

We demonstrate an example illustrating the Theorem 1.

Example 5. Let $Y = \mathfrak{R}^+ \cup \{0\}$, $\top \mathbf{c} = \frac{\mathbf{c}}{5} \forall \mathbf{c} \in Y$. Define a function $\delta_b : Y \times Y \rightarrow [0, +\infty)$ by the formula

$$\delta_b(\mathbf{c}, \sigma) = (\mathbf{c} - \sigma)^2, \text{ for all } \mathbf{c}, \sigma \in Y.$$

Define the binary relation \perp on Y by $c \perp \sigma$ if $c\sigma \leq (c \vee \sigma)$, where $c \vee \sigma = c$ or σ . Then, (Y, δ_b) is an O -complete b_{\perp} -MS. The mapping $\mathcal{V} : Y \times Y \times [0, 1] \rightarrow Y$ is defined as

$$\mathcal{V}(c, \sigma; \alpha) = \alpha c + (1 - \alpha)\sigma, \text{ for all } c, \sigma \in Y.$$

Set $\mathbb{k} = \frac{1}{1+2^4}$ and $c_{\vartheta} = \mathcal{V}(c_{\vartheta-1}, \top c_{\vartheta-1}; \alpha_{\vartheta-1})$, where $c_0 = 1$ and $\alpha_{\vartheta-1} = \frac{1}{2^4} - \mathbb{k}$. Then, $(Y, \perp, \delta_b, \mathcal{V})$ is an O -complete convex b_{\perp} -MS with $q = 2$, and \top has a unique fixed point in Y .

It shows that (Y, \perp, δ_b) is an b_{\perp} -MS with $q = 2$, from Example 4. For each $\mathfrak{d}, c, \sigma \in Y$ with $c \perp \sigma, \mathfrak{d} \perp c, \mathfrak{d} \perp \sigma$, we obtain

$$\begin{aligned} \delta_b(\mathfrak{d}, \mathcal{V}(c, \sigma; \alpha)) &= [\alpha(\mathfrak{d} - c) + (1 - \alpha)(\mathfrak{d} - \sigma)]^2 \\ &\leq [\alpha|\mathfrak{d} - c| + (1 - \alpha)|\mathfrak{d} - \sigma|]^2 \\ &= (\alpha|\mathfrak{d} - c|)^2 + ((1 - \alpha)|\mathfrak{d} - \sigma|)^2 + 2\alpha(1 - \alpha)|\mathfrak{d} - c||\mathfrak{d} - \sigma| \\ &\leq (\alpha|\mathfrak{d} - c|)^2 + ((1 - \alpha)|\mathfrak{d} - \sigma|)^2 + \alpha(1 - \alpha)(|\mathfrak{d} - c|^2 + |\mathfrak{d} - \sigma|^2) \\ &= \alpha(\mathfrak{d} - c)^2 + (1 - \alpha)(\mathfrak{d} - \sigma)^2 \\ &= \alpha\delta_b(\mathfrak{d}, c) + (1 - \alpha)\delta_b(\mathfrak{d}, \sigma). \end{aligned}$$

Hence, $(Y, \delta_b, \mathcal{V})$ is a orthogonal convex b_{\perp} -MS with $q = 2$. It is easy to see that \top satisfies $\delta_b(\top c, \top \sigma) = \frac{1}{25}\delta_b(c, \sigma) \leq \mathbb{k}\delta_b(c, \sigma)$, where $\mathbb{k} = \frac{1}{17}$. We choose $c_0 \in Y \setminus \{0\}$. Combining with $c_{\vartheta} = \mathcal{V}(c_{\vartheta-1}, \top c_{\vartheta-1}; \alpha_{\vartheta-1})$ and $\top c = \frac{c}{5}$, we have

$$c_{\vartheta} = \alpha_{\vartheta-1}c_{\vartheta-1} + (1 - \alpha_{\vartheta-1})\top c_{\vartheta-1} = \left(\frac{1}{5} + \frac{4}{5}\alpha_{\vartheta-1}\right)c_{\vartheta-1},$$

and

$$c_{\vartheta-1} = \left(\frac{1}{5} + \frac{4}{5}\alpha_{\vartheta-2}\right)c_{\vartheta-2}, c_{\vartheta-2} = \left(\frac{1}{5} + \frac{4}{5}\alpha_{\vartheta-3}\right)c_{\vartheta-3}, \dots, c_1 = \left(\frac{1}{5} + \frac{4}{5}\alpha_0\right)c_0.$$

Since $\alpha_{\vartheta-1} = \frac{1}{2^4} - \mathbb{k}$ for all $\vartheta \in \mathcal{N}$, we obtain

$$c_{\vartheta} = \left(\frac{69}{340}\right)^{\vartheta}c_0 \text{ and } \top c_{\vartheta} = \frac{1}{5} \cdot \left(\frac{69}{340}\right)^{\vartheta}c_0.$$

Letting $\vartheta \rightarrow \infty$, we have $c_{\vartheta} \rightarrow 0 \in Y$ and $\top c_{\vartheta} \rightarrow 0 \in Y$. Hence, 0 is a fixed point of \top in Y . Next, prove \top has a unique fixed point. Postulate that $c^*, \mathfrak{z} \in Y$ are two distinct fixed points of \top . Then,

$$\delta_b(c^*, \mathfrak{z}) > 0, \delta_b(c^*, \mathfrak{z}) = \delta_b(\top c^*, \top \mathfrak{z}) = \delta_b\left(\frac{1}{5}c^*, \frac{1}{5}\mathfrak{z}\right) = \frac{1}{25}\delta_b(c^*, \mathfrak{z}),$$

a contradiction. Hence, \top has a unique fixed point 0 in Y .

We prove the Kannan theorem for an O -complete convex b_{\perp} -MS.

Theorem 2. Let $(Y, \perp, \delta_b, \mathcal{V})$ be an O -complete convex b_{\perp} -MS with constant $q > 1$ and let $\top : Y \rightarrow Y$ be a contraction mapping. Suppose $\exists b \in [0, 1)$ such that the following axioms hold:

1. \top is \perp -preserving,
2. For all $c, \sigma \in Y$ with $c \perp \sigma$, and for some $\omega \in [0, \frac{1}{2})$,

$$\delta_b(\top c, \top \sigma) > 0, \delta_b(\top c, \top \sigma) \leq \omega[\delta_b(c, \top c) + \delta_b(\sigma, \top \sigma)]. \tag{7}$$

Take $c_0 \in Y$ such that $\delta_b(c_0, \top c_0) = \mathcal{K} < \infty$ and $c_{\vartheta} = \mathcal{V}(c_{\vartheta-1}, \top c_{\vartheta-1}; \alpha_{\vartheta-1})$ for $\vartheta \in \mathcal{N}$ and $\alpha_{\vartheta-1} \in (0, \frac{1}{4q^2}]$. If $\omega \in (0, \frac{1}{4q^2}]$, then \top has a unique fixed point in Y .

Proof. Since (Y, \perp) is an \mathfrak{D} -set,

$$\exists c_0 \in Y : (\forall c \in Y, c \perp c_0) \quad \text{or} \quad (\forall c \in Y, c_0 \perp c).$$

It follows that $c_0 \perp \top c_0$ or $\top c_0 \perp c_0$. Let $c_1 := \top c_0, c_2 := \top c_1, \dots, c_{\vartheta+1} := \top c_{\vartheta}$, for all $\vartheta \in \mathcal{N} \cup \{0\}$. If $c_{\vartheta} = c_{\vartheta+1}$ for any $\vartheta \in \mathcal{N} \cup \{0\}$, then it is clear that c_{ϑ} is a fixed point of \top . Postulate that $c_{\vartheta} \neq c_{\vartheta+1} \forall \vartheta \in \mathcal{N} \cup \{0\}$. Thus, we have $\delta_b(c_{\vartheta}, c_{\vartheta+1}) > 0 \forall \vartheta \in \mathcal{N} \cup \{0\}$. By condition (1), we get

$$c_{\vartheta} \perp c_{\vartheta+1} \quad \text{or} \quad c_{\vartheta+1} \perp c_{\vartheta}$$

for all $\vartheta \in \mathcal{N} \cup \{0\}$. Hence $\{c_{\vartheta}\}$ is an \mathfrak{D} -sequence.

For any $\vartheta \in \mathcal{N}$, we get

$$\delta_b(c_{\vartheta}, c_{\vartheta+1}) = \delta_b(c_{\vartheta}, \mathcal{V}(c_{\vartheta}, \top c_{\vartheta}; a_{\vartheta})) \leq (1 - a_{\vartheta})\delta_b(c_{\vartheta}, \top c_{\vartheta}) \tag{8}$$

and

$$\begin{aligned} \delta_b(c_{\vartheta}, \top c_{\vartheta}) &= \delta_b(\mathcal{V}(c_{\vartheta-1}, \top c_{\vartheta-1}; a_{\vartheta-1}), \top c_{\vartheta}) \\ &\leq a_{\vartheta-1}\delta_b(c_{\vartheta-1}, \top c_{\vartheta}) + (1 - a_{\vartheta-1})\delta_b(\top c_{\vartheta-1}, \top c_{\vartheta}) \\ &\leq \varrho a_{\vartheta-1}\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}) + \varrho a_{\vartheta-1}\delta_b(\top c_{\vartheta-1}, \top c_{\vartheta}) + \delta_b(\top c_{\vartheta-1}, \top c_{\vartheta}) \\ &\leq \varrho a_{\vartheta-1}\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}) + (\varrho a_{\vartheta-1} + 1)\omega[\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}) + \delta_b(c_{\vartheta}, \top c_{\vartheta})] \\ &= (\varrho a_{\vartheta-1} + \varrho a_{\vartheta-1}\omega + \omega)\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}) + (\varrho a_{\vartheta-1}\omega + \omega)\delta_b(c_{\vartheta}, \top c_{\vartheta}); \end{aligned}$$

i.e., $[1 - (\varrho a_{\vartheta-1}\omega + \omega)]\delta_b(c_{\vartheta}, \top c_{\vartheta}) \leq (\varrho a_{\vartheta-1} + \varrho a_{\vartheta-1}\omega + \omega)\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1})$.

Since $\varrho a_{\vartheta-1}\omega + \omega \leq (\frac{1}{4\varrho} + 1)\omega < \frac{4}{5} \cdot \frac{1}{4\varrho^2} < 1$, then

$$\delta_b(c_{\vartheta}, \top c_{\vartheta}) \leq \frac{\varrho a_{\vartheta-1} + \varrho a_{\vartheta-1}\omega + \omega}{1 - (\varrho a_{\vartheta-1}\omega + \omega)}\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}). \tag{9}$$

Denote $\tilde{\zeta}_{\vartheta-1} = \frac{\varrho a_{\vartheta-1} + \varrho a_{\vartheta-1}\omega + \omega}{1 - \varrho a_{\vartheta-1}\omega + \omega}$ for $\vartheta \in \mathcal{N}$. We deduce that

$$\begin{aligned} \tilde{\zeta}_{\vartheta-1} &= \frac{\varrho a_{\vartheta-1} + \varrho a_{\vartheta-1}\omega + \omega}{1 - \varrho a_{\vartheta-1}\omega + \omega} \\ &< \frac{\frac{5}{4}}{1 - \varrho a_{\vartheta-1}\omega + \omega} - 1 \\ &< \frac{\frac{5}{4}}{1 - \frac{5}{4} \cdot \frac{1}{4\varrho^2}} - 1 < \frac{9}{11}. \end{aligned}$$

Combining the above two inequalities, we get

$$\delta_b(c_{\vartheta}, \top c_{\vartheta}) \leq \tilde{\zeta}_{\vartheta-1}\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}) < \frac{9}{11}\delta_b(c_{\vartheta-1}, \top c_{\vartheta-1}), \tag{10}$$

which shows that $\{\delta_b(c_{\vartheta}, \top c_{\vartheta})\}$ is decreasing O-sequence of non-negative reals. Hence, $\exists \lambda \geq 0$ such that $\lim_{\vartheta \rightarrow \infty} \delta_b(c_{\vartheta}, \top c_{\vartheta}) = \lambda$. Prove that $\lambda = 0$. Let $\lambda > 0$. Taking $\vartheta \rightarrow \infty$ in Equation (10), we have $\lambda \leq \frac{9}{11}\lambda < \lambda$, a contradiction. Hence, $\lambda = 0$; i.e., $\lim_{\vartheta \rightarrow \infty} \delta_b(c_{\vartheta}, \top c_{\vartheta}) = 0$. Moreover, by inequality (8), we obtain $\delta_b(c_{\vartheta}, c_{\vartheta+1}) \leq (1 - a_{\vartheta})\delta_b(c_{\vartheta}, \top c_{\vartheta}) < \delta_b(c_{\vartheta}, \top c_{\vartheta})$, which shows that $\lim_{\vartheta \rightarrow \infty} \delta_b(c_{\vartheta}, c_{\vartheta+1}) = 0$. Next, prove that an O-sequence $\{c_{\vartheta}\}$ is Cauchy. Contrary, we assume an O-sequence $\{c_{\vartheta}\}$ is not Cauchy, then $\exists \varepsilon_0 > 0, \{c_{s(\alpha)}\}$ and $\{c_{t(\alpha)}\}$ are the sub sequences of $\{c_{\vartheta}\}$ such that $s(\alpha)$ is not greatest number with $s(\alpha) > t(\alpha) > l$,

$$\delta_b(c_{s(\alpha)}, c_{t(\alpha)}) \geq \varepsilon_0$$

and

$$\delta_b(\mathbf{c}_{s(\alpha)-1}, \mathbf{c}_{t(\alpha)}) < \varepsilon_0.$$

Then, we conclude that

$$\varepsilon_0 \leq \delta_b(\mathbf{c}_{s(\alpha)}, \mathbf{c}_{t(\alpha)}) \leq \varrho[\delta_b(\mathbf{c}_{s(\alpha)}, \mathbf{c}_{t(\alpha)+1}) + db(\mathbf{c}_{t(\alpha)+1}, \mathbf{c}_{t(\alpha)})],$$

which implies that

$$\frac{\varepsilon_0}{\varrho} \leq \limsup_{\alpha \rightarrow \infty} \delta_b(\mathbf{c}_{s(\alpha)}, \mathbf{c}_{t(\alpha)+1}).$$

Note that

$$\begin{aligned} \delta_b(\mathbf{c}_{s(\alpha)}, \mathbf{c}_{t(\alpha)+1}) &= \delta_b\left(\mathcal{V}(\mathbf{c}_{s(\alpha)-1}, \top \mathbf{c}_{s(\alpha)-1}; \mathbf{a}_{s(\alpha)-1}, \mathbf{c}_{t(\alpha)+1})\right) \\ &\leq \mathbf{a}_{s(\alpha)-1} \delta_b(\mathbf{c}_{s(\alpha)-1}, \mathbf{c}_{t(\alpha)+1}) + (1 - \mathbf{a}_{s(\alpha)-1}) \delta_b(\top \mathbf{c}_{s(\alpha)-1}, \mathbf{c}_{t(\alpha)+1}) \\ &\leq \mathbf{a}_{s(\alpha)-1} \delta_b(\mathbf{c}_{s(\alpha)-1}, \mathbf{c}_{t(\alpha)+1}) + (1 - \mathbf{a}_{s(\alpha)-1}) \varrho \left[\delta_b(\top \mathbf{c}_{s(\alpha)-1}, \top \mathbf{c}_{t(\alpha)+1}) \right. \\ &\quad \left. + \delta_b(\top \mathbf{c}_{t(\alpha)+1}, \mathbf{c}_{t(\alpha)+1}) \right] \\ &\leq \mathbf{a}_{s(\alpha)-1} \delta_b(\mathbf{c}_{s(\alpha)-1}, \mathbf{c}_{t(\alpha)+1}) + (1 - \mathbf{a}_{s(\alpha)-1}) \varrho \left[\omega \delta_b(\mathbf{c}_{s(\alpha)-1}, \top \mathbf{c}_{s(\alpha)-1}) \right. \\ &\quad \left. + (\omega + 1) \delta_b(\top \mathbf{c}_{t(\alpha)+1}, \mathbf{c}_{t(\alpha)+1}) \right] \text{ (for some } \omega \in \left[0, \frac{1}{2}\right) \text{ satisfying (3))} \\ &\leq \mathbf{a}_{s(\alpha)-1} \left[\varrho \delta_b(\mathbf{c}_{s(\alpha)-1}, \mathbf{c}_{t(\alpha)}) + \varrho \delta_b(\mathbf{c}_{t(\alpha)}, \mathbf{c}_{t(\alpha)+1}) \right] \\ &\quad + (1 - \mathbf{a}_{s(\alpha)-1}) \varrho \left[\omega \delta_b(\mathbf{c}_{s(\alpha)-1}, \top \mathbf{c}_{s(\alpha)-1}) + (\omega + 1) \delta_b(\top \mathbf{c}_{t(\alpha)+1}, \mathbf{c}_{t(\alpha)+1}) \right], \end{aligned}$$

we obtain $\lim_{\alpha \rightarrow \infty} \delta_b(\mathbf{c}_{s(\alpha)}, \mathbf{c}_{t(\alpha)+1}) \leq \frac{1}{4\varrho^2} \varrho \varepsilon_0 < \frac{1}{\varrho} \varepsilon_0$, a negation. Thus, an O-sequence $\{\mathbf{c}_\vartheta\}$ is a Cauchy in Y . By completeness property, implies that $\exists \mathbf{c}^* \in Y$ such that

$$\lim_{\vartheta \rightarrow \infty} \delta_b(\mathbf{c}_\vartheta, \mathbf{c}^*) = 0.$$

Now prove that \mathbf{c}^* is a fixed point of \top . Since

$$\begin{aligned} \delta_b(\mathbf{c}^*, \top \mathbf{c}^*) &\leq \varrho[\delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + \delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}^*)] \\ &\leq \varrho[\delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + \varrho^2[\delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}_\vartheta) + \delta_b(\top \mathbf{c}_\vartheta, \top \mathbf{c}^*)]] \\ &\leq \varrho \delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + \varrho^2 \delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}_\vartheta) + \varrho^2 \omega [\delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}_\vartheta) + \delta_b(\mathbf{c}^*, \top \mathbf{c}^*)], \end{aligned}$$

we conclude that

$$\begin{aligned} (1 - \varrho^2 \omega) \delta_b(\mathbf{c}^*, \top \mathbf{c}^*) &\leq \varrho \delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + (\varrho^2 + \varrho^2 \omega) \delta_b(\mathbf{c}_\vartheta, \top \mathbf{c}_\vartheta) \\ &\leq \varrho \delta_b(\mathbf{c}^*, \mathbf{c}_\vartheta) + (\varrho^2 + \varrho^2 \omega) \left(\frac{9}{11}\right)^\vartheta \delta_b(\mathbf{c}_0, \top \mathbf{c}_0). \end{aligned}$$

Consequently, we have $\lim_{\vartheta \rightarrow \infty} \delta_b(\mathbf{c}^*, \top \mathbf{c}^*) = 0$, so \mathbf{c}^* is a fixed point of \top . Next, prove the uniqueness part. Let $\mathbf{c}^*, \mathfrak{z}$ be two fixed points of \top and assume that $\top^\vartheta \mathbf{c}^* = \mathbf{c}^* \neq \mathfrak{z} = \top^\vartheta \mathfrak{z} \forall \vartheta \in \mathcal{N}$. By choice of \mathbf{c}_0 , we have

$$(\mathbf{c}_0 \perp \mathbf{c}^* \text{ and } \mathbf{c}_0 \perp \mathfrak{z}) \text{ or } (\mathbf{c}^* \perp \mathbf{c}_0 \text{ and } \mathfrak{z} \perp \mathbf{c}_0).$$

Since \top is \perp -preserving, we have

$$(\top^\vartheta c_0 \perp \top^\vartheta c^* \text{ and } \top^\vartheta c_0 \perp \mathfrak{z}) \text{ or } (\top^\vartheta c^* \perp c_0 \text{ and } \top^\vartheta \mathfrak{z} \perp c_0)$$

for all $\vartheta \in \mathcal{N}$. Now

$$\delta_b(c^*, \mathfrak{z}) = \delta_b(\top^\vartheta c^*, \top^\vartheta \mathfrak{z}) \leq \varrho[\delta_b(\top^\vartheta c^*, \top^\vartheta c_0) + \delta_b(\top^\vartheta c_0, \top^\vartheta \mathfrak{z})].$$

As $\vartheta \rightarrow \infty$, we obtain $\delta_b(c^*, \mathfrak{z}) \leq 0$. Thus, $c^* = \mathfrak{z}$. Hence, \top has a unique fixed point in Y . \square

Now, we demonstrate an illustration of Theorem 2.

Example 6. Let $Y = \mathfrak{R}^+ \cup \{0\}$, $\top : Y \rightarrow Y$ define by

$$\top c = \begin{cases} 0, & \text{if } c \in [0, \frac{\sqrt{5}}{2}), \\ \frac{1}{4c}, & \text{if } c \in [\frac{\sqrt{5}}{2}, +\infty). \end{cases}$$

For any $c, \sigma \in Y$, let $\delta_b : Y \times Y \rightarrow [0, +\infty)$ be a function defined by $\delta_b(c, \sigma) = (c - \sigma)^2$, for all $c, \sigma \in Y$. Define the binary relation \perp on Y by $c \perp \sigma$ if $c\sigma \leq (c \vee \sigma)$, where $c \vee \sigma = c$ or σ and the mapping $\mathcal{V} : Y \times Y \times [0, 1] \rightarrow Y$ as

$$\mathcal{V}(c, \sigma; a) = ac + (1 - a)\sigma.$$

Let c_0 be the initial value and $c_\vartheta = \mathcal{V}(c_{\vartheta-1}, \top c_{\vartheta-1}; a_{\vartheta-1})$, where $a_{\vartheta-1} = \frac{1}{4\varrho}$. If $\varrho = \frac{1}{4\varrho^2}$, then \top has a unique fixed point in Y .

Proof. From Example 5, we know that $(Y, \perp, \delta_b, \mathcal{V})$ is a orthogonal convex b_\perp -MS with $\varrho = 2$. We show that \top satisfies the follows

$$\delta_b(\top c, \top \sigma) \leq \varrho[\delta_b(c, \top c) + \delta_b(\sigma, \top \sigma)] \tag{11}$$

for any $c, \sigma \in Y$. Now, we arise the below cases.

- (i) If $c, \sigma \in [0, \frac{\sqrt{5}}{2})$, then, we shows that Equation (11) holds.
- (ii) If $c \in [0, \frac{\sqrt{5}}{2})$, $\sigma \in [\frac{\sqrt{5}}{2}, +\infty)$, then

$$\begin{aligned} \delta_b(\top c, \top \sigma) - \frac{1}{16}[\delta_b(c, \top c) + \delta_b(\sigma, \top \sigma)] &= (\frac{1}{4\sigma})^2 - \frac{1}{16}[c^2 + (\sigma - \frac{1}{4\sigma})^2] \\ &\leq (\frac{1}{4\sigma})^2 - \frac{1}{16}(\sigma - \frac{1}{4\sigma})^2 \\ &\leq 0, \end{aligned}$$

which implies that

$$\delta_b(\top c, \top \sigma) \leq \frac{1}{16}[\delta_b(c, \top c) + \delta_b(\sigma, \top \sigma)]$$

holds for any $c \in [0, \frac{\sqrt{5}}{2})$ and $\sigma \in [\frac{\sqrt{5}}{2}, +\infty)$.

- (iii) If $c \in [\frac{\sqrt{5}}{2}, +\infty)$ and $\sigma \in [0, \frac{\sqrt{5}}{2})$, then, similarly to case (ii), we can also get that inequality (11) holds.

(iv) If $c, \sigma \in [\frac{\sqrt{5}}{2}, +\infty)$, then

$$\begin{aligned} \delta_b(\top c, \top \sigma) - \frac{1}{16}[\delta_b(c, \top c) + \delta_b(\sigma, \top \sigma)] &= \frac{1}{16}(\frac{1}{c} - \frac{1}{\sigma})^2 - \frac{1}{16}[(c - \frac{1}{4c})^2 + (\sigma - \frac{1}{4\sigma})^2] \\ &= \frac{1}{16} \frac{15}{16}(\frac{1}{c^2} + \frac{1}{\sigma^2}) + 1 - [(c^2 + \sigma^2) + \frac{2}{c\sigma}] \\ &\leq \frac{1}{16}[\frac{15}{16}(\frac{1}{c^2} + \frac{1}{\sigma^2}) + 1 - [2c\sigma + \frac{2}{c\sigma}]] \\ &\leq \frac{1}{16}[\frac{15}{16}(\frac{1}{c^2} + \frac{1}{\sigma^2}) + 1 - 4] < 0, \end{aligned}$$

which shows that

$$\delta_b(\top c, \top \sigma) < \frac{1}{16}[\delta_b(c, \top c) + \delta_b(\sigma, \top \sigma)]$$

holds for all $c, \sigma \in [\frac{\sqrt{5}}{2}, +\infty)$. We conclude that, Equation (11) holds for any $c, \sigma \in Y$. Find the unique fixed point in Y . Now, we arises the following two cases.

Case (a): If $c_0 < \frac{\sqrt{5}}{2}$, then

$$\begin{aligned} \top c_0 &= 0, \\ c_1 &= \frac{1}{8}c_0 + \frac{7}{8}\top c_0 = \frac{1}{8}c_0, \\ c_2 &= \frac{1}{8}c_1 + \frac{7}{8}\top c_1 = (\frac{1}{8})^2c_0, \\ c_3 &= \frac{1}{8}c_2 + \frac{7}{8}\top c_2 = (\frac{1}{8})^3c_0, \\ &\dots \\ c_\vartheta &= \frac{1}{8}c_{\vartheta-1} + \frac{7}{8}\top c_{\vartheta-1} = (\frac{1}{8})^\vartheta c_0. \end{aligned}$$

Obviously, $c_\vartheta \rightarrow 0$ as $\vartheta \rightarrow \infty$.

Case (b): If $c_0 \geq \frac{\sqrt{5}}{2}$, then

$$\begin{aligned} \top c_0 &= \frac{1}{4c_0}, \\ c_1 &= \frac{1}{8}c_0 + \frac{7}{8}\top c_0, \\ \frac{c_1}{c_0} &= \frac{1}{8} + \frac{7}{32} \frac{1}{c_0^2} \leq \frac{3}{10}. \end{aligned}$$

If $0 \leq c_1 < \frac{\sqrt{5}}{2}$, then $\top c_1 = 0$. From Case (a), it follows that $c_\vartheta \rightarrow 0$ as $\vartheta \rightarrow \infty$. If $c_1 \geq \frac{\sqrt{5}}{2}$, then $\frac{c_2}{c_1} = \frac{1}{8} + \frac{7}{32} \cdot \frac{1}{c_1^2} \leq \frac{3}{10}$. The above procedure, we conclude that $c_{\vartheta-1} \geq \frac{\sqrt{5}}{2}$. Then, we get

$$\frac{c_\vartheta}{c_{\vartheta-1}} = \frac{1}{8} + \frac{7}{32} \cdot \frac{1}{c_{\vartheta-1}^2} \leq \frac{3}{10}$$

and

$$\frac{c_\vartheta}{c_0} = \frac{c_1}{c_0} \cdot \frac{c_2}{c_1} \dots \frac{c_\vartheta}{c_{\vartheta-1}} \leq (\frac{3}{10})^\vartheta,$$

which implies that $c_\theta \leq (\frac{3}{10})^\theta c_0$. Hence, $\lim_{\theta \rightarrow \infty} c_\theta = 0$, where 0 is a fixed point of \mathbb{T} . Clearly, the unique fixed point of \mathbb{T} in Y is 0. Assume that z is also a fixed point of \mathbb{T} in $[\sqrt{52}, +\infty)$. Then $\mathbb{T}z = z$; i.e., $z = \mathbb{T}z = \frac{1}{4z} \implies z = \frac{1}{2} < \sqrt{52}$, a rebuttal. \square

4. Application

Consider the critical damped motion of the spring-mass m system under the action of an external force Θ is

$$m\left(\frac{d^2\mu}{dv^2}\right) + \Pi\frac{d\mu}{dv} - \Theta(v, \mu(v)) = 0; \text{ with } \mu(0) = 0; \mu'(0) = 0. \tag{12}$$

where $\Pi > 0$ is the dumping constant and $\Theta : [0, s] \times \mathbb{R}^+ \rightarrow \mathbb{R}$ be a continuous map.

Consider the following integral equation equivalent to (12) is

$$\mu(v) = \int_0^s \Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho))\delta\varrho, \tag{13}$$

with $v, \varrho \in [0, s]$.

The Green's function $\Lambda(v, \varrho)$ is defined as

$$\Lambda(v, \varrho) = \begin{cases} \frac{1 - e^{\zeta(v-\varrho)}}{\zeta}, & \text{for } 0 \leq \varrho \leq v \leq s; \\ 0, & \text{for } v \leq \varrho \leq s; \end{cases}$$

where $\zeta = \frac{\pi}{i}$ is a constant ratio. Define $Y = \mathcal{C}([0, s], \mathbb{R})$ be the set of real continuous functions defined on $[0, s]$. Then, for $s \geq 1$, define b_\perp -MS by

$$\delta_b(\mu, \eta) = \sup_{v \in [0, s]} (|\mu(v)| + |\eta(v)|)^2, \tag{14}$$

for all $\mu, \eta \in Y$ with $\varkappa > 1$ and $v \in [0, s]$.

Then, it is simple to verify that $(Y, \delta_b, \varrho \geq 1)$ forms an \mathfrak{D} -complete b_\perp -MS with $\varrho = 2$. The triple $(Y, \delta_b, \varrho \geq 1)$ is denoted by Y .

Then, we show that the Equation (12) admits a solution iff $\exists \mu^* \in Y$ is a solution of the equation

$$\mu(v) = \int_0^s \Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho))\delta\varrho,$$

with $v, \varrho \in [0, s]$.

Theorem 3. Suppose that the problem (12) and define $\mathbb{T} : \mathcal{C}([0, s], \mathbb{R}) \rightarrow \mathcal{C}([0, s], \mathbb{R})$ by

$$\mathbb{T}\mu(v) = \int_0^s \Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho))\delta\varrho,$$

with $v, \varrho \in [0, s]$. Postulate that:

- (i) $\Theta : [0, s] \times \mathbb{R}^+ \rightarrow \mathbb{R}$ is a \perp -continuous function;
- (ii) For all $\mu, \eta \in Y, \exists \varkappa > 0$ such that $\delta_b(\mathbb{T}\mu, \mathbb{T}\eta) > 0$ and $\delta_b(\mu, \eta) > 0$ yields

$$|\Theta(v, \mu(v))| + |\Theta(v, \eta(v))| \leq \mathbb{k}\sqrt{\delta_b(\mu, \eta)}, \tag{15}$$

for all $v \in [0, s]$ and $v > 1$.

- (iii) For all $v \in [0, s]$ and $\mu, \eta \in \mathcal{C}([0, s], \mathbb{R}), \delta_b(\mu(v), \eta(v)) \geq 0 \implies \delta_b(\mathbb{T}\mu(v), \mathbb{T}\eta(v)) \geq 0$.

Then, the integral Equation (12) has a unique solution.

Proof. Define \perp on Y by $\mu \perp \eta \implies \mu(v)\eta(v) \geq \mu(v)$ or $\mu(v)\eta(v) \geq \eta(v)$ for all $v \in [0, s]$:

Now define $\delta_b : Y \times Y \rightarrow (0, \infty)$ by

$$\delta_b(\mu, \eta) = \sup_{v \in [0, s]} (|\mu(v)| + |\eta(v)|)^2, \tag{16}$$

for all $\mu, \eta \in Y$, with $\varkappa > 1$. Therefore, (Y, δ_b) is a complete b -MS. Define $\Upsilon : Y \rightarrow Y$ by

$$\Upsilon\mu(v) = \int_0^s \Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho))\delta\varrho.$$

Now, prove Υ is \perp -preserving. For $\mu, \eta \in Y$ with $\mu \perp \eta$ and $v \in [0, s]$, we get

$$\Upsilon\mu(v) = \int_0^s \Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho))\delta\varrho \geq 1.$$

It shows that $\Upsilon\mu(v)\Upsilon\eta(v) \geq \Upsilon\mu(v)$ and so $\Upsilon\mu(v) \perp \Upsilon\eta(v)$. Then, Υ is \perp -preserving.

We show that Υ is orthogonal convex structure contraction on $\mathcal{C}([0, s], \mathfrak{R})$. By stipulation (iii) we have $\delta_b(\Upsilon\mu, \Upsilon\eta) > 0$. By the stipulations (i) and (ii) of the theorem, we obtain

$$\begin{aligned} \left(\left| \Upsilon\mu(v) \right| + \left| \Upsilon\eta(v) \right| \right)^2 &= \left(\left| \int_0^s \Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho))\delta\varrho \right| + \left| \int_0^s \Lambda(v, \varrho)\Theta(\varrho, \eta(\varrho))\delta\varrho \right| \right)^2 \\ &\leq \left(\int_0^s \left| \Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho)) \right| \delta\varrho + \int_0^s \left| \Lambda(v, \varrho)\Theta(\varrho, \eta(\varrho)) \right| \delta\varrho \right)^2 \\ &\leq \left(\int_0^s \Lambda(v, \varrho) \left(\left| \Theta(\varrho, \mu(\varrho)) \right| + \left| \Theta(\varrho, \eta(\varrho)) \right| \right) \delta\varrho \right)^2 \\ &\leq \mathbb{k} \left(\int_0^s \Lambda(v, \varrho) \left(\sqrt{\delta_b(\mu, \eta)} \right) \delta\varrho \right)^2 \\ &\leq \mathbb{k}\delta_b(\mu, \eta) \left(\int_0^s \Lambda(v, \varrho)\delta\varrho \right)^2. \end{aligned} \tag{17}$$

Then, we have

$$\left(\left| \Upsilon\mu(v) \right| + \left| \Upsilon\eta(v) \right| \right)^2 \leq \mathbb{k}\delta_b(\mu, \eta) \left(\int_0^s \Lambda(v, \varrho)\delta\varrho \right)^2. \tag{18}$$

Since $\int_0^s \Lambda(v, \varrho)\delta\varrho \leq 1$ and applying supremum on both sides, we get

$$\left(\left| \Upsilon\mu(v) \right| + \left| \Upsilon\eta(v) \right| \right)^2 \leq \mathbb{k}\delta_b(\mu, \eta). \tag{19}$$

Then $\delta_b(\Upsilon\mu, \Upsilon\eta) \leq \mathbb{k}\delta_b(\mu, \eta)$. Thus the condition (2) is satisfied. Therefore, all the conditions of Theorem 1 are satisfied. Hence the operator has a unique fixed point, which means that the integral Equation (12) has a unique solution. This completes the proof. \square

5. Example

Let us consider the following nonlinear integral equation

$$\mu(v) = \int_0^v [(v(1 - \varrho))^{\varsigma-1} - (v - \varrho)^{\varsigma-1}] \cos(\mu(\varrho))\delta\varrho, \tag{20}$$

with $0 \leq \varrho \leq v \leq 1$. Define $\Upsilon : \mathcal{C}([0, s], \mathfrak{R}) \rightarrow \mathcal{C}([0, s], \mathfrak{R})$ by

$$\Upsilon\mu(v) = \int_0^v [(v(1 - \varrho))^{\varsigma-1} - (v - \varrho)^{\varsigma-1}] \cos(\mu(\varrho))\delta\varrho.$$

Given the conditions of Theorem 3, it is simple to demonstrate that Equation (20) has a unique solution for $\zeta = 1$ and $\varrho = 1$. Additionally, we will emphasize the viability of our strategies using the iteration process.

$$\mu_{\vartheta+1}(v) = \int_0^1 [(v(1 - \varrho))^{\zeta-1} - (v - \varrho)^{\zeta-1}] \cos(\mu_{\vartheta}(\varrho)) \delta\varrho.$$

Let $\zeta \in (1, 2)$. Let us take $\zeta = 1.5$ and initial point $\mu_0(v) = 0$. The sequence $\mu_{\vartheta+1}(v) = \int_0^1 [(v(1 - \varrho))^{\zeta-1} - (v - \varrho)^{\zeta-1}] \cos(\mu_{\vartheta}(\varrho)) \delta\varrho$ is shown in Table 1 for $v = 0.1$ converge to the exact solution $\mu(0.1) = \mathbb{T}(\mu(0.1)) = 0.033$.

Table 1. For $v = 0.1$ exact solution is $\mu(0.1) = 0.033$.

ϑ	$\mu_{\vartheta+1}(0.1)$	Approximate Solution	Absolute Error
0	$\mu_1(0.1)$	0.0308	2.5×10^{-3}
1	$\mu_2(0.1)$	0.0307	2.6×10^{-3}
2	$\mu_3(0.1)$	0.0307	2.6×10^{-3}

We obtain the interpolated graphs of nonlinear integral equation for $v = 0.1$, we get the following interpolated graphs, Figure 1 respectively.




Figure 1. Interpolated graph for $t = 0.1$.

Example 7. Assume the following nonlinear integral equation.

$$\mu(v) = \int_0^1 \Lambda(v, \varrho) \Theta(\varrho, \mu(\varrho)) \delta\varrho, \quad \text{for all } \varrho \in [0, 1]$$

Then it has a solution in \mathbb{T} .

Proof. Let $\mathbb{T} : Y \rightarrow Y$ be defined by

$$\mathbb{T}\mu(v) = \int_0^1 \Lambda(v, \varrho) \Theta(\varrho, \mu(\varrho)) \delta\varrho,$$

and set $\Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho)) = \frac{3}{4}\varrho\Theta(\varrho, \mu(\varrho))$ and $\Lambda(v, \varrho)\Theta(\varrho, \eta(\varrho)) = \frac{3}{4}\varrho\Theta(\varrho, \eta(\varrho))$, for all $\mu, \eta \in [0, 1]$. Then we have

$$\begin{aligned} |\Lambda(v, \varrho)\Theta(\varrho, \mu(\varrho)) + \Lambda(v, \varrho)\Theta(\varrho, \eta(\varrho))|^2 &= \left| \frac{3}{4}\varrho\Theta(\varrho, \mu(\varrho)) + \frac{3}{4}\varrho\Theta(\varrho, \eta(\varrho)) \right|^2 \\ &= \frac{9}{16}\varrho^2 |\Theta(\varrho, \mu(\varrho)) + \Theta(\varrho, \eta(\varrho))|^2 \leq \delta_b(\mu, \eta). \end{aligned}$$

Furthermore, see that $\frac{3}{4} \int_0^1 \varrho^2 \delta \varrho = \frac{3}{4} \left(\frac{1^3}{3} - \frac{0^3}{3} \right) = \frac{1}{4} \leq 1$. Then, it is easy to see that all other conditions of the above application are easy to examine and the above problem has a solution in \mathbb{T} . \square

6. Conclusions

In this manuscript, we established a orthogonal convex structural contraction mapping, proved a number of fixed point theorems utilizing orthogonal b -metric space, and demonstrated the existence of a solution to a unique integral equation developed in mechanical engineering.

Author Contributions: Conceptualization, G.N., A.J.G., G.M. and K.J.; methodology, G.N., A.J.G., G.M. and A.S.B.; validation, E.A., K.J. and N.M.; formal analysis, G.N., A.S.B., A.J.G., G.M. and E.A.; investigation, G.N., A.J.G. and N.M.; writing—original draft preparation, G.N., A.S.B., A.J.G., G.M., K.J., E.A. and N.M.; writing—review and editing, E.A. and N.M.; supervision, E.A. and N.M.; project administration, E.A. and N.M.; funding acquisition, A.S.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors are grateful to all the referees for their valuable suggestions to help improve this paper.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Agarwal, R.P.; Meehan, M.; O’regan, D. *Fixed Point Theory and Applications*; Cambridge University Press: Cambridge, UK, 2001.
2. Alghamdi, M.A.; Hussain, N.; Salimi, P. Fixed point and coupled fixed point theorems on b -metric-like spaces. *J. Inequalities Appl.* **2013**, *2013*, 1–25. [[CrossRef](#)]
3. Bota, M.F.; Karapinar, E. A note on “Some results on multi-valued weakly Jungck mappings in b -metric space”. *Open Math.* **2013**, *11*, 1711–1712. [[CrossRef](#)]
4. Alsulami, H.H.; Karapinar, E.; Piri, H. Fixed points of generalized-Suzuki type contraction in complete-metric spaces. *Discret. Dyn. Nat. Soc.* **2015**, *2015*, 1–8. [[CrossRef](#)]
5. Kadelburg, Z.; Radenović, S. Notes on some recent papers concerning F-contractions in b -metric spaces. *Constr. Math. Anal.* **2018**, *1*, 108–112. [[CrossRef](#)]
6. Mitrović, Z.D. A note on the results of Suzuki, Miculescu and Mihail. *J. Fixed Point Theory Appl.* **2019**, *21*, 1–4. [[CrossRef](#)]
7. Popescu, O.; Stan, G. Two fixed point theorems concerning F-contraction in complete metric spaces. *Symmetry* **2020**, *12*, 58. [[CrossRef](#)]
8. Saleem, N.; Iqbal, I.; Iqbal, B.; Radenović, S. Coincidence and fixed points of multivalued F-contractions in generalized metric space with application. *J. Fixed Point Theory Appl.* **2020**, *22*, 81. [[CrossRef](#)]
9. Czerwik, S. Contraction mappings in b -metric spaces. *Acta Math. Inform. Univ. Ostrav.* **1993**, *1*, 5–11.
10. Debnath, P.; Konwar, N.; Radenović, S. *Metric Fixed Point Theory: Applications in Science, Engineering and Behavioural Sciences*; Springer: Singapore, 2021 ; pp. 314–316.
11. Latif, A.; Al Subaie, R.F.; Alansari, M.O. Fixed points of generalized multi-valued contractive mappings in metric type spaces. *J. Nonlinear Var. Anal.* **2022**, *6*, 123–138.
12. Jiang, S.; Wu, Y. An example of a metric space with finite decomposition complexity. *J. Nonlinear Funct. Anal.* **2022**, *2022*, 27.
13. Shatanawi, W.; Rajić, V.Ć.; Radenović, S.; Al-Rawashhdeh, A. Mizoguchi-Takahashi-type theorems in tvs-cone metric spaces. *Fixed Point Theory Appl.* **2012**, *2012*, 1–7. [[CrossRef](#)]
14. Takahashi, W. A convexity in metric space and nonexpansive mappings. I. *Inkodai Math. Semin. Rep.* **1970**, *22*, 142–149. [[CrossRef](#)]

15. Goebel, K.; Kirk, W.A. Iteration processes for nonexpansive mappings. *J. Contemp. Math.* **1983**, *21*, 115–123.
16. Ding, X.P. Iteration processes for nonlinear mappings in convex metric spaces. *J. Math. Anal. Appl.* **1988**, *132*, 114–122. [[CrossRef](#)]
17. Reich, S.; Shafir, I. Nonexpansive iterations in hyperbolic spaces. *Nonlinear Anal.* **1990**, *19*, 537–558. [[CrossRef](#)]
18. Mureşan, V.; Mureşan, A.S. On the theory of fixed point theorems for convex contraction mappings. *Carpathian J. Math.* **2015**, *31*, 365–371. [[CrossRef](#)]
19. Latif, A.; Sintunavarat, W.; Ninsri, A. Approximate fixed point theorems for partial generalized convex contraction mappings in α -complete metric spaces. *Taiwan. J. Math.* **2015**, *19*, 315–333. [[CrossRef](#)]
20. Georgescu, F. Iterated function systems consisting of generalized convex contractions in the framework of complete strong b -metric spaces. *Ann. West Univ. Timis. Math. Comput. Sci.* **2017**, *55*, 119–142. [[CrossRef](#)]
21. Karaca, N.; Yildirim, I. Fixed point theorem for Reich contraction mapping in convex b -metric spaces. *Aligarh Bull. Math.* **2021**, *40*, 29–39.
22. Chen, L.; Yang, N.; Zhao, Y. Fixed point theorems for the Mann’s iteration scheme in convex graphical rectangular b -metric spaces. *Optimization* **2021**, *70*, 1359–1373. [[CrossRef](#)]
23. Dolićanin-Đekić, D.; Bin-Mohsin, B. Some new fixed point results for convex contractions in b -metric spaces. *Univ. Thought Publ. Nat. Sci.* **2019**, *9*, 67–71. [[CrossRef](#)]
24. Asif, A.; Khan, S.U.; Abdeljawad, T.; Arshad, M.; Savas, E. 3D analysis of modified F-contractions in convex b -metric spaces with application to Fredholm integral equations. *AIMS Math.* **2020**, *5*, 6929–6948. [[CrossRef](#)]
25. Berinde, V. Approximating fixed points of almost convex contractions in metric spaces. *Ann. Acad. Rom. Sci. Ser. Math. Appl.* **2020**, *12*, 11–23. [[CrossRef](#)]
26. Bisht, R.K.; Rakočević, V. Fixed points of convex and generalized convex contractions. *Rend. Circ. Mat. Palermo Ser.* **2020**, *69*, 21–28. [[CrossRef](#)]
27. Chen, L.; Li, C.; Kaczmarek, R.; Zhao, Y. Several fixed point theorems in convex b -metric spaces and applications. *Mathematics* **2020**, *8*, 2422. [[CrossRef](#)]
28. Khan, M.S.; Singh, Y.M.; Maniu, G.; Postolache, M. On generalized convex contractions of type-2 in b -metric and 2-metric spaces. *J. Nonlinear Sci. Appl.* **2017**, *10*, 2902–2913. [[CrossRef](#)]
29. Nallaselli, G.; Gnanaprakasam, A.J.; Mani, G.; Mitrović, Z.D.; Aloqaily, A.; Mlaiki, N. Integral equation via fixed point theorems on a new type of convex contraction in b -metric and 2-metric spaces. *Mathematics* **2023**, *11*, 344. [[CrossRef](#)]
30. Gordji, M.E.; Ramezani, M.; De La Sen, M.; Cho, Y.J. On orthogonal sets and Banach fixed point theorem. *Fixed Point Theory* **2017**, *18*, 569–578. [[CrossRef](#)]
31. Gordji, M.E.; Habibi, H. Fixed point theory in generalized orthogonal metric space. *J. Linear Topol. Algebra* **2017**, *6*, 251–260.
32. Sawangsup, K.; Sintunavarat, W.; Cho, Y.J. Fixed point theorems for orthogonal F -contraction mappings on O -complete metric spaces. *J. Fixed Point Theory Appl.* **2020**, *22*, 1–14. [[CrossRef](#)]
33. Arul Joseph, G.; Gunasekaran, N.; Absar, U.H.; Gunaseelan, M.; Imran, A.B.; Kamsing, N. Common fixed-points technique for the existence of a solution to fractional integro-differential equations via orthogonal Branciari metric spaces. *Symmetry* **2022**, *14*, 1859.
34. Beg, I.; Mani, G.; Gnanaprakasam, A.J. Fixed point of orthogonal F -Suzuki contraction mapping on O -complete b -metric spaces with applications. *J. Funct. Spaces* **2021**, *2021*, 1–12. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.