

Article

Randomly Stopped Sums, Minima and Maxima for Heavy-Tailed and Light-Tailed Distributions

Remigijus Leipus ^{1,†}, Jonas Šiaulyš ^{2,*,†}, Svetlana Danilenko ^{3,†} and Jūratė Karasevičienė ^{2,†}

¹ Institute of Applied Mathematics, Vilnius University, Naugarduko 24, LT-03225 Vilnius, Lithuania; remigijus.leipus@mif.vu.lt

² Institute of Mathematics, Vilnius University, Naugarduko 24, LT-03225 Vilnius, Lithuania; jurate.karaseviciene@mif.vu.lt

³ Department of Mathematical Statistics, Vilnius Gediminas Technical University, Saulėtekio 11, LT-10223 Vilnius, Lithuania; svetlana.danilenko@vilniustech.lt

* Correspondence: jonas.siaulyš@mif.vu.lt

† These authors contributed equally to this work.

Abstract: This paper investigates the randomly stopped sums, minima and maxima of heavy- and light-tailed random variables. The conditions on the primary random variables, which are independent but generally not identically distributed, and counting random variable are given in order that the randomly stopped sum, random minimum and maximum is heavy/light tailed. The results generalize some existing ones in the literature. The examples illustrating the results are provided.

Keywords: heavy tail; light tail; randomly stopped sums; randomly stopped minima; randomly stopped maxima

MSC: 60E05; 60G50; 91B05



Citation: Leipus, R.; Šiaulyš, J.; Danilenko, S.; Karasevičienė, J. Randomly Stopped Sums, Minima and Maxima for Heavy-Tailed and Light-Tailed Distributions. *Axioms* **2024**, *13*, 355. <https://doi.org/10.3390/axioms13060355>

Academic Editors: Natasa Kontrec, Jelena Vujakovic and Hranislav Milosevic

Received: 14 April 2024

Revised: 16 May 2024

Accepted: 22 May 2024

Published: 25 May 2024

1. Introduction

This paper is devoted to the randomly stopped sums, minima and maxima of heavy- and light-tailed random variables (r.v.s). Such objects appear when the number of the random variables under consideration is unknown and is described by some random integer. In particular, randomly stopped sums appear in such fields as insurance and financial mathematics, survival analysis, risk theory, computer and communication networks, etc. The area of randomly stopped sums for heavy-tailed r.v.s has been well developed for more than 50 years and covers mainly the case of independent identically distributed (i.i.d.) r.v.s. In this paper, we consider the case where the underlying r.v.s are not necessarily identically distributed, although they are independent.

Specifically, suppose that X_1, X_2, \dots are r.v.s defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Define a sequence of partial sums $\{S_n, n \geq 0\}$ by

$$S_0 := 0, \quad S_n := X_1 + \dots + X_n, \quad n \geq 1. \quad (1)$$

The main subject of the paper lies in the study of *randomly stopped sums*:

$$S_\nu := X_1 + \dots + X_\nu,$$

where n in (1) is replaced by a random variable ν , taking values in $\mathbb{N}_0 := \{0, 1, 2, \dots\}$. Throughout this paper, we assume that ν is not degenerate at zero, i.e., $\mathbb{P}(\nu > 0) > 0$. We will call such ν a *counting random variable*.

Further, we will assume that r.v.s X_1, X_2, \dots are independent and counting r.v. ν is independent of the sequence $\{X_1, X_2, \dots\}$. In general, r.v.s X_1, X_2, \dots can be not identically



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

distributed, each having a distribution function (d.f.) $F_{X_k}(x) = \mathbb{P}(X_k \leq x)$, respectively. Consider the d.f.

$$F_{S_\nu}(x) = \mathbb{P}(S_\nu \leq x) = \sum_{n=0}^{\infty} \mathbb{P}(S_n \leq x) \mathbb{P}(\nu = n).$$

The main task considered in this paper is to give conditions guaranteeing that F_{S_ν} is heavy-/light-tailed, provided that some of the d.f.s F_{X_k} or F_ν are heavy-/light-tailed.

Other objects of the paper are the randomly stopped minima and maxima. By the *randomly stopped minimum of sums*, we call the minimum of partial sums:

$$S_{(\nu)} = \begin{cases} \min\{S_1, \dots, S_\nu\}, & \nu \geq 1, \\ 0, & \nu = 0, \end{cases}$$

and by the *randomly stopped maximum of sums*, we call the maximum of partial sums:

$$S^{(\nu)} = \max\{0, S_1, \dots, S_\nu\}.$$

Also, we provide some results for the *randomly stopped minimum*,

$$X_{(\nu)} = \begin{cases} \min\{X_1, \dots, X_\nu\}, & \nu \geq 1, \\ 0, & \nu = 0, \end{cases}$$

and the *randomly stopped maximum*,

$$X^{(\nu)} = \max\{0, X_1, \dots, X_\nu\}.$$

Similarly, we are interested in when $F_{X_{(\nu)}}$, $F_{X^{(\nu)}}$, $F_{S_{(\nu)}}$ and $F_{S^{(\nu)}}$ are heavy-tailed or light-tailed. The most attention we pay is to the closure of heavy-tailed and light-tailed classes of distributions with respect to random transformations under consideration. For example, Proposition 1 (see parts (iii), (iv)) below implies that a randomly stopped sum remains heavy-tailed if at least one of the primary r.v.s $\{X_1, X_2, \dots\}$ reached by the counting r.v. ν is heavy-tailed. Proposition 2 (see parts (i), (ii)) shows that the randomly stopped maximum has an analogous property. Meanwhile, Proposition 3 (i) shows that the randomly stopped minimum remains heavy-tailed if the first primary r.v. X_1 is heavy-tailed, and the tails of other primary r.v.s are asymptotically compared to the distribution tail of the first primary r.v. Proposition 5 (iii) implies that the randomly stopped maximum of sums for any counting r.v. remains heavy-tailed if the first primary r.v. X_1 is heavy-tailed. Meanwhile, according to Proposition 4 (i), in order for the randomly stopped maximum to remain heavy-tailed, it is necessary that the other primary r.v.s $\{X_2, X_3, \dots\}$ obtain some nonnegative values. Similar facts about the closure of the class of light-tailed distributions with respect to the considered transformations can also be obtained from Propositions 1–5 below. For various distribution classes, similar questions on the closure with respect to various transformations have been studied in [1–30]. In particular, regularly varying distributions were considered in [23], consistently varying distributions in [2,15], long-tailed distributions in [18,19,21] and dominatedly varying distributions in [6,18,19]. Maxima and sums of nonstationary random-length sequences of random variables with regularly varying tails were studied in [31]. We mention also paper [32], where two independent heavy-tailed r.v.s, such that their minimum is not heavy-tailed, were constructed.

One of the incentives to study the randomly stopped structures is related to the models describing the insurance business. According to the well-known Sparre Andersen model [33], the insurer’s wealth $W_u(t)$ is described by the risk renewal model:

$$W_u(t) = u + pt - \sum_{k=1}^{N_\theta(t)} Z_k, \quad t \geq 0,$$

where $u \geq 0$ is the initial capital, $p > 0$ is a constant premium rate, $N_\theta(t)$ is a counting process generated by a sequence of not negative r.v.s $\{\theta_1, \theta_2, \dots\}$ and $\{Z_1, Z_2, \dots\}$ is a sequence of independent random claims. Due to such a model, the behavior of the insurer's wealth $W_u(t)$ is driven by the randomly stopped sums

$$S_\theta(t) = \sum_{k=1}^{N_\theta(t)} Z_k, \quad t \geq 0,$$

and the model ruin probability,

$$\psi(u) = \mathbb{P}\left(\inf_{t>0} W_u(t) < 0\right)$$

is related to maximum of the randomly stopped sums

$$\max_{0 < t \leq T} S_\theta(t).$$

It is well known that the behavior of $S_\theta(t)$, the selection of the premium rate p and the estimation of the ruin probability depends on whether the generating elements $\{\theta_1, \theta_2, \dots\}$, $\{Z_1, Z_2, \dots\}$ and $S_\theta(t)$ have light tails or heavy tails, even in the case that the distributions generating the model are identically distributed. For details, see [34–38].

We also note the well known duality of the homogeneous risk renewal model and the G/G/1 model from queuing theory, where the arrivals follow the counting process generated by distribution F_θ and service times have distribution F_Z . Then the probability of ruin $\psi(u)$ coincides with the probability that the stationary waiting time exceeds u . For details see [34].

The structure of the paper is as follows. In Section 2, we introduce heavy- and light-tailed distributions and formulate two auxiliary lemmas. The main results are formulated in Section 3. Some examples of nonstandard heavy-tailed and light-tailed distributions are presented in Section 4. The heaviness of the distribution tails presented in Section 4 is determined on the basis of the statements formulated in Section 3. The proofs of the main results are presented in Section 5. The last Section 6 is devoted to the discussion of the obtained results in the broadest context together with the highlighting of future research directions.

2. Heavy-Tailed and Light-Tailed Distributions

For any distribution F , define its Laplace–Stieltjes transform as

$$\widehat{F}(\lambda) := \int_{-\infty}^{\infty} e^{\lambda x} dF(x), \quad \lambda \in \mathbb{R}.$$

A distribution F is said to be *heavy-tailed*, denoted $F \in \mathcal{H}$, if

$$\widehat{F}(\lambda) = \infty \quad \text{for any } \lambda > 0.$$

Otherwise, F is said to be *light-tailed*. Common examples of heavy-tailed distributions are Pareto, log-normal, Weibull with shape parameter $\tau \in (0, 1)$, Burr and Student's t distributions. For a detailed exposition of the heavy-tailed distributions and their properties, we refer to monographs [36,39–44].

We formulate two lemmas that will be used in the proofs of several main propositions. Although the results of the lemmas are well known and can be found, e.g., in [41,43,44], we provide the proofs for the sake of convenience. The first lemma gives equivalent conditions for the distribution F to be heavy-/light-tailed.

Lemma 1. *Suppose that F is a d.f. of a real-valued r.v. The following statements are equivalent:*

- (i) F is heavy-tailed,

- (ii) $\limsup_{x \rightarrow \infty} e^{\lambda x} \bar{F}(x) = \infty$ for any $\lambda > 0$,
- (iii) $\limsup_{x \rightarrow \infty} x^{-1} \log \bar{F}(x) = 0$.

Similarly, the equivalent are the following statements:

- (i') F is light-tailed,
- (ii') $\limsup_{x \rightarrow \infty} e^{\lambda x} \bar{F}(x) < \infty$ for some $\lambda > 0$,
- (iii') $\limsup_{x \rightarrow \infty} x^{-1} \log \bar{F}(x) < 0$.

Proof. We prove only the first part of the lemma.

(i) \Rightarrow (iii). Suppose that $\hat{F}(\lambda) = \infty$ for any $\lambda > 0$. Let, on the contrary,

$$\limsup_{x \rightarrow \infty} \frac{\log \bar{F}(x)}{x} < 0.$$

Then, there exist constants $c > 0$ and $x_c > 0$ such that $x^{-1} \log \bar{F}(x) \leq -c$ for $x \geq x_c$, or, equivalently,

$$\bar{F}(x) \leq e^{-cx}, \quad x \geq x_c. \tag{2}$$

For any $\delta \in (0, c)$, using (2) and the alternative expectation formula (see [45], for instance), we obtain

$$\begin{aligned} \int_{[0, \infty)} e^{\delta u} dF(u) &= 1 + \delta \int_0^\infty e^{\delta u} \bar{F}(u) du \\ &= 1 + \left(\int_1^{e^{\delta x_c}} + \int_{e^{\delta x_c}}^\infty \right) \bar{F}(\delta^{-1} \log u) du \\ &\leq e^{\delta x_c} + \int_{e^{\delta x_c}}^\infty e^{-c\delta^{-1} \log u} du \\ &= e^{\delta x_c} + \int_{e^{\delta x_c}}^\infty u^{-c\delta^{-1}} du. \end{aligned}$$

Since $c\delta^{-1} > 1$, the last integral is finite; hence,

$$\hat{F}(\delta) \leq F(0) + \int_{[0, \infty)} e^{\delta u} dF(u) < \infty,$$

leading to a contradiction.

(iii) \Rightarrow (ii). From the condition

$$\limsup_{x \rightarrow \infty} x^{-1} \log \bar{F}(x) = 0$$

we obtain that there exists an infinitely increasing sequence $\{x_n\}$ such that

$$\lim_{n \rightarrow \infty} x_n^{-1} \log \bar{F}(x_n) = 0.$$

For any given $\lambda > 0$, this implies that there exists $n_\lambda \geq 1$ such that

$$x_n^{-1} \log \bar{F}(x_n) \geq -\lambda/2$$

for all $n \geq n_\lambda$. Equivalently,

$$e^{\lambda x_n} \bar{F}(x_n) \geq e^{\lambda x_n/2}, \quad n \geq n_\lambda.$$

Hence, $e^{\lambda x_n} \bar{F}(x_n)$ tends to infinity as $n \rightarrow \infty$, and thus,

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \bar{F}(x) \geq \lim_{n \rightarrow \infty} e^{\lambda x_n} \bar{F}(x_n) = \infty.$$

Since this holds for any $\lambda > 0$, we have (ii).

(ii) \Rightarrow (i). Let

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \bar{F}(x) = \infty$$

for any $\lambda > 0$. For $x \in \mathbb{R}$, write

$$\int_{-\infty}^{\infty} e^{\lambda u} dF(u) \geq \int_{(x, \infty)} e^{\lambda u} dF(u) \geq e^{\lambda x} \bar{F}(x).$$

Thus,

$$\hat{F}(\lambda) \geq \limsup_{x \rightarrow \infty} e^{\lambda x} \bar{F}(x) = \infty \text{ for any } \lambda > 0,$$

and Lemma 1 is proved. \square

The next lemma implies that \mathcal{H} and \mathcal{H}^c are closed with respect to weak tail equivalence.

Lemma 2. Let F and G be two distributions of real-valued r.v.s.

(i) If $F \in \mathcal{H}$ and

$$\liminf_{x \rightarrow \infty} \frac{\bar{G}(x)}{\bar{F}(x)} > 0, \tag{3}$$

then $G \in \mathcal{H}$.

(ii) If $F \in \mathcal{H}^c$, and $\bar{G}(x) \leq \tilde{c} \bar{F}(x)$ for some $\tilde{c} > 0$ and large x ($x > x_{\tilde{c}}$), then $G \in \mathcal{H}^c$.

Proof. Consider part (i). By condition (3), we obtain that

$$\bar{G}(x) \geq \hat{c} \bar{F}(x)$$

for some \hat{c} and sufficiently large x ($x > x_{\hat{c}}$). Therefore,

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \bar{G}(x) \geq \hat{c} \limsup_{x \rightarrow \infty} e^{\lambda x} \bar{F}(x) = \infty$$

for any positive λ implying $G \in \mathcal{H}$ by Lemma 1 (ii).

The proof of part (ii) can be constructed in a similar way by using Lemma 1 (ii'), showing that

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \bar{G}(x) < \infty$$

for some $\lambda > 0$. Lemma 2 is proved. \square

3. Main Results

In this section, we formulate the main results of the paper. We start with the randomly stopped sums. We notice that the d.f. F_{S_ν} can become heavy-tailed because of the heavy tail of some element in $\{F_{X_1}, F_{X_2}, \dots\}$ or because of the heavy tail of the counting random variable ν .

Proposition 1. Let X_1, X_2, \dots be independent real-valued r.v.s and let ν be a counting r.v. independent of the sequence $\{X_1, X_2, \dots\}$. Distribution F_{S_ν} is heavy-tailed if at least one of the following conditions is satisfied:

(i) $\inf_{k \geq 1} \mathbb{E} e^{\lambda X_k} > 1$ for any $\lambda > 0$, and $F_\nu \in \mathcal{H}$;

- (ii) $\inf_{k \geq 1} \mathbb{P}(X_k \geq a) = 1$ for some $a > 0$, and $F_v \in \mathcal{H}$;
- (iii) $F_{X_\varkappa} \in \mathcal{H}$ for some $\varkappa \geq 1$, and $\overline{F_v}(x) > 0$ for all $x \in \mathbb{R}$;
- (iv) $F_{X_\varkappa} \in \mathcal{H}$ for some $1 \leq \varkappa \leq \max\{\text{supp}(v)\}$ and $\text{supp}(v) < \infty$.

Distribution F_{S_v} is light-tailed if at least one of the following conditions is satisfied:

- (v) $F_{X_1} \in \mathcal{H}^c, F_v \in \mathcal{H}^c, \overline{F_{X_1}}(x) > 0$ for all $x \in \mathbb{R}$ and

$$\limsup_{x \rightarrow \infty} \sup_{k \geq 1} \frac{\overline{F_{X_k}}(x)}{\overline{F_{X_1}}(x)} < \infty; \tag{4}$$

- (vi) $\sup_{k \geq 1} \mathbb{E} e^{\lambda X_k} < \infty$ for some $\lambda > 0$, and $F_v \in \mathcal{H}^c$.

Our next statement is about the randomly stopped maximum of r.v.s. We observe that some conditions under which the distribution of the randomly stopped maximum $F_{X^{(v)}}$ becomes heavy-tailed are the same as in Proposition 1. Unfortunately, we did not find how to make a heavy-tailed distribution $F_{X^{(v)}}$ from the light-tailed primary r.v.s $\{X_1, X_2, \dots\}$.

Proposition 2. Let X_1, X_2, \dots be independent real-valued r.v.s and let v be a counting r.v. independent of the sequence $\{X_1, X_2, \dots\}$.

- (i) If $F_{X_\varkappa} \in \mathcal{H}$ for some $\varkappa \geq 1$ and $\overline{F_v}(x) > 0$ for all $x \in \mathbb{R}$, then $F_{X^{(v)}} \in \mathcal{H}$;
- (ii) If $F_{X_\varkappa} \in \mathcal{H}$ for some $\varkappa \leq \max\{\text{supp}(v)\} < \infty$, then $F_{X^{(v)}} \in \mathcal{H}$;
- (iii) Distribution $F_{X^{(v)}}$ belongs to the class \mathcal{H}^c if $F_{X_1} \in \mathcal{H}^c, \overline{F_{X_1}}(x) > 0$ for all $x \in \mathbb{R}, \mathbb{E}v < \infty$ and

$$\limsup_{x \rightarrow \infty} \sup_{n \geq 1} \frac{1}{n} \sum_{k=1}^n \frac{\overline{F_{X_k}}(x)}{\overline{F_{X_1}}(x)} < \infty. \tag{5}$$

The statement below is on the distribution of the randomly stopped minimum of r.v.s. From the formulation below, we observe that the tail of the d.f. $F_{X^{(v)}}$ has much less chance of becoming heavy compared to the d.f.s F_{S_v} and $F_{X^{(v)}}$.

Proposition 3. Let X_1, X_2, \dots be independent real-valued r.v.s and let v be a counting r.v. independent of the sequence $\{X_1, X_2, \dots\}$.

- (i) If $F_{X_1} \in \mathcal{H}$ and

$$\liminf_{x \rightarrow \infty} \min_{1 \leq k \leq \varkappa} \frac{\overline{F_{X_k}}(x)}{\overline{F_{X_1}}(x)} > 0$$

for $\varkappa = \min\{\text{supp}(v) \setminus \{0\}\}$, then $F_{X^{(v)}} \in \mathcal{H}$ and

$$\overline{F_{X^{(v)}}}(x) \underset{x \rightarrow \infty}{\sim} \mathbb{P}(v = \varkappa) \overline{F_{X^{(\varkappa)}}}(x);$$

- (ii) If $F_{X_k} \in \mathcal{H}^c$ for $1 \leq k \leq \varkappa = \min\{\text{supp}(v) \setminus \{0\}\}$, then $F_{X^{(v)}} \in \mathcal{H}^c$.

The next two statements are on the heaviness of randomly stopped minimum of sums and randomly stopped maximum of sums. It can be seen from the presented formulations that some of the conditions were already present in the previous statements. However, for the sake of clarity, we present the full statements on the heaviness of $F_{S^{(v)}}$ and $F_{S^{(v)}}$.

Proposition 4. Let X_1, X_2, \dots be independent real-valued r.v.s and let v be a counting r.v. independent of the sequence $\{X_1, X_2, \dots\}$.

- (i) If $F_{X_1} \in \mathcal{H}$ and $\min_{1 \leq k \leq \varkappa} \mathbb{P}(X_k \geq 0) > 0$ for $\varkappa = \min\{\text{supp}(v) \setminus \{0\}\}$, then $F_{S^{(v)}} \in \mathcal{H}$ and

$$\overline{F_{S^{(v)}}}(x) \underset{x \rightarrow \infty}{\asymp} \overline{F_{X_1}}(x). \tag{6}$$

(ii) If $F_{X_1} \in \mathcal{H}^c$, then $F_{S_{(v)}} \in \mathcal{H}^c$ for any r.v. v .

Proposition 5. Let $\{X_1, X_2, \dots\}$ and v be r.v.s. such as in Propositions 1–4. Then $F_{S_{(v)}} \in \mathcal{H}$ if at least one of the following conditions is satisfied:

- (i) $\inf_{k \geq 1} \mathbb{E} e^{\lambda X_k} > 1$ for all $\lambda > 0$ and $F_v \in \mathcal{H}$;
- (ii) $\inf_{k \geq 1} \mathbb{P}(X_k \geq a) = 1$ for some $a > 0$ and $F_v \in \mathcal{H}$;
- (iii) $F_{X_1} \in \mathcal{H}$;
- (iv) $F_{X_\varkappa} \in \mathcal{H}$ for some $\varkappa \geq 1$ in the case of infinite $\text{supp}(v)$ or for some $1 \leq \varkappa \leq \max\{\text{supp}(v)\}$ in the case of finite $\text{supp}(v)$.

Distribution $F_{S_{(v)}}$ is light-tailed if:

- (v) $\sup_{k \geq 1} \mathbb{E} e^{\lambda X_k} < \infty$ for some $\lambda > 0$ and $F_v \in \mathcal{H}^c$.

In the i.i.d. case, Proposition 1 immediately implies the following corollaries. Note that the first two corollaries can be found in monograph [41] as Problems 2.12 and 2.13.

Corollary 1. Let X_1, X_2, \dots be i.i.d. real-valued r.v.s with common distribution F_{X_1} , and let v be a counting r.v. independent of $\{X_1, X_2, \dots\}$. If $F_{X_1} \in \mathcal{H}^c$ and $F_v \in \mathcal{H}^c$, then $F_{S_v} \in \mathcal{H}^c$.

Corollary 2. Let X_1, X_2, \dots be i.i.d. nonnegative not degenerate at zero r.v.s, and let v be a counting r.v. independent of $\{X_1, X_2, \dots\}$. If $F_v \in \mathcal{H}$, then $F_{S_v} \in \mathcal{H}$.

Corollary 3. Let X_1, X_2, \dots be i.i.d. real-valued r.v.s with common distribution F_{X_1} , and let v be a counting r.v. independent of $\{X_1, X_2, \dots\}$. If $F_{X_1} \in \mathcal{H}$ then $F_{S_v} \in \mathcal{H}$.

Analogous corollaries can be formulated for randomly stopped minima and maxima.

4. Examples

In this section, we present two examples showing how one concretely can construct heavy-tailed distributions by using the above randomly stopped structures.

Example 1. Let $\{X_1, X_2, \dots\}$ be a sequence of independent r.v.s such that the first member X_1 has the Pareto distribution

$$F_{X_1}(x) = \left(1 - \frac{1}{(1+x)^3}\right) \mathbb{1}_{[0,\infty)}(x),$$

and other elements of the sequence are identically exponentially distributed:

$$F_{X_k}(x) = (1 - e^{-x}) \mathbb{1}_{[0,\infty)}(x), \quad k \in \{2, 3, \dots\}$$

According to Proposition 1 (parts (iii) and (iv)) and Proposition 5 (iii), distributions F_{S_v} and $F_{S_{(v)}}$ are heavy-tailed for any counting r.v. independent of the sequence $\{X_1, X_2, \dots\}$. This is due to the fact that the first of all primary distributions has a significantly heavier tail than the other elements of the infinite primary sequence. For instance, in the case of the discrete uniform counting r.v. with parameter $N \geq 2$, we have that distributions with the tail

$$\begin{aligned} \overline{F_{S_v}}(x) &= \overline{F_{S_{(v)}}}(x) \\ &= \mathbb{1}_{(-\infty,0)}(x) + \left(\frac{1}{(1+x)^3} + \frac{1}{N} \sum_{k=1}^{N-1} \frac{1}{(k-1)!} \int_0^x \frac{y^{k-1} e^{-y}}{(1+(x-y))^3} dy\right) \mathbb{1}_{[0,\infty)}(x) \end{aligned}$$

belong to the class \mathcal{H} . Proposition 2 (ii) implies that distribution $F_{X(v)}$ belongs to the class \mathcal{H} for any counting r.v. independent of $\{X_1, X_2, \dots\}$. Meanwhile Proposition 3 (i) and

Proposition 4 (i) imply that $F_{X_{(\nu)}}$ and $F_{S_{(\nu)}}$ are heavy-tailed for counting r.v. under condition $1 \in \text{supp}(\nu)$. In the case of the discrete uniform counting r.v. ν with parameter $N = 3$, we have that $F_{S_{(\nu)}} = F_{X_1}$ and distributions with the following tails are heavy-tailed:

$$\begin{aligned} \overline{F_{X_{(\nu)}}}(x) &= \mathbb{1}_{(-\infty,0)}(x) + \left(\frac{1}{(1+x)^3} + \left(e^{-x} - \frac{e^{-2x}}{3} \right) \left(1 - \frac{1}{(1+x)^3} \right) \right) \mathbb{1}_{[0,\infty)}(x), \\ \overline{F_{S_{(\nu)}}}(x) &= \mathbb{1}_{(-\infty,0)}(x) + \frac{1}{3(1+x)^3} (1 + e^{-x} + e^{-2x}) \mathbb{1}_{[0,\infty)}(x). \end{aligned}$$

Example 2. Let $\{X_1, X_2, \dots\}$ be a sequence of independent r.v.s uniformly distributed on the interval $[0, 1]$, i.e.,

$$F_{X_k}(x) = x \mathbb{1}_{[0,1)}(x) + \mathbb{1}_{[1,\infty)}(x)$$

for each $k \in \mathbb{N}$.

Obviously,

$$\mathbb{E} e^{\lambda X_k} = \frac{e^\lambda - 1}{\lambda} > 1$$

for any $\lambda > 0$ and all $k \in \mathbb{N}$. Therefore, by Proposition 1 (i) and Proposition 5 (i), we obtain that distributions F_{S_ν} and $F_{S^{(\nu)}}$ are heavy-tailed for an arbitrary heavy-tailed counting r.v. ν independent of $\{X_1, X_2, \dots\}$. Suppose that counting r.v. ν is distributed according to the zeta distribution with parameter 2:

$$\mathbb{P}(\nu = n) = \frac{1}{n^2} \frac{1}{\zeta(2)}, \quad n \in \mathbb{N},$$

where

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad s \in \mathbb{C},$$

denotes the Riemann zeta function. Such ν is heavy-tailed. Propositions 1 (i) and 5 (i) imply that distribution

$$F_{S_\nu}(x) = F_{S^{(\nu)}}(x) = \frac{1}{\zeta(2)} \sum_{n=1}^{\infty} \frac{1}{n^2} F_{X_1}^{*n}(x) \mathbb{1}_{[0,n]}(x)$$

belongs to class \mathcal{H} , where

$$F_{X_1}^{*n}(x) = \frac{1}{n!} \sum_{k=0}^{\lfloor x \rfloor} (-1)^k \binom{n}{k} (x-k)^n$$

is the well-known Irwin–Hall distribution with parameter n ; see [46,47] or Section 26.9 in [48]. Meanwhile, Propositions 3 (ii) and 4 (ii) imply that distributions with tails

$$\begin{aligned} \overline{F_{S^{(\nu)}}}(x) &= \overline{F_{X_1}}(x), \\ \overline{F_{X_{(\nu)}}}(x) &= \mathbb{1}_{(-\infty,0)}(x) + \frac{1}{\zeta(2)} \sum_{n=1}^{\infty} \frac{1}{n^2} (1-x)^n \mathbb{1}_{[0,1)}(x) \end{aligned}$$

are light-tailed despite the fact that the counting r.v. ν distributed according to the zeta distribution is heavy-tailed.

Example 3. Let $\{X_1, X_2, \dots\}$ be a sequence of independent r.v.s distributed according to the Burr type XII law, i.e.,

$$F_{X_k}(x) = \left(1 - \left(\frac{1}{1 + \sqrt{kx}} \right)^{3/2} \right) \mathbb{1}_{[0,\infty)}(x), \quad k = 1, 2, \dots,$$

and let the counting r.v. v be independent of $\{X_1, X_2, \dots\}$ and distributed according to the shifted Poisson law, i.e.,

$$\mathbb{P}(v = k) = \frac{1}{e(k-3)!}, k = 3, 4, \dots \tag{7}$$

Since $F_{X_1} \in \mathcal{H}$ and

$$\liminf_{x \rightarrow \infty} \min_{1 \leq k \leq 3} \frac{\overline{F_{X_k}}(x)}{\overline{F_{X_1}}(x)} = \min_{1 \leq k \leq 3} \left(\frac{1}{\sqrt{k}}\right)^{3/2} = 3^{-3/4} > 0,$$

we obtain from Proposition 3 (i) that $F_{X(v)} \in \mathcal{H}$ and

$$\overline{F_{X(v)}}(x) \underset{x \rightarrow \infty}{\sim} \frac{1}{e} \overline{F_{X(3)}}(x) \tag{8}$$

with

$$\begin{aligned} \overline{F_{X(3)}}(x) &= \left(\frac{1}{(1 + \sqrt{x})(1 + \sqrt{2x})(1 + \sqrt{3x})} \right)^{3/2}, \\ \overline{F_{X(v)}}(x) &= \frac{1}{e} \sum_{n=3}^{\infty} \frac{1}{(n-3)!} \prod_{k=1}^n \left(\frac{1}{1 + \sqrt{kx}} \right)^{3/2}. \end{aligned}$$

A graphical representation of the asymptotic (8) is shown in Figure 1.




Figure 1. Comparison of tails $\overline{F_{X(v)}}$ (blue line) and $\overline{F_{X(3)}}$ (red line) from Example 3.

We note that Proposition 3 (i) can also be applied to other Burr type XII distributions whose distribution functions have the form

$$F(x) = \left(1 - \left(1 + \left(\frac{x}{\beta} \right)^\alpha \right)^{-\gamma} \right) \mathbb{I}_{[0, \infty)}(x),$$

where α, β, γ are positive parameters; see [49], for instance.

Example 4. Let $\{X_1, X_2, \dots\}$ be a sequence of independent r.v.s such that F_{X_1} is distributed according to the Weibull law with the scale parameter 1 and the shape parameter 1/2, i.e.,

$$\overline{F_{X_1}}(x) = \mathbb{I}_{(-\infty, 0)}(x) + e^{-\sqrt{x}} \mathbb{I}_{[0, \infty)}(x).$$

Since $F_{X_1} \in \mathcal{H}$, due to Proposition 4 (i), we obtain that the d.f. of the randomly stopped minimum of sums $F_{S(v)}$ is heavy-tailed and

$$\overline{F_{S(v)}}(x) \underset{x \rightarrow \infty}{\asymp} \overline{F_{X_1}}(x)$$

if $\min_{2 \leq k \leq \varkappa} \mathbb{P}(X_k \geq 0) > 0$ for $\varkappa = \min\{\text{supp}(v) \setminus \{0\}\}$.

For example, if

$$\mathbb{P}(X_k = -1) = \mathbb{P}(X_k = 1) = \frac{1}{2}, \quad k \in \{2, 3, \dots\},$$

and ν is distributed according to the shifted Poisson law (7), then $F_{S(v)} \in \mathcal{H}$ and

$$\overline{F_{S(v)}}(x) \underset{x \rightarrow \infty}{\asymp} e^{-\sqrt{x}}.$$

A graphical representation of the last relation is shown in Figure 2, having in mind that

$$\frac{1}{4e} e^{-\sqrt{x}} \leq \overline{F_{S(v)}}(x) \leq e^{-\sqrt{x}}, \quad x \geq 0,$$

and

$$\overline{F_{S(v)}}(x) = \frac{1}{e} \sum_{n=3}^{\infty} \mathbb{P}\left(\bigcap_{k=1}^n \{S_k > x\}\right) \frac{1}{(n-3)!} = \frac{1}{e} \sum_{n=3}^{\infty} \frac{\Delta_n(x)}{(n-3)!},$$

where

$$\begin{aligned} \Delta_{2m}(x) &= \frac{1}{2^{2m-1}} \sum_{k=0}^{m-1} \binom{2m-1}{k} \left(e^{-\sqrt{x+2(m-k)-1}} + e^{-\sqrt{x+2(m-k)-2}} \right), \quad m \in \{2, 3, \dots\}; \\ \Delta_{2m+1}(x) &= \frac{1}{2^{2m}} \sum_{k=0}^{m-1} \binom{2m}{k} \left(e^{-\sqrt{x+2(m-k)}} + e^{-\sqrt{x+2(m-k)-1}} \right) \\ &\quad + \frac{1}{2^{2m}} \binom{2m}{m} e^{-\sqrt{x}}, \quad m \in \{1, 2, \dots\}. \end{aligned}$$




Figure 2. Tail of d.f. $\overline{F_{S(v)}}$ (blue line) and its bounds (red lines) from Example 4.

5. Proofs of the Main Results

In this section, we present the proofs of all main propositions. We assign a separate subsection to the proof of each proposition.

5.1. Proof of Proposition 1

Proof of part (i). For any $\lambda > 0$ and an arbitrary $K \geq 1$, we have

$$\begin{aligned} \mathbb{E} e^{\lambda S_\nu} &= \mathbb{E} \left(e^{\lambda S_\nu} \sum_{n=0}^{\infty} \mathbb{1}_{\{v=n\}} \right) = \mathbb{E} \left(\sum_{n=0}^{\infty} e^{\lambda S_n} \mathbb{1}_{\{v=n\}} \right) \\ &\geq \mathbb{E} \left(\sum_{n=0}^K e^{\lambda S_n} \mathbb{1}_{\{v=n\}} \right) \\ &= \sum_{n=0}^K \mathbb{E} e^{\lambda S_n} \mathbb{P}(v = n). \end{aligned} \tag{9}$$

From the condition

$$\inf_{k \geq 1} \mathbb{E} e^{\lambda X_k} > 1$$

we derive that the estimate

$$\min_{1 \leq k \leq K} \mathbb{E} e^{\lambda X_k} \geq \Delta$$

holds for some $\Delta = \Delta(\lambda) > 1$. Therefore, for all $n \in \{1, \dots, K\}$, we obtain

$$\mathbb{E} e^{\lambda S_n} = \prod_{k=1}^n \mathbb{E} e^{\lambda X_k} \geq \Delta^n. \tag{10}$$

This, together with (9), implies that

$$\mathbb{E} e^{\lambda S_\nu} \geq \sum_{n=0}^K \Delta^n \mathbb{P}(v = n).$$

Since $F_\nu \in \mathcal{H}$, we have

$$\sum_{n=0}^K \Delta^n \mathbb{P}(v = n) = \mathbb{E} e^{v \log \Delta} \mathbb{1}_{\{v \leq K\}} \xrightarrow{K \rightarrow \infty} \infty.$$

Hence, $\mathbb{E} e^{\lambda S_\nu} = \infty$ implying $F_{S_\nu} \in \mathcal{H}$ by definition. Part (i) of the proposition is proved. \square

Proof of part (ii). Let us fix an arbitrary $\lambda > 0$. Due to the conditions of part (ii), for such λ , we have

$$\begin{aligned} \inf_{k \geq 1} \mathbb{E} e^{\lambda X_k} &= \inf_{k \geq 1} (\mathbb{E} e^{\lambda X_k} \mathbb{1}_{\{X_k \geq a\}} + \mathbb{E} e^{\lambda X_k} \mathbb{1}_{\{X_k < a\}}) \\ &\geq \inf_{k \geq 1} \mathbb{E} e^{\lambda X_k} \mathbb{1}_{\{X_k \geq a\}} \\ &\geq \inf_{k \geq 1} e^{\lambda a} \mathbb{P}(X_k \geq a) \\ &= e^{\lambda a} > 1. \end{aligned}$$

Hence, the assertion of part (ii) follows from part (i) of the proposition. \square

Proof of part (iii). The requirement $\overline{F}_v(x) > 0$ for all $x \in \mathbb{R}$ implies that counting r.v. v has an unbounded support. Thus, we can find $K \geq \varkappa$ such that $\mathbb{P}(v = K) > 0$. Let λ be any positive number and $M \geq 1$. Then,

$$\begin{aligned} \mathbb{E} e^{\lambda S_K} &\geq \mathbb{E} \exp \left\{ \lambda \sum_{k=1}^K X_k \mathbb{I}_{\{X_k \leq M\}} \right\} \\ &= \mathbb{E} e^{\lambda X_{\varkappa} \mathbb{I}_{\{X_{\varkappa} \leq M\}}} \prod_{\substack{k=1 \\ k \neq \varkappa}}^K \mathbb{E} e^{\lambda X_k \mathbb{I}_{\{X_k \leq M\}}} \xrightarrow{M \rightarrow \infty} \infty \end{aligned}$$

because $F_{\varkappa} \in \mathcal{H}$ and $\mathbb{E} e^{\lambda X_k} > 0$ for each $k \in \{1, \dots, K\}$. Therefore, $F_{S_K} \in \mathcal{H}$. By representation (9), we obtain that

$$\mathbb{E} e^{\lambda S_v} \geq \mathbb{P}(v = K) \mathbb{E} e^{\lambda S_K}$$

implying $F_{S_v} \in \mathcal{H}$. This completes the proof of part (iii) of the proposition. \square

Proof of part (iv). Let K be such that $\mathbb{P}(v = K) > 0$ and $\varkappa \leq K$. Clearly, the conditions of part (iv) imply the existence of such K . To finish the proof of this part, it is sufficient to repeat the arguments of part (iii). \square

Proof of part (v). Suppose that $0 < \delta \leq \lambda$, and $\lambda > 0$ is such that $\mathbb{E} e^{\lambda X_1^+} < \infty$ with $X_1^+ := X_1 \mathbb{I}_{\{X_1 \geq 0\}}$. By the standard representation (9), we have

$$\begin{aligned} \mathbb{E} e^{\delta S_v} &= \sum_{n=0}^{\infty} \mathbb{E} e^{\delta S_n} \mathbb{P}(v = n) \\ &\leq \sum_{n=0}^{\infty} \mathbb{E} e^{\delta S_n^+} \mathbb{P}(v = n), \end{aligned} \tag{11}$$

where $S_0^+ = 0$ and

$$S_n^+ = \sum_{k=1}^n X_k^+ = \sum_{k=1}^n X_k \mathbb{I}_{\{X_k \geq 0\}}, \quad n \in \{1, 2, \dots\}.$$

Condition (4) implies

$$\overline{F}_{X_k}(x) \leq c_1 \overline{F}_{X_1}(x) \tag{12}$$

for some $c_1 > 0$, all $k \geq 1$ and all $x \in \mathbb{R}$. Therefore, by the alternative expectation formula (see, for instance, [45]), we derive from (12) that

$$\begin{aligned} \mathbb{E} e^{\delta X_k^+} &= 1 + \delta \int_0^{\infty} e^{\delta u} \overline{F}_{X_k^+}(u) \, du \\ &\leq 1 + \delta c_1 \int_0^{\infty} e^{\lambda u} \overline{F}_{X_1}(u) \, du \\ &= 1 + \frac{\delta}{\lambda} c_1 (\mathbb{E} e^{\lambda X_1^+} - 1) := c_2(\delta) \end{aligned}$$

for any $k \geq 1$, where $1 < c_2(\delta) < \infty$ for $0 < \delta \leq \lambda$, and

$$\lim_{\delta \downarrow 0} c_2(\delta) = 1.$$

Since X_1^+, X_2^+, \dots are independent r.v.s, we obtain

$$\mathbb{E} e^{\delta S_n^+} = \prod_{k=1}^n \mathbb{E} e^{\delta X_k^+} \leq (c_2(\delta))^n.$$

Hence, by inequality (11) and condition $F_\nu \in \mathcal{H}^c$ we derive that

$$\mathbb{E} e^{\delta S_\nu} \leq \sum_{n=0}^{\infty} (c_2(\delta))^n \mathbb{P}(\nu = n) = \mathbb{E} e^{\nu \log c_2(\delta)} < \infty$$

if $\delta \in (0, \lambda]$ is chosen as sufficiently small.

This implies that $F_{S_\nu} \in \mathcal{H}^c$. \square

Proof of part (vi). The statement of this part can be proved analogously to the statement of part (v). Namely, the conditions of part (vi) imply that

$$\sup_{k \geq 1} \mathbb{E} e^{\lambda X_k^+} = c_\lambda$$

for some constants $\lambda > 0$ and $c_\lambda \geq 1$. Therefore, using the alternative expectation formula, we derive

$$\begin{aligned} \mathbb{E} e^{\delta X_k^+} &= 1 + \delta \int_{[0, \infty)} e^{\delta u} \overline{F_{X_k}}(u) \, du \\ &\leq 1 + \frac{\delta}{\lambda} \left(\lambda \int_{[0, \infty)} e^{\lambda u} \overline{F_{X_k}}(u) \, du \right) \\ &= 1 + \frac{\delta}{\lambda} (c_\lambda - 1) \end{aligned}$$

for all $\delta \in (0, \lambda)$ and $k \geq 1$. The last estimation and inequality (11) imply that

$$\mathbb{E} e^{\delta S_\nu} \leq \sum_{n=0}^{\infty} \prod_{k=1}^n \mathbb{E} e^{\delta X_k^+} \mathbb{P}(\nu = n) \leq \mathbb{E} e^{\nu \log \left(1 + \frac{\delta}{\lambda} (c_\lambda - 1) \right)}.$$

If $\delta \in (0, \lambda]$ is sufficiently small, then the last expectation is finite because of $F_\nu \in \mathcal{H}^c$. Hence, $F_{S_\nu} \in \mathcal{H}^c$ as well. Part (vi) of the proposition is proved. \square

5.2. Proof of Proposition 2

Proof of part (i). By the standard representation, we have

$$\begin{aligned} \overline{F_{X^{(\nu)}}}(x) &= \sum_{n=1}^{\infty} \mathbb{P}(X^{(n)} > x) \mathbb{P}(\nu = n) \\ &\geq \mathbb{P}(X^{(K)} > x) \mathbb{P}(\nu = K) \end{aligned} \tag{13}$$

for $x > 0$ and any K such that $\mathbb{P}(\nu = K) > 0$, $K \geq \varkappa$. Due to the conditions of part (ii), there exists a sequence of numbers K with the above property. Obviously,

$$\begin{aligned} \mathbb{P}(X^{(K)} > x) &= \mathbb{P}(\max\{0, X_1, \dots, X_K\} > x) \\ &\geq \mathbb{P}(X_\varkappa > x). \end{aligned} \tag{14}$$

Consequently, for an arbitrary $\lambda > 0$, we obtain from (13) and (14)

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X^{(\nu)}}}(x) \geq \mathbb{P}(\nu = K) \limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_\varkappa}}(x).$$

The assertion of part (i) follows now by Lemma 1. \square

Proof of part (ii). The proof of this part is similar to the proof of part (i), because the conditions of part (ii) imply that there exists at least one K such that $K \geq \varkappa$ and $\mathbb{P}(\nu = K) > 0$. \square

Proof of part (iii). The standard representation implies that

$$\overline{F_{X^{(v)}}}(x) = \sum_{n=1}^{\infty} \mathbb{P}(X^{(n)} > x) \mathbb{P}(v = n) \tag{15}$$

$$= \sum_{n=1}^{\infty} \mathbb{P}\left(\bigcup_{k=1}^n \{X_k > x\}\right) \mathbb{P}(v = n) \\ \leq \sum_{n=1}^{\infty} \mathbb{P}(v = n) \sum_{k=1}^n \overline{F_{X_k}}(x) \tag{16}$$

for positive x .

Due to Lemma 1, there is $\lambda > 0$ such that

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_1}}(x) < \infty. \tag{17}$$

It follows from the estimate (15) that

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X^{(v)}}}(x) \leq \limsup_{x \rightarrow \infty} e^{\lambda x} \sum_{n=1}^{\infty} \mathbb{P}(v = n) \sum_{k=1}^n \overline{F_{X_k}}(x).$$

Condition (5) of part (iii) implies that

$$\sum_{k=1}^n \overline{F_{X_k}}(x) \leq c_4 n \overline{F_{X_1}}(x) \tag{18}$$

for all $n \geq 1$, for some $c_4 > 0$ and for sufficiently large x ($x \geq x_1$). Therefore, by (17) and (18), we obtain that

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X^{(v)}}}(x) \leq c_4 \mathbb{E}v \limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_1}}(x) < \infty.$$

The assertion of part (iii) follows now by Lemma 1. \square

5.3. Proof of Proposition 3

Proof of part (i). By the standard representation we have

$$\overline{F_{X^{(v)}}}(x) = \sum_{n=1}^{\infty} \mathbb{P}(\min\{X_1, \dots, X_n\} > x) \mathbb{P}(v = n) \\ = \sum_{n=1}^{\infty} \mathbb{P}(v = n) \prod_{k=1}^n \overline{F_{X_k}}(x) \\ = \overline{F_{X^{(\varkappa)}}}(x) \mathbb{P}(v = \varkappa) + \sum_{n=\varkappa+1}^{\infty} \mathbb{P}(v = n) \overline{F_{X^{(\varkappa)}}}(x) \prod_{k=\varkappa+1}^n \overline{F_{X_k}}(x) \\ \leq \overline{F_{X^{(\varkappa)}}}(x) \mathbb{P}(v = \varkappa) \left(1 + \overline{F_{X_{\varkappa+1}}}(x) \frac{\mathbb{P}(v > \varkappa)}{\mathbb{P}(v = \varkappa)}\right), \tag{19}$$

and

$$\overline{F_{X^{(v)}}}(x) \geq \overline{F_{X^{(\varkappa)}}}(x) \mathbb{P}(v = \varkappa)$$

for each positive x . In addition, conditions of part (i) give that $\overline{F_{X^{(\varkappa)}}}(x) > 0$ for all positive x . Therefore,

$$\overline{F_{X^{(v)}}}(x) \underset{x \rightarrow \infty}{\sim} \mathbb{P}(v = \varkappa) \overline{F_{X^{(\varkappa)}}}(x).$$

We obtain from this, by using Lemma 2, that $F_{X_{(\nu)}} \in \mathcal{H}$ if $F_{X_{(\varkappa)}} \in \mathcal{H}$. Hence, to prove the assertion of part (i) it is enough to prove that $F_{X_{(\varkappa)}} \in \mathcal{H}$ for $1 \leq \varkappa \leq \min\{\text{supp}(\nu) \setminus \{0\}\}$.

Due to the condition $F_{X_1} \in \mathcal{H}$ and Lemma 1, we have

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_1}}(x) = \infty \tag{20}$$

for an arbitrary $\lambda > 0$. The requirement

$$\liminf_{x \rightarrow \infty} \min_{1 \leq k \leq \varkappa} \frac{\overline{F_{X_k}}(x)}{\overline{F_{X_1}}(x)} > 0$$

implies that

$$\overline{F_{X_k}}(x) \geq c_5 \overline{F_{X_1}}(x)$$

for some positive c_5 , sufficiently large x ($x \geq x_2$) and for all $1 \leq k \leq \varkappa$. Therefore, for any positive λ and large x ($x \geq x_2$) we obtain

$$\begin{aligned} e^{\lambda x} \overline{F_{X_{(\varkappa)}}}(x) &= e^{\lambda x} \prod_{k=1}^{\varkappa} \overline{F_{X_k}}(x) \\ &\geq c_5^{\varkappa} e^{\lambda x} (\overline{F_{X_1}}(x))^{\varkappa} \\ &= (c_5 e^{\lambda x / \varkappa} \overline{F_{X_1}}(x))^{\varkappa}. \end{aligned}$$

By relation (20) we derive that

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_{(\varkappa)}}}(x) = \infty$$

implying that $F_{X_{(\varkappa)}} \in \mathcal{H}$. Part (i) of the proposition is proved. \square

Proof of part (ii). According to inequality (19) and Lemma 2, $F_{X_{(\nu)}} \in \mathcal{H}^c$ if $F_{X_{(\varkappa)}} \in \mathcal{H}^c$. Since \varkappa is finite, conditions $F_{X_k} \in \mathcal{H}^c$, $k \in \{1, 2, \dots, \varkappa\}$ and Lemma 1 imply that

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_k}}(x) < \infty \tag{21}$$

for some $\lambda > 0$ and each $k \in \{1, 2, \dots, \varkappa\}$. For this λ and an arbitrary positive x , we have

$$e^{\lambda x} \overline{F_{X_{(\varkappa)}}}(x) = \prod_{k=1}^{\varkappa} \left(e^{\lambda x / \varkappa} \overline{F_{X_k}}(x) \right).$$

Since $\lambda / \varkappa \leq \lambda$, due to (21),

$$\limsup_{x \rightarrow \infty} e^{\lambda x / \varkappa} \overline{F_{X_k}}(x) < \infty$$

for each $k \in \{1, 2, \dots, \varkappa\}$. Therefore,

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_{(\varkappa)}}}(x) < \infty$$

implying that $F_{X_{(\varkappa)}} \in \mathcal{H}^c$ by Lemma 1. Hence, $F_{X_{(\nu)}} \in \mathcal{H}^c$ as well, and part (ii) of the proposition is proved. \square

5.4. Proof of Proposition 4

Proof of part (i). If $\varkappa = 1$, then for $x > 0$, we have

$$\begin{aligned} \overline{F_{S(v)}}(x) &= \sum_{n \in \text{supp}(v) \setminus \{0\}} \overline{F_{S(n)}}(x) \mathbb{P}(v = n) \\ &\geq \overline{F_{S(1)}}(x) \mathbb{P}(v = 1) \\ &= \overline{F_{X_1}}(x) \mathbb{P}(v = 1), \end{aligned}$$

and

$$\begin{aligned} \overline{F_{S(v)}}(x) &= \sum_{n=1}^{\infty} \overline{F_{S(n)}}(x) \mathbb{P}(v = n) \\ &= \sum_{n=1}^{\infty} \mathbb{P}(\min\{S_1, \dots, S_n\} > x) \mathbb{P}(v = n) \\ &= \sum_{n=1}^{\infty} \mathbb{P}\left(\bigcap_{k=1}^n \{S_k > x\}\right) \mathbb{P}(v = n) \\ &\leq \sum_{n=1}^{\infty} \mathbb{P}(S_1 > x) \mathbb{P}(v = n) \\ &= \overline{F_{X_1}}(x) \mathbb{P}(v \geq 1). \end{aligned}$$

The derived estimates imply the asymptotic relation (6) in the case $\varkappa = 1$.

Let us now suppose that $\varkappa > 1$. Due to the conditions of part (i)

$$\mathbb{P}(X_k \geq 0) \geq c_6$$

for some $c_6 > 0$ and all $1 \leq k \leq \varkappa$. Hence, by the standard decomposition, we obtain that for positive x

$$\begin{aligned} \overline{F_{S(v)}}(x) &= \sum_{n=1}^{\infty} \overline{F_{S(n)}}(x) \mathbb{P}(v = n) \\ &\geq \overline{F_{S(\varkappa)}}(x) \mathbb{P}(v = \varkappa) \\ &= \mathbb{P}(\min\{S_1, \dots, S_{\varkappa}\} > x) \mathbb{P}(v = \varkappa) \\ &= \mathbb{P}\left(\bigcap_{k=1}^{\varkappa} \{X_1 + \dots + X_k > x\}\right) \mathbb{P}(v = \varkappa) \\ &\geq \mathbb{P}(X_1 > x, X_2 \geq 0, \dots, X_{\varkappa} \geq 0) \mathbb{P}(v = \varkappa) \\ &= \mathbb{P}(X_1 > x) \prod_{k=2}^{\varkappa} \mathbb{P}(X_k \geq 0) \mathbb{P}(v = \varkappa) \\ &\geq c_6^{\varkappa-1} \mathbb{P}(v = \varkappa) \overline{F_{X_1}}(x). \end{aligned} \tag{22}$$

On the other hand, similarly, as in the case $\varkappa = 1$, we have

$$\begin{aligned} \overline{F_{S(v)}}(x) &= \sum_{n \in \text{supp}(v) \setminus \{0\}} \mathbb{P}\left(\bigcap_{k=1}^n \{S_k > x\}\right) \mathbb{P}(v = n) \\ &\leq \sum_{n \in \text{supp}(v) \setminus \{0\}} \mathbb{P}(S_1 > x) \mathbb{P}(v = n) \\ &= \overline{F_{X_1}}(x) \mathbb{P}(v \geq \varkappa). \end{aligned} \tag{23}$$

Estimates (22) and (23) imply that the asymptotic relation (6) holds for any possible \varkappa . In addition, we observe that, by Lemma 2, distribution $F_{S^{(v)}}$ belongs to \mathcal{H} together with F_{X_1} . Part (i) of the proposition is proved. \square

Proof of part (ii). The statement of this part follows immediately from the estimate (23) and Lemma 1 because

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{S^{(v)}}}(x) \leq \mathbb{P}(v \geq 1) \limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_1}}(x)$$

for any $\lambda > 0$. \square

5.5. Proof of Proposition 5

Proof of part (i). Proof of this part is similar to the proof of part (i) of Proposition 1. Namely, for $\lambda > 0$ and $K \geq 2$ by using (10), we obtain that

$$\begin{aligned} \mathbb{E} e^{\lambda S^{(v)}} &\geq \mathbb{E} \left(e^{\lambda S^{(v)}} \mathbb{1}_{\{v \leq K\}} \right) \\ &= \sum_{n=0}^K \mathbb{E} e^{\lambda S^{(n)}} \mathbb{P}(v = n) \\ &\geq \sum_{n=0}^K \mathbb{E} e^{\lambda S_n} \mathbb{P}(v = n) \\ &\geq \sum_{n=0}^K \Delta^n \mathbb{P}(v = n) \\ &= \mathbb{E} \left(e^{v \log \Delta} \mathbb{1}_{\{v \leq K\}} \right) \end{aligned}$$

with $\Delta = \Delta(\lambda) = \inf_{k \geq 1} \mathbb{E} e^{\lambda X_k} > 1$. The condition $F_v \in \mathcal{H}$ implies that

$$\lim_{K \rightarrow \infty} \mathbb{E} \left(e^{v \log \Delta} \mathbb{1}_{\{v \leq K\}} \right) = \infty.$$

Therefore, $\mathbb{E} e^{\lambda S^{(v)}} = \infty$ for an arbitrary $\lambda > 0$, i.e., $F_{S^{(v)}} \in \mathcal{H}$. Part (i) of the proposition is proved. \square

Proof of part (ii). The assertion of this part is obvious because condition $\inf_{k \geq 1} \mathbb{P}(X_k \geq a) = 1$ with $a > 0$ implies that $\inf_{k \geq 1} \mathbb{E} e^{\lambda X_k} > 1$ for any $\lambda > 0$. The details of this implication are presented in the proof of Proposition 1 (ii). \square

Proof of part (iii). For positive x , we have

$$\begin{aligned} \overline{F_{S^{(v)}}}(x) &= \sum_{n=1}^{\infty} \overline{F_{S^{(n)}}}(x) \mathbb{P}(v = n) \\ &= \sum_{n=1}^{\infty} \mathbb{P} \left(\bigcup_{k=1}^n \{S_k > x\} \right) \mathbb{P}(v = n) \\ &\geq \sum_{n=1}^{\infty} \mathbb{P}(S_1 > x) \mathbb{P}(v = n) \\ &= \overline{F_{X_1}}(x) \mathbb{P}(v \geq 1). \end{aligned} \tag{24}$$

The assertion of part (iii) follows now from Lemma 1 because by (24)

$$\limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{S^{(v)}}}(x) \geq \mathbb{P}(v \geq 1) \limsup_{x \rightarrow \infty} e^{\lambda x} \overline{F_{X_1}}(x)$$

for an arbitrary positive λ . \square

Proof of part (iv). Conditions of this part and Proposition 1 (parts (iii) and (iv)) imply that $F_{S_\nu} \in \mathcal{H}$. In addition, for positive x ,

$$\begin{aligned} \overline{F_{S^{(\nu)}}}(x) &= \sum_{n=1}^{\infty} \mathbb{P}(\max\{S_1, S_2, \dots, S_n\} > x) \mathbb{P}(\nu = n) \\ &\geq \sum_{n=1}^{\infty} \mathbb{P}(S_n > x) \mathbb{P}(\nu = n) \\ &= \overline{F_{S_\nu}}(x). \end{aligned}$$

Hence, $F_{S^{(\nu)}} \in \mathcal{H}$, according to the Lemma 2. Part (iv) of the proposition is proved. \square

Proof of part (v). Let $\lambda > 0$ be a positive number from the condition of part (v), i.e.,

$$\sup_{k \geq 1} \mathbb{E} e^{\lambda X_k} = \hat{c}_\lambda$$

with some positive constant \hat{c}_λ . For this λ , we have

$$\begin{aligned} \sup_{k \geq 1} \mathbb{E} e^{\lambda X_k^+} &= \sup_{k \geq 1} \mathbb{E} \left(e^{\lambda X_k^+} \mathbb{1}_{\{X_k \geq 0\}} + e^{\lambda X_k^+} \mathbb{1}_{\{X_k < 0\}} \right) \\ &= \sup_{k \geq 1} \mathbb{E} \left(e^{\lambda X_k} \mathbb{1}_{\{X_k \geq 0\}} + \mathbb{1}_{\{X_k < 0\}} \right) \\ &\leq \hat{c}_\lambda + 1, \end{aligned}$$

where $X_k^+ = X_k \mathbb{1}_{\{X_k \geq 0\}}$ for $k \in \{1, 2, \dots\}$. Due to Proposition 1(vi), d.f. $F_{S_\nu^+}$ belongs to the class \mathcal{H}^c with r.v. $S_\nu^+ = X_1^+ + \dots + X_\nu^+$.

According to the standard representation, for positive x , we have

$$\begin{aligned} \overline{F_{S^{(\nu)}}}(x) &= \sum_{n=1}^{\infty} \mathbb{P}(\max\{S_1, S_2, \dots, S_n\} > x) \mathbb{P}(\nu = n) \\ &\leq \sum_{n=1}^{\infty} \mathbb{P}(\max\{S_1^+, S_2^+, \dots, S_n^+\} > x) \mathbb{P}(\nu = n) \\ &= \sum_{n=1}^{\infty} \mathbb{P}(S_n^+ > x) \mathbb{P}(\nu = n) \\ &= \overline{F_{S_\nu^+}}(x). \end{aligned}$$

By applying Lemma 2, we obtain that d.f. $F_{S^{(\nu)}}$ is light-tailed due to the light tail of d.f. $F_{S_\nu^+}$. Part (v) of the proposition is proved. \square

6. Concluding Remarks

In this paper, we show that both heavy-tailed and light-tailed classes of distributions have quite a number of interesting properties related to the randomly stopped structures. Based on our results, various heavy-tailed or light-tailed distributions can be constructed. On the other hand, according to the propositions we proved, in most cases, it is easier to determine whether the considered distribution is light-tailed or heavy-tailed. The main novelty of our work consists in the fact that we study randomly stopped structures in a set of independent but possibly *differently distributed* primary random variables. In Section 1, it was mentioned that randomly stopped structures together with heavy-tailed distributions appear in such fields as insurance and financial activity, survival analysis, risk management, computer and communication networks, etc. Recently, many articles have been written on the heavy-tailed distributions, both in scientific and popular science journals. Let us mention a few such works. Heavy-tailed distributions applied to financial losses and stochastic returns are described and discussed in [50–52]. The influence of

heavy-tailed distributions on actuarial statistics is examined in [53–56]. The performance of heavy-tailed distributions in social and medical research is discussed in [57,58]. The application of heavy-tailed distributions of a special form to study computer systems and telecommunication networks is presented in [59–61]. More concretely, the results of the current paper related to the randomly stopped sums are applied not only to the standard areas such as insurance models ([62,63], etc.), but also to information ranking algorithms ([64,65]) and teletraffic arrivals [66].

From the content of the mentioned works, it can be seen that in many cases, it is quite difficult to fit heavy-tailed distributions to the real data. Therefore, our proposed transformations of heavy-tailed distributions increase the chances of choosing the right distribution. So, in our opinion, it makes sense to continue research on transformations for heavy-tailed distributions. In addition to the randomly stopped structures examined in this paper, moment transformations, random effects, and randomly stopped products can be considered, for instance.

Author Contributions: Conceptualization, R.L. and J.Š.; methodology, J.Š.; software, S.D.; validation, R.L., S.D. and J.K.; formal analysis, J.K.; investigation, S.D. and J.K.; resources, J.Š.; writing—original draft preparation, S.D.; writing—review and editing, R.L.; visualization, S.D. and J.K.; supervision, J.Š.; project administration, R.L.; funding acquisition, J.Š. and J.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Data are contained within the article.

Acknowledgments: The authors are very grateful to the editors and the anonymous referees for their constructive and valuable suggestions and comments which have helped to improve the previous version of the paper.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Albin, J.M.P. A note on the closure of convolution power mixtures (random sums) of exponential distributions. *J. Aust. Math. Soc.* **2008**, *84*, 1–7. [\[CrossRef\]](#)
2. Andrulytė, I.M.; Manstavičius, M.; Šiaulys, J. Randomly stopped maximum and maximum of sums with consistently varying distributions. *Mod. Stoch. Theory Appl.* **2017**, *4*, 65–78. [\[CrossRef\]](#)
3. Cheng, D.; Ni, F.; Pakes, A.G.; Wang, Y. Some properties of the exponential distribution class with application to risk theory. *J. Korean Stat. Soc.* **2012**, *41*, 515–527. [\[CrossRef\]](#)
4. Cline, D.B.H. Convolutions of the distributions with exponential tails. *J. Aust. Math. Soc.* **1987**, *43*, 347–365. [\[CrossRef\]](#)
5. Danilenko, S.; Šiaulys, J. Random convolution of O -exponential distributions. *Nonlinear Anal. Model.* **2015**, *20*, 447–454. [\[CrossRef\]](#)
6. Danilenko, S.; Šiaulys, J. Randomly stopped sums of not identically distributed heavy tailed random variables. *Stat. Probab. Lett.* **2016**, *113*, 84–93. [\[CrossRef\]](#)
7. Danilenko, S.; Markevičiūtė, J.; Šiaulys, J. Randomly stopped sums with exponential-type distributions. *Nonlinear Anal. Model.* **2017**, *22*, 793–807. [\[CrossRef\]](#)
8. Danilenko, S.; Paškauskaitė, S.; Šiaulys, J. Random convolution of inhomogeneous distributions with O -exponential tail. *Mod. Stoch. Theory Appl.* **2016**, *3*, 79–94. [\[CrossRef\]](#)
9. Danilenko, S.; Šiaulys, J.; Stepanauskas, G. Closure properties of O -exponential distributions. *Stat. Probab. Lett.* **2018**, *140*, 63–70. [\[CrossRef\]](#)
10. Dirma, M.; Nakliuda, N.; Šiaulys, J. Generalized moments of sums with heavy-tailed random summands. *Lith. Math. J.* **2023**, *63*, 254–271. [\[CrossRef\]](#)
11. Embrechts, P.; Goldie, C.M. On the closure and factorization properties of subexponential and related distributions. *J. Aust. Math. Soc. Ser. A* **1980**, *29*, 243–256. [\[CrossRef\]](#)
12. Geng, B.; Liu, Z.; Wang, S. A Kesten-type inequality for randomly weighted sums of dependent subexponential random variables with applications to risk theory. *Lith. Math. J.* **2023**, *63*, 81–91. [\[CrossRef\]](#)
13. Karasevičienė, J.; Šiaulys, J. Randomly stopped sums with generalized subexponential distribution. *Axioms* **2023**, *12*, 641. [\[CrossRef\]](#)
14. Karasevičienė, J.; Šiaulys, J. Randomly stopped minimum, maximum, minimum of sums and maximum of sums with generalized subexponential distributions. *Axioms* **2024**, *13*, 85. [\[CrossRef\]](#)
15. Kizinevič, E.; Sprindys, J.; Šiaulys, J. Randomly stopped sums with consistently varying distributions. *Mod. Stoch. Theory Appl.* **2016**, *3*, 165–179. [\[CrossRef\]](#)

16. Konstantinides, D.; Leipus, R.; Šiaulyš, J. A note on product-convolution for generalized subexponential distributions. *Nonlinear Anal. Model.* **2022**, *27*, 11054–11067. [[CrossRef](#)]
17. Konstantinides, D.; Leipus, R.; Šiaulyš, J. On the non-closure under convolution for strong subexponential distributions. *Nonlinear Anal. Model.* **2023**, *28*, 97–115. [[CrossRef](#)]
18. Leipus, R.; Šiaulyš, J. Closure of some heavy-tailed distribution classes under random convolution. *Lith. Math. J.* **2012**, *52*, 249–258. [[CrossRef](#)]
19. Leipus, R.; Šiaulyš, J. On the random max-closure for heavy-tailed random variables. *Lith. Math. J.* **2017**, *57*, 208–221. [[CrossRef](#)]
20. Lin, J.; Wang, Y. New examples of heavy-tailed O-subexponential distributions and related closure properties. *Stat. Probab. Lett.* **2012**, *82*, 427–432. [[CrossRef](#)]
21. Ragulina, O.; Šiaulyš, J. Randomly stopped minima and maxima with exponential-type distributions. *Nonlinear Anal. Model.* **2019**, *24*, 297–313. [[CrossRef](#)]
22. Shimura, T.; Watanabe, T. Infinite divisibility and generalized subexponentiality. *Bernoulli* **2005**, *11*, 445–469. [[CrossRef](#)]
23. Sprindys, J.; Šiaulyš, J. Regularly distributed randomly stopped sum, minimum and maximum. *Nonlinear Anal. Model.* **2020**, *25*, 509–522. [[CrossRef](#)]
24. Teicher, H. Moments of randomly stopped sums revisited. *J. Theor. Probab.* **1995**, *8*, 779–793. [[CrossRef](#)]
25. Tesemnikov, P.I. On the distribution tail of the sum of the maxima of two randomly sums in the presence of heavy tails. *Sib. Elektron. Mat. Izv.* **2019**, *16*, 1785–1794. [[CrossRef](#)]
26. Watanabe, T. Convolution equivalence and distributions of random sums. *Probab. Theory Relat. Fields* **2008**, *142*, 367–397. [[CrossRef](#)]
27. Watanabe, T. The Wiener condition and the conjectures of Embrechts and Goldie. *Ann. Probab.* **2019**, *47*, 1221–1239. [[CrossRef](#)]
28. Watanabe, T.; Yamamuro, K. Ratio of the tail of an infinity divisible distribution on the line to that of its Lévy measure. *Electron. J. Probab.* **2010**, *15*, 44–74. [[CrossRef](#)]
29. Xu, H.; Foss, S.; Wang, Y. Convolution and convolution-root properties of long-tailed distributions. *Extremes* **2015**, *18*, 605–628. [[CrossRef](#)]
30. Xu, H.; Wang, Y.; Cheng, D.; Yu, C. On the closure under infinitely divisible distribution roots. *Lith. Math. J.* **2022**, *62*, 258–287. [[CrossRef](#)]
31. Markovich, N.M.; Rodionov, I.V. Maxima and sums of non-stationary random length sequences. *Extremes* **2020**, *23*, 451–464. [[CrossRef](#)]
32. Leipus, R.; Šiaulyš, J.; Konstantinides, D. Minimum of heavy-tailed random variables is not heavy-tailed. *AIMS Math.* **2023**, *8*, 13066–13072. [[CrossRef](#)]
33. Andersen, E.S. On the collective theory of Risk in case of contagion between claims. *Bull. Inst. Math. Appl.* **1957**, *12*, 275–279.
34. Asmussen, S.; Albrecher, H. *Ruin Probabilities*, 2nd ed.; World Scientific: Singapore, 2010.
35. Denisov, D.; Foss, S.; Korshunov, D. Tail asymptotics for the supremum of random walk when the mean is not finite. *Queueing Syst.* **2004**, *46*, 15–33. [[CrossRef](#)]
36. Embrechts, P.; Klüppelberg, C.; Mikosch, T. *Modelling Extremal Events for Insurance and Finance*; Springer: New York, NY, USA, 1997.
37. Embrechts, P.; Veraverbeke, N. Estimates for the probability of ruin with special emphasis and the possibility of large claims. *Insur. Math. Econ.* **1982**, *1*, 55–72. [[CrossRef](#)]
38. Klüppelberg, C. Subexponential distributions and integrated tails. *J. Appl. Probab.* **1988**, *25*, 132–141. [[CrossRef](#)]
39. Bingham, N.H.; Goldie, C.M.; Teugels, J.L. *Regular Variation*; Cambridge University Press: Cambridge, UK, 1987.
40. Borovkov, A.A.; Borovkov, K.A. *Asymptotic Analysis of Random Walks: Heavy-Tailed Distributions*; Cambridge University Press: Cambridge, UK, 2008.
41. Foss, S.; Korshunov, D.; Zachary, S. *An Introduction to Heavy-Tailed and Subexponential Distributions*, 2nd ed.; Springer: New York, NY, USA, 2013.
42. Konstantinides, D.G. *Risk Theory: A Heavy Tail Approach*; World Scientific: Hackensack, NJ, USA, 2018.
43. Leipus, R.; Šiaulyš, J.; Konstantinides, D. *Closure Properties for Heavy-Tailed and Related Distributions: An Overview*; Springer: Cham, Switzerland, 2023.
44. Nair, J.; Wierman, A.; Zwart, B. *The Fundamentals of Heavy Tails: Properties, Emergence, and Estimation*; Cambridge University Press: Cambridge, UK, 2022.
45. Liu, Y. A general treatment of alternative expectation formulae. *Stat. Probab. Lett.* **2020**, *166*, 108863. [[CrossRef](#)]
46. Hall, P. The distribution of means for samples of sizes N drawn from a population in which the variate takes values between 0 and 1, all such values being equally probable. *Biometrika* **1927**, *19*, 240–245. [[CrossRef](#)]
47. Irwin, J.O. On the frequency distribution of the means of samples from a population having any law of frequency with finite moments, with special reference to Pearson's type II. *Biometrika* **1927**, *19*, 225–239. [[CrossRef](#)]
48. Johnson, N.L.; Kotz, S.; Balakrishnan, N. *Continuous Univariate Distributions*, 2nd ed.; Wiley: New York, NY, USA, 1995; Volume 2.
49. Tadikamalla, Pandu R. A look at the Burr and related distributions. *Int. Stat. Rev.* **1980**, *48*, 337–344. [[CrossRef](#)]
50. Chen, H.; Fan, K. Tail Value-at-Risk based profiles for extreme risks and their application in distributionally robust portfolio selections. *Mathematics* **2023**, *11*, 91. [[CrossRef](#)]
51. Mehta, M.J.; Yang, F. Portfolio optimization for extreme risks with maximum diversification: An empirical analysis. *Risks* **2022**, *10*, 101. [[CrossRef](#)]

52. Sepanski, J.H.; Wang, X. New classes of distortion risk measures and their estimation. *Risks* **2023**, *11*, 194. [[CrossRef](#)]
53. Mahdavi, A.; Kharazm, O.; Contreras-Reyes, J.E. On the contaminated weighted exponential distribution: Applications to modelling insurance claim data. *J. Risk Financ. Manag.* **2022**, *15*, 500. [[CrossRef](#)]
54. Olmos, N.M.; Gómez-Déniz, E.; Venegas, O. The heavy-tailed Gleser model: Properties, estimation, and applications. *Mathematics* **2022**, *10*, 4577. [[CrossRef](#)]
55. Grün, B.; Miljkovic, T. Extending composite loss models using a general framework of advanced computational tools. *Scand. Actuar. J.* **2019**, *2019*, 642–660. [[CrossRef](#)]
56. Marambakuyana, W.A.; Shongwe, S.C. Composite and mixture distributions for heavy-tailed data—An application to insurance claims. *Mathematics* **2024**, *12*, 335. [[CrossRef](#)]
57. Klebanov, L.B.; Kuvaeva-Gudoshnikova, Y.V.; Rachev, S.T. Heavy-tailed probability distributions: Some examples of their appearance. *Mathematics* **2023**, *11*, 3094. [[CrossRef](#)]
58. Santoro, K.I.; Gallardo, D.I.; Venegas, O.; Cortés, I.E.; Gomes, H.W. A heavy-tailed distribution based on the Lomax-Reyleigh distribution with application to medical data. *Mathematics* **2023**, *11*, 4626. [[CrossRef](#)]
59. Markovich, N.; Vaičiulis, M. Extreme value statistics for envolving random networks. *Mathematics* **2023**, *11*, 2171. [[CrossRef](#)]
60. Rusev, V.; Skorikov, A. The asymptotics of moments for the remaining time of heavy-tail distributions. *Comput. Sci. Math. Forum* **2023**, *7*, 52. [[CrossRef](#)]
61. Sousa-Vieira, M.E.; Fernández-Veiga, M. Study of coded ALOHA with multi-user detection under heavy-tailed and correlated arrivals. *Future Internet* **2023**, *15*, 132. [[CrossRef](#)]
62. Klüppelberg, C.; Mikosch, T. Large deviations of heavy-tailed random sums with applications in insurance and finance. *J. Appl. Probab.* **1997**, *34*, 293–308. [[CrossRef](#)]
63. Aleškevičienė, A.; Leipus, R.; Šiaulyš, J. Tail behavior of random sums under consistent variation with applications to the compound renewal risk model. *Extremes* **2008**, *11*, 261–279. [[CrossRef](#)]
64. Olvera-Cravioto, M. Asymptotics for weighted random sums. *Adv. Appl. Probab.* **2012**, *44*, 1142–1172. [[CrossRef](#)]
65. Volkovich, Y.; Litvak, N. Asymptotic analysis for personalized Web search. *Adv. Appl. Probab.* **2010**, *42*, 577–604. [[CrossRef](#)]
66. Faÿ, G.; González-Arévalo, B.; Mikosch, T.; Samorodnitsky, G. Modeling teletraffic arrivals by a Poisson cluster process. *Queueing Syst.* **2006**, *54*, 121–140. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.