

Supplementary File S3: Participants and Procedure

Article title: Virtual reality-based early neurocognitive stimulation in critically ill patients: A pilot randomized clinical trial

Authors: Guillem Navarra-Ventura, Gemma Gomà, Candelaria de Haro, Mercè Jodar, Leonardo Sarlabous, David Hernando, Raquel Bailón, Ana Ochagavía, Lluís Blanch, Josefina López-Aguilar*, & Sol Fernández-Gonzalo*

* Co-senior authors

Corresponding author: Guillem Navarra-Ventura (gnavarra@tauli.cat)

Critical Care Center, Hospital Universitari Parc Taulí, Institut d'Investigació i Innovació Parc Taulí I3PT, Universitat Autònoma de Barcelona, 08208 Sabadell, Catalonia (Spain). Tel. +34 937 23 66 73

Journal name: Journal of Personalized Medicine

Methods

Participants and Procedure

Previous cognitive impairment or dementia was checked in the patients' clinical records and by interviewing the relatives. In patients aged ≥ 60 , pre-existing cognitive impairment was recorded using the short form of the Spanish version of the Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE) [1] filled in by relatives. Patients with scores ≥ 3.56 were excluded as they were at risk of probable cognitive impairment.

References

1. Jorm AF (2004) The Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE): a review. *Int Psychogeriatrics* 16:275–293