

Table S2 Log-linear regression analysis including only the female stratum, of the association between symptom cluster (mutually adjusted) and the FAS-Score as dependent variable (n=245)

Characteristic	Change in Percent	95% CI ¹	p-value	q-value ²
Clustered symptoms: Loss of sense: taste and/ or smell				
No	—	—		
Yes	5.13	-0.04, 0.14	0.3	0.4
Clustered symptoms: Ear, nose and throat	0.00	-0.02, 0.03	0.6	0.7
Clustered symptoms: Cardiopulmonary	3.05	0.00, 0.07	0.048	0.2
Clustered symptoms: Cognitive and mental	6.18	0.03, 0.08	<0.001	<0.001
Clustered symptoms: Locomotor system	1.01	-0.03, 0.05	0.6	0.7
Clustered symptoms: Gastrointestinal	-1.00	-0.04, 0.03	0.7	0.7
Clustered symptoms: Eyes/ Hair/ Skin/ Stings in arms & legs	3.05	-0.01, 0.08	0.2	0.3
Age (years)	0.00	-0.01, 0.00	0.054	0.2
Body mass index (kg/m ²)	1.01	0.00, 0.01	0.2	0.3
Smoker status				
Never smoked	—	—		
Ex-smoker	4.08	-0.04, 0.13	0.3	0.5
Current smoker	-1.98	-0.15, 0.12	0.8	0.8
Prior diagnosis of depression disorder				
No	—	—		
Yes	12.75	-0.01, 0.25	0.080	0.3
Prior diagnosis of anxiety disorder				
No	—	—		
Yes	12.75	-0.04, 0.28	0.14	0.3

¹CI = Confidence Interval

²False discovery rate correction for multiple testing