

Table S1 Log-linear regression analysis including only the male stratum: Association between symptom cluster (mutually adjusted) and the FAS-Score as dependent variable (n=201)

Characteristic	Change in Percent	95% CI ¹	p-value	q-value ²
Clustered symptoms: Loss of sense: taste and/ or smell				
No	—	—		
Yes	-1.00	-0.09, 0.08	0.9	0.9
Clustered symptoms: Ear, nose and throat	0.00	-0.02, 0.03	0.7	0.8
Clustered symptoms: Cardiopulmonary	2.02	-0.03, 0.06	0.5	0.8
Clustered symptoms: Cognitive and mental	5.13	0.02, 0.08	<0.001	0.009
Clustered symptoms: Locomotor system	-1.00	-0.05, 0.03	0.5	0.8
Clustered symptoms: Gastrointestinal	3.05	-0.01, 0.08	0.12	0.5
Clustered symptoms: Eyes/ Hair/ Skin/ Stings in arms & legs	2.02	-0.03, 0.08	0.5	0.8
Age (years)	0.00	0.00, 0.00	0.4	0.8
Body mass index (kg/m ²)	0.00	-0.01, 0.01	0.8	0.8
Smoker status				
Never smoked	—	—		
Ex-smoker	2.02	-0.06, 0.11	0.6	0.8
Current smoker	2.02	-0.11, 0.15	0.8	0.8
Prior diagnosis of depression disorder				
No	—	—		
Yes	7.25	-0.12, 0.26	0.5	0.8
Prior diagnosis of anxiety disorder				
No	—	—		
Yes	29.69	0.04, 0.49	0.020	0.13

¹CI = Confidence Interval

²False discovery rate correction for multiple testing