

Article

Volumetric Biomarkers of Visual Outcome after Surgical Repair in Lamellar Macular Holes

Myrta Lippera ¹, George Moussa ¹, Tsveta Ivanova ¹, Mariantonia Ferrara ^{2,3,4}, Karina Spiess ¹, Naseer Ally ⁵, Kirti Jasani ¹, Felipe Dhawahir-Scala ¹, Niall Patton ¹ and Assad Jalil ^{1,*}

¹ Manchester Royal Eye Hospital, Manchester University Hospitals NHS Foundation Trust, Manchester M13 9WL, UK; karinaspiess@gmx.net (K.S.); kirti.jasani@mft.nhs.uk (K.J.)

² School of Medicine, University of Malaga, 29071 Malaga, Spain

³ Department of Medical and Surgical Specialties, Radiological Sciences and Public Health, University of Brescia, 25123 Brescia, Italy

⁴ Eye Unit, ASST Spedali Civili di Brescia, Piazzale Spedali Civili, 25123 Brescia, Italy

⁵ Division of Ophthalmology, Department of Neurosciences, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, 7 York Road, Parktown, Johannesburg 2193, South Africa

* Correspondence: assad.jalil@mft.nhs.uk

Abstract: Background: We investigate novel OCT parameters, based on the volumetric analysis of lamellar macular holes (LMHs), as prognostic indicators for visual outcomes after surgery. Methods: LMHs were divided into degenerative LMHs (D-LMHs) and ERM-foveoschisis (ERM-FS). Pre-operative clinical, OCT linear and volumetric parameters were collected. Volumes were obtained using the OCT automatic segmentation, such as central retinal volume (CRV) and outer nuclear layer (ONL) volume, or using a novel method to calculate volumes of specific LMH entities like epiretinal proliferation (ERP), foveal cavity (FC) in D-LMH and schitic volume (SV) in ERM-FS. Univariate and multivariate linear regression analysis evaluated the factors predictive for post-operative best-corrected visual acuity (BCVA). Results: We included 31 eyes of 31 patients (14 D-LMH, 17 ERM-FS). A pre-operative BCVA ≤ 0.48 logMAR was a predictor for achieving ≤ 0.30 logMAR at final follow-up. A lower pre-operative BCVA ($p = 0.008$) and the presence of ERP ($p = 0.002$) were associated with worse visual outcomes post-surgery. Moreover, novel pre-operative OCT parameters significantly associated with worse post-operative BCVA, such as increased FC volume ($p = 0.032$) and lower CRV ($p = 0.034$) in the D-LMH subtype and lower CRV ($p < 0.001$) and ERP volume ($p < 0.001$), higher SV ($p < 0.001$) and foveal ONL volume ($p < 0.001$) in the ERM-FS subtype. Conclusions: Novel volumetric OCT parameters can be prognostic indicators of visual outcome following surgery in LMHs.

Keywords: lamellar macular hole (LMH); optical coherence tomography (OCT); predictive prognostic factors; biomarkers; retinal layer segmentation; retinal volume; volumetric analysis



Citation: Lippera, M.; Moussa, G.; Ivanova, T.; Ferrara, M.; Spiess, K.; Ally, N.; Jasani, K.; Dhawahir-Scala, F.; Patton, N.; Jalil, A. Volumetric Biomarkers of Visual Outcome after Surgical Repair in Lamellar Macular Holes. *J. Pers. Med.* **2024**, *14*, 755. <https://doi.org/10.3390/jpm14070755>

Academic Editor: Ciprian Danielescu

Received: 21 June 2024

Revised: 7 July 2024

Accepted: 11 July 2024

Published: 16 July 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Lamellar macular hole (LMH) is morphologically characterized by the presence of a partial defect in the inner foveal layers, not extending to the entire retina, with irregular foveal contour [1–3]. With the advent of optical coherence tomography (OCT), different classifications and subtypes of LMH have been proposed [1–4]. More recently, an OCT-based consensus renamed two clinical subtypes: “epiretinal membrane-foveoschisis” (ERM-FS), characterized by the presence of a contractile ERM and foveoschisis at the level of Henle fiber layer (HFL) and a (degenerative) LMH, characterized by a foveal cavity with undermined edges and signs evoking loss of foveal tissue [2]. The pathophysiology of LMH is still largely unknown [2,5]; however, two main mechanisms have been described: (i) a contractile membrane causing traction in ERM-FS and (ii) loss of retinal tissue in D-LMH [2]. Both mechanisms produce asymmetric three-dimensional (3D) changes on the macula.

LMH is a slowly progressive condition that can impair visual acuity (VA) in a subset of patients [6]. Although pars plana vitrectomy (PPV) and internal limiting membrane (ILM) peeling is the established surgical approach for LMH, there is no general consensus regarding quantitative parameters that can help decide on the timing of surgery or predict the visual outcome [7–9]. It has been highlighted that LMH is an asymmetric, 3D disease of the macula, better defined by volumetric analysis rather than linear measurements [6]. Most previous studies on LMH, were limited by the analysis of measurements from linear OCT macular scans, which do not provide adequate information on retinal tissue remodeling [6,10,11].

The aim of our study was to investigate novel OCT parameters, based on the volumetric analysis of ERM foveoschisis (ERM-FS) and degenerative LMH (D-LMH), as prognostic indicators for post-operative best-corrected visual acuity (BCVA) logMAR after surgery. Additionally, due to the functional importance of 0.30 logMAR BCVA, such as for driving, we investigate the effect of pre-operative BCVA on achieving this threshold postoperatively.

2. Materials and Methods

This is a retrospective interventional, single-center case series that adhered to the guidelines of the Declaration of Helsinki. Under UK guidance, retrospective data collection is regarded as an audit for the purpose of service evaluation, and as such ethical approval was not required. Diagnosis and treatment were conducted according to local guidelines with no new or experimental protocols. Clinical records were extracted from an electronic surgical database of consecutive eyes that underwent PPV for LMH repair at Manchester Royal Eye Hospital, UK, from January 2020 to January 2023. Eyes with LMH which had pre-operative macular OCT using the “dense macular volume” scan with Heidelberg Spectralis (Engineering GmbH, Heidelberg, Germany), were included. Exclusion criteria were (1) high myopia (more than 6 diopters); (2) advanced glaucoma; (3) any concomitant retinal disease involving the macula, such as diabetic maculopathy, retinal vein occlusion, age-related macular degeneration; (4) any comorbidity potentially impacting on final functional outcomes, such as uveitis or amblyopia; (5) history of trauma; (6) previous intraocular surgery other than cataract surgery performed more than 6 months before vitrectomy; (7) OCT with poor image quality; and (8) post-operative follow-up (FU) less than 4 weeks.

We subsequently divided the included eyes in two subtypes, degenerative LMH (D-LMH) and ERM foveoschisis (ERM-FS), based on the OCT-based consensus definition for LMH [2], as defined and shown in Figure 1.




Figure 1. Sub-categories of lamellar macular holes. (A) Degenerative lamellar macular hole (D-LMH): defined by the presence of an irregular foveal contour (the normal smooth and regular contour of the

fovea is disrupted), a foveal cavity with undermined edges (a cavity or space, highlighted by the white dashed line, is observed in the fovea with edges that appear undermined) and apparent loss of retinal tissue on OCT. (B) Epiretinal membrane foveoschisis (ERM-FS): defined by the presence of a contractile ERM (an ERM can be detected on the retinal surface, exerting traction on the retina) with foveoschisis (the schisis within the retinal layers, specifically at the level of Henle's fiber layer due to the traction from the ERM, is highlighted by the white dashed line). In summary, key differentiating features between D-LMH and ERM-FS include: ERM presence (ERM-FS is defined by the presence of a contractile ERM, while D-LMH might lack this feature); foveal schisis vs. cavity (in ERM-FS, the foveal splitting or schisis is due to traction, often at Henle's fiber layer, whereas in D-LMH, the foveal cavity with undermined edges is due to degeneration); retinal tissue loss (D-LMH typically shows apparent loss of retinal tissue, whereas ERM-FS might not show this degree of tissue loss but rather separation or schisis within layers); and contour changes (the irregular foveal contour is more pronounced in D-LMH due to degeneration, while in ERM-FS, the changes are primarily due to the tractional effects of the ERM).

For each patient, the following data were collected:

- I. Pre-operative characteristics: baseline characteristics such as age, gender, laterality; clinical characteristics such as lens status, pre-operative BCVA (logMAR), ocular findings at dilated funduscopy; and pre-operative OCT-based characteristics as detailed below.
- II. Surgical characteristics and intraoperative complications.
- III. Post-operative characteristics: post-operative BCVA (logMAR) and complications. The ETDRS (Early Treatment Diabetic Retinopathy Study) logMAR (logarithm of the Minimum Angle of Resolution) test was the standardized visual acuity test used to assess a patient's pre- and post-operative vision.

2.1. Surgical Technique

Patients with LMH were considered for surgery if they were symptomatic with significant metamorphopsia and/or worsening of BCVA or documented progression of LMH on the OCT. Small-gauge 25G PPV was performed in all cases. For all phakic patients, cataract surgery was concurrently performed. After vitrectomy, ERM and complete ILM peeling were completed. Air or 20% Sulfur-Hexafluoride (SF₆) were used as intraocular tamponade depending on surgeon preference.

2.2. OCT Parameters

The OCT examination included a macular volume scan with a 49-line horizontal raster covering an area of 30° by 30° (approximately 125 μm spacing between each scan, depending on the axial length of the eye being examined) centered on the fovea. For each scan, two vitreoretinal experts evaluated the ellipsoid zone (EZ) and external limiting membrane (ELM) and classified them as "normal", "disrupted" and "absent". Subsequently, using the caliper tool present, the two vitreoretinal experts measured the minimum retinal thickness (MRT), defined as the smallest distance between the retinal pigment epithelium (RPE) and the inner border of the retinal tissue on the fovea on a line perpendicular to the RPE measured on the linear OCT scan dissecting the fovea; the minimum and maximal linear horizontal diameter of the foveal cavity (FC) in D-LMH and of the foveoschisis in ERM-FS defined as the shortest and longest distance measured horizontally across the foveal cavity or the foveoschisis on the linear OCT scan dissecting the fovea.

Since the OCT machine does not have the capacity to automatically segment and delineate specific LMH entities like FC in D-LMH, the volume of the schitic cavities described as schitic volume (SV) in ERM-FS, or epiretinal proliferation (ERP), we adopted a novel and manual technique to calculate the volumes of those specific entities, which was validated on healthy and pathological eyes [12]. In brief, we calculate the surface area of the specific entity studied for each linear OCT scan (Figure 2). Finally, the specific volume

was calculated by multiplying the sum of the areas by the distance between the horizontal b-scans, using the formula $\text{Volume (mm}^3) = \sum \text{area [mm}^2] \times \text{OCT-scan distance [mm]}$.




Figure 2. Collection of OCT parameters used to calculate the volumes of specific LMH entities. The areas in yellow are summed and multiplied for the distance between the OCT scans to calculate the estimated volume of a specific LMH entity. (A) A specific tool was used to calculate the area of the FC in D-LMH on every single linear OCT scan. Using the formula $\text{Volume (mm}^3) = \sum \text{area [mm}^2] \times \text{OCT-scan distance [mm]}$, the volume of the FC in D-LMH was calculated. (B) A specific tool was used to calculate the area of the SV in ERM-FS on every single linear OCT scan. Using the formula $\text{Volume (mm}^3) = \sum \text{area [mm}^2] \times \text{OCT-scan distance [mm]}$, the SV in ERM-FS was calculated. (C) A specific tool was used to calculate the area of the ERP on every single linear OCT scan. Using the formula $\text{Volume (mm}^3) = \sum \text{area [mm}^2] \times \text{OCT-scan distance [mm]}$, the volume of the ERP was calculated.

As the last step, after review of the correct OCT automatized segmentation of the retinal layers, the following parameters, calculated by the OCT machine, were collected from the “Thickness Map” tabature of the OCT software (V 1.0): the central retinal thickness (CRT), defined as the average linear thickness in the central circle of the 1, 3, 6 ETDRS circle diameters; the central retinal volume (CRV), defined as the volume of the retinal tissue

included in the 30° by 30° area scanned; the average ONL thickness, defined as the average thickness of the ONL included in the central circle of the 1, 3, 6 ETDRS circle diameters; and the volume of the ONL within a diameter of 1 mm centered in fovea, defined as the volume of the ONL included in the central circle of the 1, 3, 6 ETDRS circle diameters.

2.3. Statistical Analysis

The statistical analysis was executed using IBM SPSS Statistics for Windows, Version 29.0 (IBM Corp, Armonk, NY, USA). Statistical significance was defined as $p < 0.05$. First, continuous variables were estimated for normality using the Shapiro–Wilk test and the mean (standard deviation) was reported for normally distributed variables; otherwise, for skewed data, we reported the median (interquartile range). We performed a paired *t*-test between pre-operative and post-operative continuous variables. Fisher’s exact test was performed to compare nominal variables. To build a model of predictive factors from the initial OCT-based parameters, a multivariable linear regression analysis was conducted with post-operative BCVA (logMAR) as the dependent variable. The lens status, presence and volume of ERP, FC volume in D-LMH and SV in ERM-FS, CRV, foveal ONL volume and pre-operative BCVA (logMAR) were used for the regression model. To demonstrate the different effect of volumetric analyses on D-LMH and ERM-FS, we added interaction terms for foveal ONL volume, ERP volume and CRV as moderators. The 95% confidence interval and *p* values were generated following a 16,000 sample Wild Bootstrap. A Receiver Operating Characteristic (ROC) curve using Area Under Curve (AUC) analysis was performed to report on the sensitivity and specificity of a cut-off threshold based on pre-operative BCVA as a predictor for achieving ≤ 0.30 logMAR (6/12 Snellen) post-operatively.

3. Results

We include 31 eyes of 31 patients with LMH (14 D-LMH, 17 ERM-FS). The mean (standard deviation) age at presentation was 69 [9] years old. Ten patients (32%) were males. Combined vitrectomy with cataract surgery was performed in all phakic patients, which resulted in 21 eyes (68%). No intra or post-operative complications were identified. The baseline and surgical characteristics are found in Table 1.

Table 1. Baseline demographics, OCT parameters and visual outcomes of eyes with LMHs, further divided in D-LMH and ERM-FS groups. The table shows the pre-operative parameters collected and the visual outcomes of the total number of eyes included in the study and of the two separated groups (ERM-FS and D-LMH). A comparison for each single parameter was performed between the D-LMH and the ERM-FS group, whose *p* values are reported in the last column of the table and statistical significance is highlighted in bold. Relative to D-LMH, ERM-FS had lower proportion of ERP (four [24%] compared to nine [64%]) ($p = 0.033$), a lower ERP volume (0.029 compared to 0.078 mm³) ($p = 0.044$) and higher foveal ONL average thickness and volume (113 μm compared to 90 μm, $p < 0.001$ and 0.089 mm³ compared to 0.07 mm³, $p = 0.002$). No statistically significant difference was found for the other parameters evaluated.

	Total	D-LMH	ERM-FS	<i>p</i> Value
Total	31	14	17	-
Baseline Characteristics				
Age (years)	69 (9)	65 (8)	73 (8)	0.016
Gender (% male)	10 (32%)	6 (43%)	4 (24%)	0.441
Laterality (% right)	14 (45%)	8 (57%)	6 (35%)	0.289
Phakic (% yes)	21 (68%)	12 (86%)	9 (53%)	0.068

Table 1. Cont.

	Total	D-LMH	ERM-FS	p Value
Total	31	14	17	-
Ellipsoid Zone				
Normal	20 (65%)	7 (50%)	13 (77%)	0.153
Disrupted	8 (26%)	6 (43%)	2 (12%)	0.097
Absent	3 (10%)	1 (7%)	2 (12%)	1.000
External Limiting Membrane				
Normal	22 (71%)	9 (64%)	13 (77%)	0.693
Disrupted	6 (19%)	3 (21%)	3 (18%)	1.000
Absent	3 (10%)	2 (14%)	1 (6%)	0.576
ERP (% yes)	13 (42%)	9 (64%)	4 (24%)	0.033
Pre-operative BCVA (logMAR)	0.49 (0.22)	0.50 (0.24)	0.49 (0.20)	0.897
Central Retinal Thickness	197 (82)	166 (41)	223 (99)	0.052
Minimum Retinal Thickness	155 (74)	110 (36)	191 (78)	0.001
Cavity or schisis max horizontal diameter	1292 (617)	967 (473)	1560 (603)	0.006
Cavity or schisis min horizontal diameter	428 (212)	490 (249)	377 (166)	0.139
ONL Average Thickness	102 (20)	90 (13)	113 (19)	<0.001
Foveal Cavity Volume	0.0716 (0.0426)	0.0716 (0.0426)	-	-
Schitic Volume	0.1057 (0.0860)	-	0.1057 (0.0860)	-
ERP Volume	0.0632 (0.0811)	0.0783 (0.0929)	0.0292 (0.0324)	0.044
Foveal ONL Volume	0.0803 (0.0176)	0.0700 (0.0118)	0.0888 (0.0173)	0.002
Central Retinal Volume	2.7945 (0.4031)	2.6509 (0.4442)	2.9127 (0.3339)	0.071
Visual outcomes				
Post-operative BCVA (logMAR)	0.37 (0.19)	0.42 (0.18)	0.32 (0.20)	0.184
LogMAR gain	0.12 (0.17)	0.08 (0.14)	0.16 (0.18)	0.184

Legend. ERP: epiretinal proliferation; ONL: outer nuclear layer; BCVA: best corrected visual acuity. Continuous variables are reported as mean (standard deviation, SD) and compared using independent *t*-test. Nominal variables were compared using Fisher’s exact test.

3.1. Pre-Operative Clinical Parameters and Visual Outcomes

The median follow-up period was of 69 days. While in the D-LMH group, BCVA improved from 0.50 (0.24) logMAR pre-operatively to 0.42 (0.18) logMAR at last follow-up ($p = 0.056$); in the ERM-FS group, BCVA improved from 0.49 (0.20) logMAR to 0.32 (0.20) logMAR ($p = 0.002$). Across the whole cohort, mean pre-operative BCVA improved from 0.48 (0.22) logMAR to 0.36 (0.19) logMAR ($p < 0.001$). Pre-operative and post-operative BCVA had a significant correlation (Supplementary Materials, Figure S1). Regarding the potential confounding effect of cataract surgery on visual outcomes, we do not report significant differences in pre-operative ($p = 0.447$), post-operative BCVA ($p = 0.195$) or logMAR gain ($p = 0.643$) between patients that were phakic ($n = 21$) or pseudophakic ($n = 10$) pre-operatively (see table, Supplementary Materials, Table S1). BCVA significantly improved in both patients that were phakic pre-operatively (from 0.47 [0.22] logMAR to 0.33 [0.16] logMAR) ($p = 0.003$) and in patients that were pseudophakic preoperatively (from 0.53 [0.22] logMAR to 0.43 [0.24] logMAR) ($p = 0.049$). Furthermore, we included pre-operative lens status as factor in our multivariate linear regression model and found that cataract surgery was not linked to significant improvement in post-operative BCVA (Table 2).

Table 2. Univariate and multivariate linear regression model for post-operative BCVA. On univariate tests, ERP, increased volume of schitic cavity and worse pre-operative BCVA were associated with worse post-operative BCVA. On multivariate linear regression, the presence of ERP and worse pre-operative BCVA were associated with worse visual outcomes. Interaction terms show that lower retinal volume was associated with worse visual outcomes in both subtypes. Specifically, for ERM-FS, ERP volume, increased SV and ONL volume were linked to worse post-operative BCVA. For D-LMH, we found that larger FC was associated with worse post-operative BCVA.

Independent Variable	Univariate		Multivariate	
	Post-Operative BCVA (r)	p Value	B Coefficient (95% CI)	p Value
Pre-operative BCVA (logMAR)	0.666	<0.001	0.316 (0.155 to 0.477)	0.008
Combined	0.33 (0.16) *	0.195 *	−0.064 (−0.123 to −0.008)	0.118
Phacovitrectomy				
Pseudophakic baseline	0.43 (0.24) *		-	-
ERP (present)	0.47 (0.19) *	0.006 *	0.155 (0.082 to 0.227)	0.002
ERP (absent)	0.29 (0.15) *		-	-
Volume of ERP	0.203	0.274	-	-
D-LMH	0.169	0.564	−0.793 (−1.629 to 0.013)	0.226
ERM-FS	0.117	0.655	−7.116 (−8.401 to −5.765)	<0.001
Volume of Foveal Cavity (D-LMH)	0.033	0.91	1.335 (0.425 to 2.245)	0.032
Schitic Volume (ERM-FS)	0.618	0.008	1.197 (0.831 to 1.551)	<0.001
Retinal Volume	−0.272	0.138	-	-
D-LMH	−0.141	0.63	−0.201 (−0.332 to −0.076)	0.034
ERM-FS	−0.284	0.27	−0.370 (−0.509 to −0.232)	<0.001
ONL Volume	−0.004	0.981	-	-
D-LMH	0.089	0.763	4.994 (−1.241 to 11.104)	0.236
ERM-FS	0.194	0.457	9.486 (7.291 to 11.416)	<0.001

Multivariate Model: R² = 0.888, adjusted R²: 0.824. Univariate analysis Pearson correlation (* mean [standard deviation] logMAR reported, independent t-test). Significance defined as p < 0.05 and highlighted in bold. Presented 95% confidence interval and p values are calculated following 16,000 resampling Wild Bootstrap.

3.2. Pre-Operative OCT Parameters and Visual Outcomes

Using the univariate tests, worse pre-operative BCVA ($p < 0.001$), presence of ERP ($p = 0.006$), and increased SV in ERM-FS ($p = 0.008$) were associated with worse functional outcomes after surgery (Table 2). No statistically significant correlation was found with post-operative BCVA for the following: sex ($p = 0.241$), age, lens status ($p = 0.195$), CRT ($p = 0.212$), MRT ($p = 0.100$), ONL thickness ($p = 0.679$), pre-operative status of EZ or ELM (respectively $p = 0.097$ and $p = 0.447$ for normal, $p = 0.252$ and $p = 0.832$ for disrupted and $p = 0.339$ and $p = 0.141$ for absent), and horizontal diameter of FC in D-LMH or foveoschisis in ERM-FS ($p = 0.166$ for maximum, $p = 0.337$ for minimum horizontal diameter). With the multivariate linear regression model following a Wild Bootstrap (16,000 resamples), presence of ERP ($p = 0.002$) and worse pre-operative BCVA ($p = 0.008$) were confirmed as significantly associated with worse final visual function. Moreover, in the subtype of ERM-FS, a lower CRV ($p < 0.001$) and ERP Volume ($p < 0.001$), a higher SV ($p < 0.001$) and foveal ONL volume ($p < 0.001$) were significantly associated with worse visual outcomes after surgery. In the D-LMH subtype, increased FC volume ($p = 0.032$) and lower CRV ($p = 0.034$) were significantly associated with worse post-operative BCVA (Table 2, Figure 3). The regression model could explain 88.8% of variability in post-operative BCVA (adjusted r^2 : 0.842). In contrast pre-operative BCVA alone could explain 44.4% of the variability in post-operative BCVA (adjusted r^2 : 0.425). Standardized Beta coefficients demonstrate the relative importance of each independent and moderator variable for post-operative BCVA (dependent variable). By reporting the standardized coefficients (Figure 3), we found that CRV in both subtypes and foveal ONL volume in ERM-FS were the most significant moderators in predicting post-operative BCVA.




Figure 3. Multivariable linear regression standardized Beta coefficients and 95% confidence intervals for post-operative BCVA. This figure presents the results of a multivariable linear regression analysis examining the relationship between various predictors and post-operative BCVA. The 95% CIs provide a range of values within which we can be 95% confident that the true Beta coefficient lies. The standardized Beta coefficients represent the strength and direction of the relationship between each predictor and the post-operative BCVA, with larger absolute values indicating stronger relationships. Indeed, the table shows that retinal volume and ONL volume were the most important parameters evaluated for post-operative BCVA, as demonstrated by the higher standardized Beta coefficients, reflecting the relative importance of the independent variables for final visual outcome. The negative values of standardized Beta coefficients for pre-operative retinal volume highlight that higher retinal volumes in both D-LMH and ERM-FS subgroups are related to better post-operative BCVA (see vertical arrows on the right side of the figure). On the contrary, the positive values of standardized Beta coefficients for pre-operative ONL volume highlight that higher ONL volumes are related to worse post-operative BCVA (however, this result is statistically significant only for the ERM-FS subgroup as shown by the *p* values on the top of the figure). Legend: ONL: outer nuclear layer; ERP: epiretinal proliferation; D-LMH: degenerative lamellar macular hole; ERM-FS: epiretinal membrane foveoschisis; FC: foveal cavity; Pre-op: pre-operative. Statistical significance in bold (*p* < 0.05). The 95% CI and *p* values are based on 16,000 Wild Bootstrap.

A correlation matrix demonstrated that, at baseline, the increased volume of ERP strongly correlates to increased SV (*p* = 0.005), reduced CRV (*p* = 0.002) and foveal ONL volume (*p* = 0.008), lower MRT (*p* = 0.014) and ONL thickness (*p* = 0.007). Among other correlations between the predictors determined in the study, only CRV showed correlation with ONL volume (*p* < 0.001) (Supplementary Materials, Table S2). Additionally, we found an association between low pre-operative BCVA and absence of pre-operative EZ (*p* = 0.046) and presence of ERP with both poor pre- and post-operative BCVA (*p* = 0.026 and *p* = 0.006, respectively) (Supplementary Materials, Table S1).

Finally, through ROC curve and using AUC analysis, we determined a cut-off pre-operative BCVA of ≤0.48 logMAR as a predictor for achieving 0.30 logMAR at final follow up (sensitivity: 71.4%, specificity: 88.2%, AUC 0.828 [95% CI 0.681 to 0.974], *p* = 0.002, Youden Index [J]: 0.596) (Figure 4A). While significance was maintained for eyes with pre-operative BCVA > 0.30 logMAR, there was no significant change in BCVA in patients with good pre-operative BCVA (<0.30 logMAR) (*p* = 0.967). Eight (26%) of thirty-one eyes had 0.30 logMAR or better pre-operative BCVA, of which seven (88%) had maintained

this post-operatively. However, among the twenty-three (84%) patients with pre-operative BCVA worse than 0.30 logMAR, eight (35%) achieved post-operative BCVA of 0.30 logMAR units or better. Despite having significantly larger logMAR gain ($p = 0.027$), eyes with worse pre-operative BCVA had worse post-operative BCVA ($p = 0.002$) (Figure 4B).




Figure 4. Effect of pre-operative visual acuity on visual outcomes. (A) Receiver Operating Characteristic (ROC) analysis using Area Under Curve (AUC) determines cut-off pre-operative BCVA of ≤ 0.48 logMAR as a predictor for achieving 0.30 logMAR at final follow-up (Sensitivity: 71.4%, Specificity of 88.2%, AUC 0.828 [95%CI 0.681 to 0.974], $p = 0.002$, Youden Index [J] of 0.596). (B) Error bar chart (95% confidence interval) of pre-operative and post-operative mean BCVA (logMAR) and logMAR gain grouped by pre-operative BCVA achieving 0.30 logMAR. Despite eyes in the worse pre-operative BCVA group having significantly larger logMAR gain, ($p = 0.027$) they also had worse post-op BCVA ($p = 0.002$).

4. Discussion

We evaluated eyes that underwent vitrectomy with ERM/ILM peeling for LMH to investigate potential biomarkers of visual outcome. Surgical indications in eyes with LMH remain a controversial topic, mainly due to the contentious outcomes described following surgery [13–15]. Indeed, improvement in BCVA and/or OCT-based foveal features (such as an increase in central retinal thickness or resolution of foveal defect) have been reported after surgery for LMH [9,11,16–21]; at the same time, caution has been advised by some when choosing surgery [14,17,22–24]. Several studies have previously investigated the possible pre-operative predictors for surgical outcomes, but no consensus was achieved [13,14,17,22,25–27]. Our choice to investigate OCT volumes is based on the concept that a volumetric analysis could better represent the retinal three-dimensional structural changes in LMH.

We identified pre-operative BCVA as the main clinical parameter for final BCVA, explaining alone 44.4% of the variability in post-operative BCVA ($r^2: 0.444$, adjusted $r^2: 0.425$). This is consistent with previously described findings reporting a positive correlation between pre-operative BCVA and post-operative BCVA following LMH repair [10,21,25,28]. In our study, BCVA significantly improved from 0.48 (0.22) before surgery to 0.36 (0.19) logMAR after surgery. Moreover, we identified a cut off of a pre-operative BCVA better than 0.48 logMAR as a predictor factor to achieve a post-operative BCVA of 0.30 logMAR or better (with a sensitivity of 71.4% and a specificity of 88.2%). However, we also showed that visual gain was particularly significant in eyes with pre-operative BCVA worse than 0.30 logMAR. These findings mean that, on one hand, patients with LMH have the best chance of keeping driving standard vision (0.30 logMAR) if surgery is carried out before the BCVA drops to 0.48 or worse; but on the other hand, early surgery might imply small

or no visual gain and patients should be informed that, in some cases, early surgery just stabilizes visual acuity.

Our study confirmed the negative association between the presence of ERP and post-surgical visual outcome ($p = 0.006$ and $p = 0.020$ for univariate and multivariate regression, respectively), as previously demonstrated in the scientific literature [13,14]. Interestingly, in the D-LMH group, increased FC volume ($p = 0.032$) and lower CRV ($p = 0.034$) showed a correlation with worse post-operative BCVA. In the ERM-FS group, lower CRV ($p < 0.001$) and ERP volume ($p < 0.001$), increased SV ($p < 0.001$) and higher foveal ONL volume ($p < 0.001$) were all significant moderators associated with worse visual outcomes after surgery in the multivariate regression. Caution should be advised to not interpret these as main effects, but as significant moderators, to avoid a Table 2 fallacy error [29]. In the scientific literature, only the study of Taşlıpınar Uzel et al. analyzed retinal volumes in the context of LMH, focusing on FC and ERP volumes in the natural history of D-LMH eyes that did not undergo surgery [6]. Similarly to previous studies [30–32], Taşlıpınar Uzel et al. confirmed no correlation between vision and linear OCT measurements such as CRT in D-LMH, with FC volume the only factor associated with baseline BCVA on multivariate regression [6]. CRV or ONL volume have never been investigated in the natural history, or correlated with surgical outcomes, in LMHs. To the best of our knowledge, this is the first report on pre-operative volumetric analysis of surgically treated LMHs. In the ERM-FS subtype, correlation between higher SV and worse visual outcome could be explained by the tractional forces of the ERM: larger SV might be present in more severe stages of the pathology with augmented tractional forces due to the ERM. Similarly, in D-LMH, we supposed that higher FC volumes reflect the increased degree of retinal tissue loss in the pathogenesis of the degenerative disease, indicating an advanced stage of the pathology and consequent worse visual outcomes after surgery. Moreover, smaller pre-operative CRV in our cohort (which correlates with larger SV in ERM-FS and higher FC in D-LMH) indicates a reduction in the viable retinal tissue and its volume (Figure 2), leading to poor functional outcomes post-surgery. We demonstrate that a higher foveal ONL volume is a negative moderator for post-operative BCVA in ERM-FS. We believe that this finding may appear consistent with what has been already demonstrated for idiopathic ERM [33,34]. In the view of the causative association between tractional ERM and the development of foveoschisis in ERM-FS [2], we suggest that ERM-FS may share some characteristics with eyes affected by ERM, including the association between increased ONL volume and impaired visual function. In summary, the increased schisis and higher ONL volumes point to the increased tractional effect of ERM on the retinal tissue, with a possible effect on photoreceptors, which may explain worse functional outcomes post-surgery. In the D-LMH group, lower CRV ($p = 0.034$) and higher FC volume ($p = 0.032$), both indicating a greater loss of retinal tissue, had a correlation with reduced final vision.

Vitreotomy combined with cataract surgery is a critical confounder in studies that report on visual outcomes following surgical intervention for LMH. In our cohort, we found that the combined phacovitrectomy group, relative to patients pseudophakic at baseline, gained an additional 0.03 logMAR, which was not significant. We report significant improvement in vision, whether the patient had cataract surgery, or was pseudophakic at baseline, without measured benefit to visual outcomes on the multivariate regression model by having cataract surgery. This is most likely due to our cohort of patients not having significant lenticular opacity at baseline and cataract surgery was primarily performed prophylactically with vitrectomy.

We acknowledge that this study has several limitations, including its retrospective nature, the limited number of patients and a relatively short follow-up time. Nonetheless, we conducted a precise analysis of the asymmetric retinal morphology in LMHs by using a standardized OCT protocol. Finally, this is the first study, to our knowledge, to perform a volumetric analysis on eyes that underwent PPV for LMH and we presented a well-fitting robust regression model.

5. Conclusions

In conclusion, our study highlights clinical and novel OCT volumetric biomarkers that correlate with post-operative BCVA following surgical intervention in LMHs. Although pre-operative BCVA was confirmed as a strong predictor of final visual outcome, the volumetric pre-operative independent variables included in our regression model offered a superior fit. We describe novel pre-operative OCT parameters significantly associated with visual outcomes following surgery such as CRV, ERP volume, SV and foveal ONL volume in ERM-FS, and CRV and FC volume in D-LMH. Further studies with larger numbers and longer post-operative follow-up can better investigate the role of volumetric analysis in the surgical treatment of lamellar macular holes.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/jpm14070755/s1>; Figure S1: effect of pre-operative visual acuity on visual outcomes; Table S1: correlation between clinical and OCT-based parameters and pre- and post-operative visual acuity; Table S2: correlation matrix of pre- and post-operative BCVA and OCT parameters.

Author Contributions: Conceptualization, M.L., A.J., M.F. and G.M.; methodology, M.L. and A.J.; software, G.M.; validation, N.A. and K.S.; formal analysis, G.M.; investigation, M.L., A.J. and G.M.; resources, T.I., K.J., F.D.-S., G.M., N.P. and A.J.; data curation, G.M.; writing—original draft preparation, M.L., M.F. and G.M.; writing—review and editing, M.F., A.J. and T.I.; visualization, K.S., N.A., K.J., F.D.-S., N.P. and M.F.; supervision, A.J., M.F. and T.I.; project administration, A.J. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Ethical review and approval were waived for this study, which was a retrospective anonymized data collection, and considered service evaluation as per our local protocol and hence had ethical approval exemption.

Informed Consent Statement: Patient consent was waived due to its retrospective nature.

Data Availability Statement: Data are contained within the article and Supplementary Materials.

Conflicts of Interest: The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Witkin, A.J.; Ko, T.H.; Fujimoto, J.G.; Schuman, J.S.; Bauman, C.R.; Rogers, A.H.; Reichel, E.; Duker, J.S. Redefining lamellar holes and the vitreomacular interface: An ultrahigh-resolution optical coherence tomography study. *Ophthalmology* **2006**, *113*, 388–397. [[CrossRef](#)] [[PubMed](#)]
2. Hubschman, J.P.; Govetto, A.; Spaide, R.F.; Schumann, R.; Steel, D.; Figueroa, M.S.; Sebag, J.; Gaudric, A.; Staurenghi, G.; Haritoglou, C.; et al. Optical coherence tomography-based consensus definition for lamellar macular hole. *Br. J. Ophthalmol.* **2020**, *104*, 1741–1747. [[CrossRef](#)] [[PubMed](#)]
3. Duker, J.S.; Kaiser, P.K.; Binder, S.; de Smet, M.D.; Gaudric, A.; Reichel, E.; Sadda, S.R.; Sebag, J.; Spaide, R.F.; Stalmans, P. The International Vitreomacular Traction Study Group classification of vitreomacular adhesion, traction, and macular hole. *Ophthalmology* **2013**, *120*, 2611–2619. [[CrossRef](#)] [[PubMed](#)]
4. Govetto, A.; Dacquay, Y.; Farajzadeh, M.; Platner, E.; Hirabayashi, K.; Hosseini, H.; Schwartz, S.D.; Hubschman, J.P. Lamellar Macular Hole: Two Distinct Clinical Entities? *Am. J. Ophthalmol.* **2016**, *164*, 99–109. [[CrossRef](#)] [[PubMed](#)]
5. Catania, F.; Romano, M.R.; Crincoli, E.; Allegrini, D.; Miere, A.; Chehaibou, I.; Abdelmassih, Y.; Beaumont, W.; Chapron, T.; Souied, E.H.; et al. Phenomenology of spontaneous closure in degenerative and mixed type lamellar macular hole. *Eye* **2024**, *38*, 315–320. [[CrossRef](#)] [[PubMed](#)]
6. Taşlıpınar Uzel, A.G.; Gelisken, F.; Kühlewein, L.; Neubauer, J. Volumetric analysis of lamellar macular hole: An Optical Coherence Tomography Study. *Retina* **2023**, *43*, 209–214. [[CrossRef](#)] [[PubMed](#)]
7. Figueroa, M.S.; Govetto, A.; Steel, D.H.; Sebag, J.; Virgili, G.; Hubschman, J.P. Pars plana vitrectomy for the treatment of tractional and degenerative lamellar macular holes: Functional and Anatomical Results. *Retina* **2019**, *39*, 2090–2098. [[CrossRef](#)] [[PubMed](#)]
8. Parisi, G.; Fallico, M.; Maugeri, A.; Barchitta, M.; Agodi, A.; Russo, A.; Longo, A.; Avitabile, T.; Castellino, N.; Bonfiglio, V.; et al. Primary vitrectomy for degenerative and tractional lamellar macular holes: A systematic review and meta-analysis. *PLoS ONE* **2021**, *16*, e0246667. [[CrossRef](#)]

9. Morescalchi, F.; Russo, A.; Gambicorti, E.; Cancarini, A.; Scaroni, N.; Bahja, H.; Costagliola, C.; Semeraro, F. Peeling of the internal limiting membrane with foveal sparing for treatment of degenerative lamellar macular hole. *Retina* **2020**, *40*, 1087–1093. [[CrossRef](#)]
10. Lee, C.S.; Koh, H.J.; Lim, H.T.; Lee, K.S.; Lee, S.C. Prognostic factors in vitrectomy for lamellar macular hole assessed by spectral-domain optical coherence tomography. *Acta Ophthalmol.* **2012**, *90*, e597–e602. [[CrossRef](#)]
11. Lai, T.T.; Chen, S.N.; Yang, C.M. Epiretinal proliferation in lamellar macular holes and full-thickness macular holes: Clinical and surgical findings. *Graefes Arch. Clin. Exp. Ophthalmol.* **2016**, *254*, 629–638. [[CrossRef](#)] [[PubMed](#)]
12. Lippera, M.; Ferrara, M.; Spiess, K.; Alnafisee, N.; Ally, N.; Jalil, A.; Ivanova, T.; Moussa, G. Novel OCT-Based Method to Measure Volumes of Retinal Specific Entities. *Preprints* **2024**, 2024061063. [[CrossRef](#)]
13. Ko, J.; Kim, G.A.; Lee, S.C.; Lee, J.; Koh, H.J.; Kim, S.S.; Byeon, S.H.; Lee, C.S. Surgical outcomes of lamellar macular holes with and without lamellar hole-associated epiretinal proliferation. *Acta Ophthalmol.* **2017**, *95*, e221–e226. [[CrossRef](#)]
14. Venkatesh, R.; Pereira, A.; Jain, K.; Kumar Yadav, N. Structural and Functional Outcomes of Surgery for Lamellar Macular Holes with or without Epimacular Proliferations. *J. Ophthalmic Vis. Res.* **2022**, *17*, 42–50. [[CrossRef](#)] [[PubMed](#)]
15. Murphy, D.C.; Rees, J.; Steel, D.H. Surgical interventions for lamellar macular holes. *Cochrane Database Syst. Rev.* **2021**, 2021, CD013678.
16. Pertile, G.; Iacovello, D.; Maraone, G.; Bottega, E.; Guerriero, M.; Maggio, E. Lamellar macular defects: Are degenerative lamellar macular holes truly degenerative? *Front. Med.* **2023**, *10*, 1156410. [[CrossRef](#)] [[PubMed](#)]
17. Coassin, M.; Mastrofilippo, V.; Stewart, J.M.; Fanti, A.; Belpoliti, M.; Cimino, L.; Iovieno, A.; Fontana, L. Lamellar macular holes: Surgical outcome of 106 patients with long-term follow-up. *Graefes Arch. Clin. Exp. Ophthalmol.* **2018**, *256*, 1265–1273. [[CrossRef](#)] [[PubMed](#)]
18. Frisina, R.; Tozzi, L.; Gius, I.; Pilotto, E.; Midena, E. Novel approaches to the assessment and treatment of lamellar macular hole. *Acta Ophthalmol.* **2022**, *100*, e1287–e1297. [[CrossRef](#)]
19. Haave, H.; Petrovski, B.É.; Zając, M.; Lumi, X.; Melekidou, W.; Lytvynchuk, L.; Ruban, A.; Znaor, L.; Nawrocki, J.; Nawrocka, Z.A.; et al. Outcomes from the Retrospective Multicenter Cross-Sectional Study on Lamellar Macular Hole Surgery. *Clin. Ophthalmol.* **2022**, *16*, 1847–1860. [[CrossRef](#)]
20. Kanai, M.; Sakimoto, S.; Takahashi, S.; Nishida, K.; Maruyama, K.; Sato, S.; Sakaguchi, H.; Nishida, K. Embedding Technique versus Conventional Internal Limiting Membrane Peeling for Lamellar Macular Holes with Epiretinal Proliferation. *Ophthalmol. Retina* **2023**, *7*, 44–51. [[CrossRef](#)]
21. Chehaibou, I.; Tadayoni, R.; Hubschman, J.P.; Bottoni, F.; Caputo, G.; Chang, S.; Dell’Omo, R.; Figueroa, M.S.; Gaudric, A.; Haritoglou, C.; et al. Natural History and Surgical Outcomes of Lamellar Macular Holes. *Ophthalmol. Retina* **2024**, *8*, 210–222. [[CrossRef](#)] [[PubMed](#)]
22. Parolini, B.; Schumann, R.G.; Cereda, M.G.; Haritoglou, C.; Pertile, G. Lamellar macular hole: A clinicopathologic correlation of surgically excised epiretinal membranes. *Invest. Ophthalmol. Vis. Sci.* **2011**, *52*, 9074–9083. [[CrossRef](#)] [[PubMed](#)]
23. Witkin, A.J.; Castro, L.C.; Reichel, E.; Rogers, A.H.; Baumal, C.R.; Duker, J.S. Anatomic and visual outcomes of vitrectomy for lamellar macular holes. *Ophthalmic Surg. Lasers Imaging* **2010**, *41*, 418–424. [[CrossRef](#)] [[PubMed](#)]
24. Khodabande, A.; Riazi-Esfahani, H.; Ghassemi, H.; Mirzaei, A.; Abbas, H.; Mirghorbani, M. Structural and functional outcomes of pars plana vitrectomy in patients with lamellar macular hole. *Taiwan. J. Ophthalmol.* **2022**, *12*, 430–436. [[PubMed](#)]
25. Omoto, T.; Asahina, Y.; Zhou, H.P.; Fujino, R.; Takao, M.; Obata, R.; Inoue, T.; Asaoka, R.; Maruyama-Inoue, M.; Yanagi, Y.; et al. Visual outcomes and prognostic factors of vitrectomy for lamellar macular holes and epiretinal membrane foveoschisis. *PLoS ONE* **2021**, *16*, e0247509. [[CrossRef](#)] [[PubMed](#)]
26. Su, Y.T.; Yang, C.M.; Lai, T.T. Multimodal Imaging Evidence of Traction Component in Lamellar Macular Hole with Epiretinal Proliferation. *Ophthalmic Res.* **2023**, *66*, 835–845. [[CrossRef](#)] [[PubMed](#)]
27. Abousy, M.; Drew-Bear, L.E.; Gibbons, A.; Pan-Doh, N.; Li, X.; Handa, J.T. In-Depth Analysis of Pre-Operative OCT Markers as Prognostic Factors for Lamellar Macular Holes and Epiretinal Membrane Foveoschisis. *Ophthalmol. Retina* **2024**, *8*, 465–472. [[CrossRef](#)] [[PubMed](#)]
28. Holland, L.; Chen, J.C.; Lee, L.R. Anatomical and Functional Outcomes of Pars Plana Vitrectomy for Lamellar Macular Defects. *Asia Pac. J. Ophthalmol.* **2015**, *4*, 134–139. [[CrossRef](#)] [[PubMed](#)]
29. Westreich, D.; Greenland, S. The Table 2 fallacy: Presenting and interpreting confounder and modifier coefficients. *Am. J. Epidemiol.* **2013**, *177*, 292–298. [[CrossRef](#)]
30. dell’Omo, R.; Virgili, G.; Rizzo, S.; De Turrís, S.; Coclite, G.; Giorgio, D.; Dell’Omo, E.; Costagliola, C. Role of Lamellar Hole-Associated Epiretinal Proliferation in Lamellar Macular Holes. *Am. J. Ophthalmol.* **2017**, *175*, 16–29. [[CrossRef](#)]
31. Compera, D.; Schumann, R.G.; Cereda, M.G.; Acquistapace, A.; Lita, V.; Priglinger, S.G.; Staurenghi, G.; Bottoni, F. Progression of lamellar hole-associated epiretinal proliferation and retinal changes during long-term follow-up. *Br. J. Ophthalmol.* **2018**, *102*, 84–90. [[CrossRef](#)] [[PubMed](#)]
32. Theodossiadis, P.G.; Grigoropoulos, V.G.; Emfietzoglou, I.; Nikolaidis, P.; Vergados, I.; Apostolopoulos, M.; Theodossiadis, G.P. Evolution of lamellar macular hole studied by optical coherence tomography. *Graefes Arch. Clin. Exp. Ophthalmol.* **2009**, *247*, 13–20. [[CrossRef](#)] [[PubMed](#)]

33. Colakoglu, A.; Balci Akar, S. Potential role of Müller cells in the pathogenesis of macropsia associated with epiretinal membrane: A hypothesis revisited. *Int. J. Ophthalmol.* **2017**, *10*, 1759–1767. [[PubMed](#)]
34. Kim, J.H.; Kim, Y.M.; Chung, E.J.; Lee, S.Y.; Koh, H.J. Structural and functional predictors of visual outcome of epiretinal membrane surgery. *Am. J. Ophthalmol.* **2012**, *153*, 103–110.e1. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.