

Pine Response to Sawfly Pheromones: Effects on Sawfly's Oviposition and Larval Growth

Asifur Rahman-Soad¹, Norbert Bittner^{1,2} and Monika Hilker^{1*}

¹Applied Zoology/Animal Ecology, Institute of Biology, Freie Universität Berlin, Berlin, Germany

²Institute of Translational Genomics, Helmholtz Zentrum München Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany

Asifur Rahman-Soad: ORCID: <https://orcid.org/0000-0001-7387-5391>

Norbert Bittner: ORCID: <https://orcid.org/0000-0002-8588-185X>

Monika Hilker: ORCID: <http://orcid.org/0000-0001-7519-7395>

*Corresponding author:

Monika Hilker, Applied Zoology/Animal Ecology, Institute of Biology, Freie Universität Berlin, Berlin, Germany, Email: monika.hilker@fu-berlin.de

Table S1. Comparison of pupation of *Diprion pini* larvae that fed upon either egg-free or egg-laden *Pinus sylvestris* trees exposed to different pre-treatments with volatiles. Pupation of larvae that fed on pine trees without egg deposition: untreated (U+L), hexane-treated trees (H+L), or pheromone-treated trees (P+L). Pupation of larvae that fed on trees with sawfly egg deposition (E): untreated (U+E+L), hexane-treated (H+E+L), or pheromone-treated (P+E+L) trees. Data: Mean (\pm SE) number of pupae per tree; each treatment consisted of 7-10 trees and maximum 5 larvae per tree. Statistical differences were determined using the Mann Whitney *U* test.

	Treatments							
	U+L	U+E+L		H+L	H+E+L		P+L	P+E+L
Mean	4.00	3.27		3.80	3.54		3.70	3.00
SE	0.235	0.406		0.290	0.340		0.366	0.330
p value	0.2111			0.6402			0.1015	