

Supplementary material

Supplementary Table S1. Percentage of burnout cases according to sociodemographic groups.

SEX				
	Female	Male	Total	P
Single-item 'inclusive'	29.6%	28.6%	29.3%	0.465
Single-item 'strict'	7.7%	3.4%	6.5%	0.072
BBQ factor 2	27.3%	34.5%	29.3%	0.092
AGE				
	Under 50	Over 50	Total	P
Single-item 'inclusive'	26.4%	34.2%	30.3%	0.024
Single-item 'strict'	4.1%	8.9%	6.5%	0.014
BBQ factor 2	26.7%	36.0%	31.3%	0.018
PROFESSIONAL GROUP: HEALTH/NON-HEALTH				
	Non-health	Health	Total	P
Single-item 'inclusive'	22.3%	33.7%	30.5%	0.005
Single-item 'strict'	3.4%	7.8%	6.5%	0.033
BBQ factor 2	21.7%	35.7%	31.5%	0.000
PROFESSIONAL GROUP: CONTACT WITH PUBLIC/NO CONTACT				
	No public contact	Public contact	Total	P
Single-item 'inclusive'	25.7%	31.9%	30.5%	0.101
Single-item 'strict'	2.2%	7.8%	6.5%	0.012
BBQ factor 2	25.0%	33.5%	31.5%	0.038
CARE LEVEL				
	Primary care	Hospital	Total	P
Single-item 'inclusive'	30.5%	30.1%	30.2%	0.519
Single-item 'strict'	8.5%	5.7%	6.1%	0.220
BBQ factor 2	40.2%	29.6%	31.4%	0.040