

**Table S1. Cytopathogenic effects**

<b>MP-12</b>								
	<i>MaKo</i>	<i>TH-1</i>	<i>IgH-2</i>	<i>A6</i>	<i>SKM</i>	<i>VH-2</i>	<i>Vero76</i>	<i>CDSK</i>
<i>6hpi</i>	-	-	-	-	-	-	-	-
<i>12hpi</i>	-	-	-	-	-	-	-	-
<i>24hpi</i>	-	-	-	-	-	-	-	-
<i>48hpi</i>	-	-	+	+	-	-	+	-
<i>72hpi</i>	+	+	+	+	-	+	+	-
<b>Clone 13</b>								
	<i>MaKo</i>	<i>TH-1</i>	<i>IgH-2</i>	<i>A6</i>	<i>SKM</i>	<i>VH-2</i>	<i>BHK-21</i>	<i>CDSK</i>
<i>6hpi</i>	-	-	-	-	-	-	-	-
<i>12hpi</i>	-	-	-	-	-	-	-	-
<i>24hpi</i>	-	-	-	-	-	-	-	-
<i>48hpi</i>	-	-	-	-	-	-	-	-
<i>72hpi</i>	-	+	-	-	-	-	+	-
<b>ZH501</b>								
	<i>MaKo</i>	<i>TH-1</i>	<i>IgH-2</i>	<i>A6</i>	<i>SKM</i>	<i>VH-2</i>	<i>VeroE6</i>	<i>CDSK</i>
<i>6hpi</i>	-	-	-	-	-	-	-	-
<i>12hpi</i>	-	-	-	-	-	-	-	-
<i>24hpi</i>	-	-	-	-	-	-	-	-
<i>48hpi</i>	-	+	+	-	-	-	+	+
<i>72hpi</i>	-	+	+	-	-	+	+	+

Cytopathogenic effects were evaluated at the according time points after infection before petri dishes were frozen. Infected cells were compared to corresponding mock-infected cells, that were treated and kept in the same way like infected cells.