

Hypermucoviscous Multidrug-Resistant *Klebsiella variicola* Strain LL2208 Isolated from Chinese Longsnout Catfish (*Leiocassis longirostris*): Highly Similar to Human *K. variicola* Strains

Qingyong Li ¹, Xin Yu ¹, Lin Ye ¹, Tongyu Hou ¹, Yi Liu ¹, Guiming Liu ¹, Qing Wang ^{2,3} and Defeng Zhang ^{2,3,*}

¹ Fisheries Research and Extension Center of Huizhou, Huizhou 516055, China; byqyli@163.com (Q.L.); 27763967@126.com (X.Y.); 13528069240@139.com (L.Y.); houtongyu@hotmail.com (T.H.); 3825431221@139.com (Y.L.); 13928333187@189.com (G.L.)

² Key Laboratory of Fishery Drug Development, Ministry of Agriculture and Rural Affairs, Pearl River Fisheries Research Institute, Chinese Academy of Fishery Sciences, Guangzhou 510380, China; wangqing@prfri.ac.cn

³ Guangdong Provincial Key Laboratory of Aquatic Animal Immunology and Sustainable Aquaculture, Pearl River Fisheries Research Institute, Chinese Academy of Fishery Sciences, Guangzhou 510380, China

* Correspondence: zhangdefeng@prfri.ac.cn

Table S1. Gene islands in the genome of strain LL2208 predicted by IslandViewer 4.

Gene Islands	Positions in Genome
GI_1	1,191,331 bp – 1,203,694 bp
GI_2	1,943,658 bp – 1,967,036 bp
GI_3	3,016,227 bp – 3,040,252 bp
GI_4	3,172,831 bp – 3,182,967 bp
GI_5	4,185,931 bp – 4,215,920 bp
GI_6	4,242,977 bp – 4,254,623 bp




Figure S1. Virulence associated genes were detected by PCR.
Lane M, DNA markers (DL2000); Lane 1, *rmpA*; Lane 2, *rmpA2*; Lane 3, *magA*; Lane 4, *peg-344*;
Lane 5, *iroB*; Lane 6, *iucB*.