

Survival of Non-psychrophilic Methanogens Exposed to Extreme Temperature Changes.

Rebecca L. Mickol^{1,2}, Sarah K. Laird³, and Timothy A. Kral^{1,3}

¹Arkansas Center for Space and Planetary Sciences, University of Arkansas, Fayetteville, AR;

²American Society for Engineering Education, Washington, DC; ³Dept. of Biological Sciences,

University of Arkansas, Fayetteville, AR; [rebecca.mickol@gmail.com]

SUPPLEMENTARY FIGURES




Figure S1. Inoculation scheme for Transfer Sets 1, 2, and 3 for cultures of *Methanothermobacter wolfeii* in Experiment 2 (5 g sand, 10 mL medium).




Figure S2. Inoculation scheme for Transfer Sets 1, 2, and 3 for cultures of *Methanobacterium formicicum* in Experiment 2 (5 g sand, 10 mL medium).




Figure S3. Inoculation scheme for Transfer Sets 1, 2, and 3 for cultures of *Methanothermobacter wolfeii* in Experiment 3 (10 g sand, 5 mL medium).




Figure S4. Inoculation scheme for Transfer Sets 1, 2, and 3 for cultures of *Methanobacterium formicicum* in Experiment 3 (10 g sand, 5 mL medium).




Figure S5. Inoculation scheme for Transfer Sets 1, 2, and 3 for cultures of *Methanobacterium formicum* and *Methanothermobacter wolfeii* in Experiment 4 (5 mL medium).