



Supplementary Material

Taxogenomic and Metabolic Insights *Marinobacterium ramblicola* sp. nov., a New Slightly Halophilic Bacterium Isolated from Rambla Salada, Murcia

Ana Durán-Viseras ^{1,2*}, David J. Castro^{1,2}, José Carlos Reina^{1,2}, Victoria Béjar^{1,2} and Fernando Martínez-Checa^{1,2*}

¹ Microbial Exopolysaccharide Research Group, Department of Microbiology, Pharmacy Faculty, Campus de Cartuja s.n., 18071 Granada, Spain; anaduran@us.es (A.D.-V.), castricoxpc@gmail.com (D.J.C.), josecreina@ugr.es (J.C.R.), vbejar@ugr.es (V.B.), fmcheca@ugr.es (F.M.-C.)

² Biomedical Research Center, Institute of Biotechnology, 18016 Granada, Spain

* Correspondence: anaduran@us.es fmcheca@ugr.es




Figure S1. Two-dimensional TLC of polar lipids from strain D7^T. **Abbreviations:** PE, phosphatidylethanolamine; PG, phosphatidylglycerol; L, lipid.




Figure S2. Electron micrographs of negatively stained preparation of cells of strain D7^T. Cells were grown on MA plates at 32 °C for 24 h with shaking at 150 rpm and stained with % (w/v) uranyl acetate (Bar, 1 µm).



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).