

Supplementary File 3

Comparison of lesion distribution between **horses doing different sports activities** are shown in **Table 3a**, **Table 3b** and **Table 3c**.

MRI features from horses undergoing standing magnetic resonance imaging of the metacarpophalangeal/metatarsophalangeal region from horses with a history of undertaking different sports with number (N) and proportion (%) of forelimbs and hindlimbs affected. Features with significant differences in proportions between different sports (Pearson's Chi-squared test) are shown with overrepresented category in bold and underrepresented underlined, using a significant level of $p < 0.05$.

MCP j=metacarpophalangeal joint; MTP j=metatarsophalangeal joint; MC 3=third metacarpal bone; MT 3=third metatarsal bone; P1=proximal phalanx; PSB= proximal sesamoid bones; POD=palmar/plantar osteochondral disease; DFTS=digital flexor tendon sheath; Med SB=medial suspensory branch; Lat SB=lateral suspensory branch; MCL=medial collateral ligament of the metacarpophalangeal/metatarsophalangeal joint; LCL=lateral collateral ligament of the metacarpophalangeal/metatarsophalangeal joint; SDSL=straight distal sesamoidean ligament; MODSL=medial oblique distal sesamoidean ligament; LODSL=lateral oblique distal sesamoidean ligament; DDFT=deep digital flexor tendon; SDFT=superficial digital flexor tendon.

Table 3a: Distribution and pathology type of **bone lesions** of the metacarpophalangeal/metatarsophalangeal region from horses with a history of undertaking different sports with number (N) and proportion (%) of forelimbs and hindlimbs affected.

Table 3b: Distribution and pathology type of **metacarpophalangeal/metatarsophalangeal joint** and **digital flexor tendon sheath** lesions from horses with a history of undertaking different sports with number (N) and proportion (%) of forelimbs and hindlimbs affected.

Table 3c: Distribution and pathology type affecting the **soft tissues** of the metacarpophalangeal/metatarsophalangeal region from horses with a history of undertaking different sports with number (N) and proportion (%) of forelimbs and hindlimbs affected.

Table 3a Anatomical location Bone			MRI features	Dressage N=97	Eventing N=27	General purpose N=27	Racing N=88	Show jumping N=47	Endurance N=55	
				N and (%) of limbs	N and (%) of limbs	N and (%) of limbs	N and (%) of limbs	N and (%) of limbs	N and (%) of limbs	P value
MC/T 3	Cortex	Location	Dorsal	14 (14.4)	8 (29.6)	5 (18.5)	25 (28.4)	6 (12.8)	4 (7.3)	0.011
			Palmar/Plantar	<u>6 (6.2)</u>	5 (18.5)	4 (14.8)	28 (31.8)	4 (8.5)	4 (7.3)	<.001
			Axial (sagittal ridge)	7 (7.2)	5 (18.5)	3 (11.1)	15 (17)	5 (10.6)	2 (3.6)	
			Medial	11 (11.3)	<u>8 (29.6)</u>	4 (14.8)	29 (33)	3 (6.4)	3 (5.5)	<.001
			Lateral	<u>2 (2.1)</u>	4 (14.8)	3 (11.1)	26 (29.6)	1 (2.1)	3 (5.5)	<.001
		Aspect	Periosteal	12 (12.4)	4 (14.8)	5 (18.5)	29 (33)	6 (12.8)	<u>1 (1.8)</u>	<.001
			Endosteal	<u>12 (12.4)</u>	7 (25.9)	7 (25.9)	35 (39.8)	7 (14.9)	6 (10.9)	<.001
			All	13 (13.4)	5 (18.5)	5 (18.5)	29 (33)	6 (12.8)	<u>1 (1.8)</u>	<.001
		Pathology type	Irregularity	4 (4.1)	2 (7.4)	2 (7.4)	13 (14.8)	4 (8.5)	5 (9.1)	
			Sclerosis	10 (10.3)	6 (22.2)	5 (18.5)	25 (28.4)	4 (8.5)	5 (9.1)	0.004

			Fluid (hyperintensity)	9 (9.3)	3 (11.1)	2 (7.4)	27 (30.7)	4 (8.5)	5 (9.1)	<.001
	Trabecular	Location	Dorsal	49 (50.5)	10 (37)	10 (37)	44 (50)	21 (44.7)	25 (45.5)	
	bone		Palmar/Plantar	19 (19.6)	7 (25.9)	6 (22.2)	60 (68.2)	8 (17)	40 (72.7)	<.001
			Axial (sagittal ridge)	24 (24.7)	4 (14.8)	3 (11.1)	25 (28.4)	10 (21.3)	14 (25.5)	
			Medial	52 (53.6)	11 (40.7)	9 (33.3)	57 (64.8)	22 (46.8)	30 (54.6)	0.042
			Lateral	17 (17.5)	7 (25.9)	5 (18.5)	53 (60.2)	6 (12.8)	26 (47.3)	<.001
		Pathology type	Irregularity	13 (13.4)	2 (7.4)	2 (7.4)	10 (11.4)	5 (10.6)	14 (25.5)	
			Sclerosis	45 (46.4)	10 (37)	10 (37)	54 (61.4)	18 (38.3)	39 (70.9)	<.001
			Fluid (hyperintensity)	45 (46.4)	5 (18.5)	6 (22.2)	52 (59.1)	18 (38.3)	42 (76.4)	<.001
			Fracture incomplete	-	-	1 (3.7)	2 (2.3)	-	10 (18.2)	<.001
			Hypervascularization	16 (16.5)	6 (22.2)	1 (3.7)	59 (67.1)	6 (12.8)	26 (47.3)	<.001
P1	Cortex	Location	Dorsal	24 (24.7)	5 (18.5)	4 (14.8)	25 (28.4)	10 (21.3)	24 (43.6)	0.037
			Palmar/Plantar	1 (1)	1 (3.7)	4 (14.8)	5 (5.7)	3 (6.4)	-	0.017
			Axial (sagittal groove)	15 (15.5)	3 (11.1)	2 (7.4)	15 (17.1)	7 (14.9)	17 (30.9)	
			Medial	15 (15.5)	3 (11.1)	4 (14.8)	15 (17.1)	7 (14.9)	4 (7.3)	
			Lateral	2 (2.1)	-	5 (18.5)	7 (7.9)	2 (4.3)	4 (7.3)	0.022
		Aspect	Periosteal	19 (19.6)	4 (14.8)	7 (25.9)	22 (25)	6 (12.8)	5 (9.1)	
			Endosteal	27 (27.8)	5 (18.5)	7 (25.9)	28 (31.8)	10 (21.3)	27 (49.1)	0.02
			All	19 (19.6)	4 (14.8)	6 (22.2)	21 (23.9)	6 (12.8)	4 (7.3)	
		Pathology type	Irregularity	24 (24.7)	5 (18.5)	7 (25.9)	17 (19.3)	6 (12.8)	26 (47.3)	0.001
			Sclerosis	12 (12.4)	4 (14.8)	4 (14.8)	15 (17.1)	4 (8.5)	4 (7.3)	
			Fluid (hyperintensity)	15 (15.5)	2 (7.4)	5 (18.5)	19 (21.6)	5 (10.6)	11 (20)	
	Trabecular bone	Location	Dorsal	27 (27.8)	3 (11.1)	3 (11.1)	26 (29.6)	12 (25.5)	13 (23.6)	
			Palmar/Plantar	4 (4.1)	1 (3.7)	4 (14.8)	4 (4.6)	3 (6.4)	2 (3.6)	
			Axial (sagittal groove)	10 (10.3)	1 (3.7)	2 (7.4)	12 (13.6)	8 (17)	11 (20)	
			Medial	21 (21.7)	3 (11.1)	2 (7.4)	14 (15.9)	10 (21.3)	4 (7.3)	
			Lateral	4 (4.1)	-	4 (14.8)	8 (9.1)	4 (8.5)	4 (7.3)	
		Pathology type	Irregularity	16 (16.5)	7 (25.9)	2 (7.4)	5 (5.7)	11 (23.5)	4 (7.3)	
			Sclerosis	14 (14.4)	5 (18.5)	3 (11.1)	4 (4.6)	16 (34.1)	5 (9.1)	
			Fluid (hyperintensity)	27 (27.8)	11 (40.7)	2 (7.4)	6 (6.8)	23 (49.1)	10 (18.2)	
PSB	Medial	Location	Axial	12 (12.4)	4 (14.8)	2 (7.4)	26 (29.6)	6 (12.8)	6 (10.9)	0.008
			Abaxial	28 (28.9)	3 (11.1)	2 (7.4)	31 (35.2)	13 (27.7)	21 (38.2)	0.014
			Proximal	15 (15.5)	4 (14.8)	2 (7.4)	26 (29.6)	6 (12.8)	6 (10.9)	0.008
			Distal	16 (16.5)	1 (3.7)	3 (11.1)	24 (27.3)	7 (14.9)	5 (9.1)	0.019
			Diffuse	9 (9.3)	2 (7.4)	3 (11.1)	30 (34.1)	3 (6.4)	8 (14.6)	<.001
		Pathology type	Irregularity	23 (23.7)	5 (18.5)	-	9 (10.2)	8 (17)	16 (29.1)	0.006
			Sclerosis	6 (6.2)	1 (3.7)	2 (7.4)	23 (26.1)	3 (6.4)	3 (5.5)	<.001

			Fluid (hyperintensity)	17 (17.5)	1 (3.7)	2 (7.4)	24 (27.3)	7 (14.9)	17 (30.9)	0.009
	Lateral	Location	Axial	9 (9.3)	3 (11.1)	2 (7.4)	20 (22.7)	7 (14.9)	7 (12.7)	
			Abaxial	24 (24.7)	1 (3.7)	2 (7.4)	24 (27.3)	8 (17)	18 (32.7)	0.013
			Proximal	15 (15.5)	1 (3.7)	2 (7.4)	20 (22.7)	6 (12.8)	5 (9.1)	
		Pathology type	Distal	9 (9.3)	-	3 (11.1)	20 (22.7)	7 (14.9)	5 (9.1)	0.018
			Diffuse	7 (7.2)	1 (3.7)	2 (7.4)	25 (28.4)	4 (8.5)	6 (10.9)	<.001
			Irregularity	18 (18.6)	3 (11.1)	-	4 (4.6)	3 (6.4)	14 (25.5)	<.001
			Sclerosis	7 (7.2)	1 (3.7)	1 (3.7)	15 (17.1)	-	4 (7.3)	0.011
			Fluid (hyperintensity)	19 (19.6)	-	2 (7.4)	23 (26.1)	7 (14.9)	16 (29.1)	

Table 3b Anatomical location Joint		MRI features	N and (%) of limbs DRE (N=97)	N and (%) of limbs EVE (N=27)	N and (%) of limbs GP (N=27)	N and (%) of limbs RAC (N=88)	N and (%) of limbs SJ (N=47)	N and (%) of limbs END (N=55)	P value
MC/MTP	Location	Dorsal P1	82 (84.5)	17 (63)	15 (55.6)	56 (63.6)	40 (85.1)	42 (76.4)	0.001
		Dorsal MC/T 3	85 (87.6)	19 (70.4)	14 (51.9)	54 (61.5)	35 (74.5)	42 (76.4)	
		Palmar/Plantar P1	16 (16.5)	5 (18.5)	7 (25.9)	11 (12.5)	17 (36.2)	4 (7.3)	0.003
		Palmar/Plantar MC/T 3	21 (21.7)	12 (44.4)	5 (18.5)	63 (71.6)	16 (34)	43 (78.2)	<.001
		Proximal P1	87 (89.7)	19 (70.4)	18 (66.7)	55 (62.5)	38 (80.9)	45 (81.8)	
		Distal MC/T 3	87 (89.7)	22 (81.5)	15 (59.3)	79 (89.8)	41 (87.2)	52 (94.6)	
		Medial P1	50 (51.6)	14 (51.9)	9 (33.3)	31 (35.2)	27 (57.5)	15 (27.3)	0.006
		Medial MC/T 3	67 (69.1)	14 (51.9)	12 (44.4)	61 (69.3)	33 (70.2)	33 (60)	
		Lateral P1	23 (23.7)	3 (11.1)	5 (18.5)	14 (15.9)	15 (31.9)	12 (21.8)	
		Lateral MC/T3	21 (21.7)	10 (37)	7 (25.9)	48 (54.6)	17 (36.2)	25 (45.5)	<.001
		Axial P1 (sagittal groove)	58 (59.8)	11 (40.7)	7 (25.9)	23 (26.1)	28 (59.6)	26 (47.3)	<.001
		Medial Parasagittal groove	13 (13.4)	2 (7.4)	1 (3.7)	32 (36.4)	5 (10.6)	17 (30.9)	<.001
		Lateral Parasagittal groove	7 (7.2)	0	1 (3.7)	32 (36.4)	4 (8.5)	25 (45.5)	<.001
		Axial MC/T 3 (sagittal ridge)	64 (66)	14 (51.9)	4 (14.8)	36 (40.9)	25 (53.2)	38 (69.1)	<.001
	Pathology type	Subchondral bone Irregularity	51 (52.6)	14 (51.9)	9 (33.3)	43 (48.9)	27 (57.5)	43 (78.2)	0.002
		Subchondral bone defect	19 (19.6)	3 (11.1)	1 (3.7)	35 (39.8)	7 (14.9)	8 (14.6)	<.001
		Sclerosis	74 (76.3)	19 (70.4)	14 (51.9)	66 (75)	35 (74.5)	50 (90.9)	
		Fluid (hyperintensity)	82 (84.5)	14 (51.9)	13 (48.2)	77 (87.5)	34 (72.3)	51 (92.7)	

		Cartilage irregularity/fibri llation	7 (7.2)	2 (7.4)	1 (3.7)	4 (4.6)	8 (17)	4 (7.3)	
		Fissure	<u>5 (5.2)</u>	1 (3.7)	-	38 (43.2)	-	22 (40)	<.001
		Osteophyte	81 (83.5)	11 (40.7)	14 (51.9)	<u>18 (20.5)</u>	35 (74.5)	44 (80)	<.001
		POD	1 (1)	-	-	15 (17.1)	-	-	<.001
		Distension	84 (86.6)	16 (59.3)	16 (59.3)	<u>36 (40.9)</u>	40 (85.1)	53 (96.4)	<.001
		Chronic synovitis	75 (77.3)	18 (66.7)	16 (59.3)	<u>28 (31.8)</u>	33 (70.2)	49 (89.1)	<.001
		Capsulitis	17 (17.5)	11 (40.7)	6 (22.2)	21 (23.9)	17 (36.2)	3 (5.5)	<.001
DFTS	<i>Pathology</i>	Distension	12 (12.4)	3 (11.1)	5 (18.5)	2 (2.3)	3 (6.4)	1 (1.8)	0.013
	<i>Type</i>	Chronic synovitis	10 (10.3)	3 (11.1)	3 (11.1)	<u>1 (1.1)</u>	2 (4.3)	1 (1.8)	0.041

Table 3c Anatomical location Soft tissue		MRI features	N and (%) of limbs DRE (N=97)	N and (%) of limbs EVE (N=27)	N and (%) of limbs GP (N=27)	N and (%) of limbs RAC (N=88)	N and (%) of limbs SJ (N=47)	N and (%) of limbs END (N=55)	P value
Med SB	<i>Location</i>	Insertion	19 (19.6)	4 (14.8)	3 (11.1)	<u>5 (5.7)</u>	14 (29.8)	17 (30.9)	<.001
		Body	45 (46.4)	7 (25.9)	11 (40.7)	<u>18 (20.5)</u>	21 (44.7)	34 (61.8)	<.001
	<i>Part of the Branch affected</i>	Margin/peripher y	40 (41.2)	6 (22.2)	5 (18.5)	14 (15.9)	19 (40.4)	29 (52.7)	<.001
		Central/core	33 (34)	5 (18.5)	3 (11.1)	<u>13 (14.7)</u>	16 (34)	21 (38.2)	0.003
		Entire branch	6 (6.2)	3 (11.1)	4 (14.8)	6 (6.8)	4 (8.5)	-	
	<i>Area of the Margin affected</i>	Dorsal	11 (11.3)	-	3 (11.1)	5 (5.7)	4 (8.5)	1 (1.8)	
		Palmar/plantar	15 (15.5)	4 (14.8)	5 (18.5)	7 (8)	8 (17)	3 (5.5)	
		Axial	38 (39.2)	5 (18.5)	7 (25.9)	<u>15 (17.1)</u>	15 (31.9)	28 (50.9)	<.001
		Abaxial	13 (13.4)	1 (3.7)	4 (14.8)	6 (6.8)	3 (6.4)	2 (3.6)	
	<i>Pathology type</i>	Defect	22 (22.7)	3 (11.1)	3 (11.1)	<u>6 (6.8)</u>	4 (8.5)	29 (52.7)	<.001
		Fluid-based hyperintensity	40 (41.2)	10 (37)	9 (33.3)	<u>13 (14.8)</u>	19 (40.4)	28 (50.9)	<.001
		Enlargement	21 (21.7)	4 (14.8)	7 (25.9)	<u>12 (13.6)</u>	18 (38.3)	11 (20)	0.032
Lat SB	<i>Location</i>	Insertion	23 (23.7)	4 (14.8)	2 (7.4)	<u>3 (3.4)</u>	8 (17)	10 (18.2)	0.004
		Body	32 (33)	11 (40.7)	6 (22.2)	<u>10 (11.4)</u>	16 (34)	16 (29.1)	0.005
	<i>Part of the branch affected</i>	Margin/peripher y	30 (30.9)	5 (18.5)	4 (14.8)	<u>8 (9.1)</u>	10 (21.3)	16 (29.1)	0.007
		Central/core	26 (26.8)	8 (29.6)	3 (11.1)	<u>9 (10.2)</u>	14 (29.8)	10 (18.2)	0.019
		Entire branch	12 (12.4)	4 (14.8)	3 (11.1)	<u>2 (2.3)</u>	8 (17)	2 (3.6)	0.024
	<i>Area of the Margin affected</i>	Dorsal	13 (13.4)	3 (11.1)	3 (11.1)	1 (1.1)	4 (8.5)	4 (7.3)	
		Palmar/plantar	15 (15.5)	5 (18.5)	2 (7.4)	4 (4.6)	7 (14.9)	6 (10.9)	
		Axial	31 (32)	6 (22.2)	5 (18.5)	<u>7 (8)</u>	10 (21.3)	15 (27.3)	0.005
		Abaxial	14 (14.4)	4 (14.8)	2 (7.4)	2 (2.3)	5 (10.6)	4 (7.3)	
	<i>Pathology type</i>	Defect	18 (18.6)	2 (7.4)	-	<u>6 (6.8)</u>	1 (2.1)	15 (27.3)	
		Fluid-based hyperintensity	33 (34)	11 (40.7)	7 (25.9)	<u>10 (11.4)</u>	18 (38.3)	13 (23.6)	0.002
		Enlargement	20 (20.1)	4 (14.8)	3 (11.1)	<u>2 (2.3)</u>	11 (23.4)	8 (14.6)	0.004
MCL	<i>Location</i>	Dorsal	4 (4.1)	-	1 (3.7)	1 (1.1)	2 (4.3)	-	

		Palmar/plantar	4(4.1)	-	1(3.7)	1(1.1)	1(2.1)	-	
		Axial	5(5.2)	-	1(3.7)	1(1.1)	1(2.1)	-	
		Abaxial	4(4.1)	-	1(3.7)	1(1.1)	2(4.3)	-	
		Entire ligament	4(4.1)	1(3.7)	1(3.7)	1(1.1)	1(2.1)	-	
		Superficial (long)	4(4.1)	1(3.7)	1(3.7)	-	2(4.3)	-	
		Deep (short)	5(5.2)	1(3.7)	1(3.7)	1(1.1)	1(2.1)	-	
		Origin	4(4.1)	1(3.7)	-	1(1.1)	1(2.1)	-	
		Body	5(5.2)	1(3.7)	1(3.7)	-	1(2.1)	-	
		Insertion	5(5.2)	1(3.7)	-	-	-	-	
		Entire ligament	2(2.1)	-	-	-	-	-	
	<i>Pathology type</i>	Fluid-based hyperintensity	5(5.2)	1(3.7)	-	-	2(4.3)	-	
		Enlargement	1(1)	1(3.7)	1(3.7)	-	1(2.1)	-	
LCL	<i>Location</i>	Dorsal	9(9.3)	1(3.7)	8(29.6)	4(4.6)	3(6.4)	1(1.8)	<.001
		Palmar/plantar	9(9.3)	-	7(25.9)	4(4.6)	2(4.3)	2(1.8)	<.001
		Axial	10(10.3)	2(7.4)	8(29.6)	4(4.6)	5(10.6)	1(1.8)	<.001
		Abaxial	9(9.3)	-	7(25.9)	3(3.4)	2(4.3)	1(1.8)	<.001
		Entire ligament	8(8.3)	-	7(25.9)	2(2.3)	2(4.3)	1(1.8)	<.001
	<i>Level</i>	Superficial (long)	9(9.3)	1(3.7)	8(29.6)	4(4.6)	4(8.5)	1(1.8)	<.001
		Deep (short)	9(9.3)	2(7.4)	8(29.6)	4(4.6)	5(10.6)	1(1.8)	<.001
		Origin	7(7.2)	1(3.7)	6(22.2)	4(4.6)	3(6.4)	1(1.8)	0.015
		Body	5(5.2)	1(3.7)	8(29.6)	2(2.3)	3(6.4)	1(1.8)	<.001
		Insertion	6(6.2)	-	4(14.8)	2(2.3)	1(2.1)	1(1.8)	0.034
		Entire ligament	4(4.1)	-	3(11.1)	1(1.1)	1(2.1)	1(1.8)	
	<i>Pathology type</i>	Fluid-based hyperintensity	7(7.2)	2(7.4)	9(33.3)	4(4.6)	5(10.6)	1(1.8)	<.001
		Enlargement	8(8.3)	1(3.7)	8(29.6)	4(4.6)	3(6.4)	1(1.8)	<.001
SDSL	<i>Location</i>	Dorsal	8(8.3)	-	-	-	-	1(1.8)	0.004
		Palmar/plantar	7(7.2)	-	-	-	-	1(1.8)	0.012
		Medial	9(9.3)	-	1(3.7)	-	-	1(1.8)	0.004
		Lateral	5(5.2)	1(3.7)	-	-	-	1(1.8)	
		Entire ligament	6(6.2)	-	-	-	-	-	0.009
	<i>Level</i>	Bone ligament interface	3(3.1)	-	-	-	-	-	
		Body	11(11.3)	1(3.7)	-	-	-	1(1.8)	<.001
	<i>Pathology type</i>	Irregularity	7(7.2)	-	-	-	-	1(1.8)	0.012
		Enlargement	4(4.1)	-	-	-	-	1(1.8)	
		Fluid-based hyperintensity	7(7.2)	-	-	-	-	-	0.003
		Adhesions	3(3.1)	1(3.7)	1(3.7)	-	-	-	
Med ODSL	<i>Location</i>	Dorsal	14(14.4)	2(7.4)	1(3.7)	2(2.3)	4(8.5)	3(5.5)	
		Palmar/plantar	15(15.5)	3(11.1)	3(11.1)	<u>2(2.3)</u>	4(8.5)	2(3.6)	0.028
		Axial	25(25.8)	1(3.7)	3(11.1)	<u>2(2.3)</u>	13(27.7)	5(9.1)	<.001
		Abaxial	16(16.5)	2(7.4)	1(3.7)	<u>2(2.3)</u>	2(4.3)	2(3.6)	0.004
		Entire ligament	14(14.4)	5(18.5)	1(3.7)	3(3.4)	2(4.3)	-	0.001

	<i>Level</i>	Bone ligament interface	4(4.1)	1(3.7)	1(3.7)	1(1.1)	-	-	
		Body	27 (27.8)	5(18.5)	4(14.8)	<u>3 (3.4)</u>	12(25.5)	5(9.1)	<.001
	<i>Pathology type</i>	Irregularity	16 (16.5)	-	3(11.1)	<u>2</u>	10(21.3)	4(7.3)	<.001
		Enlargement	17 (17.5)	3(11.1)	2(7.4)	<u>1 (1.1)</u>	3(6.4)	1(1.8)	<.001
		Fluid-based hyperintensity	27 (27.8)	5(18.5)	3(11.1)	<u>2 (2.3)</u>	7(14.9)	6(10.9)	<.001
		Adhesions	3(3.1)	-	1(3.7)	-	-	-	
Lat ODSL	<i>Location</i>	Dorsal	15 (15.5)	3(11.1)	5(18.5)	<u>2</u>	3(6.4)	1(1.8)	<.001
		Palmar/plantar	17 (17.5)	3(11.1)	2(7.4)	<u>2</u>	4(8.5)	2(3.6)	<.001
		Axial	21 (21.7)	3(11.1)	6(22.2)	<u>2</u>	7(14.9)	3(5.5)	<.001
		Abaxial	16 (16.5)	2(7.4)	3(11.1)	<u>2</u>	3(6.4)	1(1.8)	<.001
		Entire ligament	15 (15.5)	2(7.4)	4(14.8)	<u>2</u>	2(4.3)	-	<.001
	<i>Level</i>	Bone ligament interface	2(2.1)	-	2(7.4)	-	-	-	0.031
		Body	20 (20.6)	3(11.1)	6(22.2)	<u>2</u>	8(17)	4(7.3)	<.001
	<i>Pathology type</i>	Irregularity	12 (12.4)	-	5(18.5)	<u>2</u>	4(8.5)	3(5.5)	0.002
		Enlargement	14 (14.3)	3(11.1)	5(18.5)	<u>2</u>	4(8.5)	4(7.3)	0.007
		Fluid-based hyperintensity	20 (20.6)	3(11.1)	4(14.8)	<u>2</u>	6(12.8)	4(7.3)	<.001
		Adhesions	-	-	2(7.4)	-	1(2.1)	-	
DDFT	<i>Location</i>	Dorsal	-	1(3.7)	-	-	2(4.3)	1(1.8)	
		Palmar/plantar	-	1(3.7)	1(3.7)	-	-	-	
		Medial	-	-	-	-	1(2.1)	-	
		Lateral	-	1(3.7)	2(7.4)	-	-	1(1.8)	0.019
	<i>Level</i>	Proximal to the PSB	-	1(3.7)	2(7.4)	-	1(2.1)	1(1.8)	0.019
		At the level of the PSB	-	1(3.7)	1(3.7)	-	1(2.1)	-	
		Distal to the PSB	-	2(7.4)	1(3.7)	-	-	-	0.003
		Body	-	-	1(3.7)	-	2(4.3)	-	
	<i>Pathology type</i>	Irregularity	-	2(7.4)	2(7.4)	-	1(2.1)	1(1.8)	0.021
		Enlargement	-	1(3.7)	-	-	-	-	0.04
		Fluid-based hyperintensity	-	1(3.7)	-	-	-	1(1.8)	
		Adhesions	-	2(7.4)	1(3.7)	-	-	1(1.8)	0.019
		Chronic fibrosis	-	-	3(11.1)	-	1(2.1)	-	<.001
SDFT	<i>Location</i>	Dorsal	2(2.1)	-	1(3.7)	2(2.3)	1(2.1)	2(3.6)	
		Palmar/plantar	-	1(3.7)	1(3.7)	-	1(2.1)	-	
		Medial	3(3.1)	1(3.7)	2(7.4)	-	3(6.4)	-	
		Lateral	1(1)	1(3.7)	-	-	1(2.1)	-	
	<i>Level</i>	Proximal to the PSB	1(1)	-	2(7.4)	-	2(4.3)	-	0.038
		At the level of the PSB	4(4.1)	-	2(7.4)	1(1.1)	3(6.4)	2(3.6)	
		Distal to the PSB	2(2.1)	-	2(7.4)	1(1.1)	2(4.3)	-	
		Body	1(1)	1(3.7)	1(3.7)	-	2(4.3)	-	

	<i>Pathology type</i>	Irregularity	4(4.1)	1(3.7)	1(3.7)	2(2.3)	4(8.5)	2(3.6)	
		Enlargement	1(1)	-	2(7.4)	-	2(4.3)	-	0.038
		Fluid-based hyperintensity	4(4.1)	1(3.7)	2(7.4)	1(1.1)	2(4.3)	2(3.6)	
		Adhesions	3(3.1)	-	2(7.4)	-	-	-	
		Chronic fibrosis	1(1)	-	1(3.7)	-	-	-	

