

Supplementary Materials

Evaluation of Two Extraction Methods for the Analysis of Hydrophilic Low Molecular Weight Compounds from *Ganoderma lucidum* Spores and Antiproliferative Activity on Human Cell Lines

Maria Michela Salvatore ^{1,†}, **Vincenza De Gregorio** ^{2,3,†}, **Monica Gallo** ⁴, **Maria Michela Corsaro** ¹, **Angela Casillo** ¹, **Raffaele Vecchione** ^{3,*}, **Anna Andolfi** ^{1,*}, **Daniele Naviglio** ^{1,‡} and **Paolo Antonio Netti** ^{2,3,5,‡}

¹ Department of Chemical Sciences, University of Naples Federico II, 80126 Naples, Italy; mariamichela.salvatore@unina.it (M.M.S.); corsaro@unina.it (M.M.C.); angela.casillo@unina.it (A.C.); naviglio@unina.it (D.N.)

² Interdisciplinary Research Centre on Biomaterials (CRIB), University of Naples Federico II, 80125 Naples, Italy; vincenza.degregorio@unina.it (V.DG.); nettipa@unina.it (P.A.N.)

³ Center for Advanced Biomaterials for Health Care @ CRIB, Istituto Italiano di Tecnologia, 80125 Naples, Italy

⁴ Department of Molecular Medicine and Medical Biotechnology, University of Naples Federico II, 80131 Naples, Italy; mongallo@unina.it

⁵ Department of Chemical, Materials and Industrial Production Engineering (DICMAPI) University of Naples Federico II, Naples 80125, Italy

* Correspondence: raffaele.vecchione@iit.it (R.V.); andolfi@unina.it (A.A.); Tel.: +39-081-2539179 (A.A.)

† These authors contributed equally to this work.

‡ These authors contributed equally to this work.




Figure S1. Comparison of proliferative activities on Caco-2 (non-invasive) and HCT-116 (invasive) cell lines of the dialyzed extracts from Rapid Solid-Liquid Dynamic Extraction (RSLDE_{IN+OUT}) after 48 h exposure.