

Supplementary Materials

Assessment of Handler Exposure to Pesticides from Stretcher-Type Power Sprayers in Orchards

Zhinan Wang ^{1,†}, Yuxi Meng ^{1,2,†}, Xiangdong Mei ¹, Jun Ning ¹, Xiaodong Ma ^{2,*}
and Dongmei She ^{1,*}

¹ State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, No.2 Yuanmingyuan West Road, Beijing 100193, China; 82101182339@caas.cn (Z.W.); myxiixi@cau.edu.cn (Y.M.); xdmei@ippcaas.cn (X.M.); jning@ippcaas.cn (J.N.)

² College of Science, China Agricultural University, Beijing 100193, China

* Correspondence: dongxm@cau.edu.cn (X.M.); shedongmei@caas.cn (D.S.)

† Zhinan Wang and Yuxi Meng are co-first authors and contributed equally to this work.

Received: date; Accepted: date; Published: date

Table S1. Mass spectrometric parameters for UPLC-MS determination of SYP-9625 concentrations.

Compound	Precursor (m/z)	Precursor (m/z)	Dwell (s)	Cone Voltage (V)	Collision Energy (V)
SYP-9625	394.34	108.97a	0.161	38	44
	394.34	310.16a	0.161	38	24

¹ The transitions were used for quantification.




Figure S1. Structure of SYP-9625.




Figure S2. The UPLC-MS chromatogram for SYP-9625.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).