

Supporting Information

Estimating Full-Coverage PM_{2.5} Concentrations Based on Himawari-8 and NAQPMS Data over Sichuan-Chongqing

Qiaolin Zeng^{1,2,3}, Hao Zhu^{1,2}, Yanghua Gao^{1,2,*}, Tianshou Xie³, Sizhu Liu³ and Liangfu Chen⁴

- 1 Chongqing Institute of Meteorological Sciences, Chongqing 401147, China; zengql@cqupt.edu.cn (Q.Z.); s180201040@stu.cqupt.edu.cn (H.Z.)
 - 2 Chongqing Engineering Research Center of Agrometeorology and Satellite Remote Sensing, Chongqing 401147, China
 - 3 The College of Computer Science and Technology, Chongqing University of Posts and Telecommunications, Chongqing 400065, China; s200231036@stu.cqupt.edu.cn (T.X.); liusz@cqupt.edu.cn (S.L.)
 - 4 The Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100094, China; chenlf@radi.ac.cn
- * Correspondence: yanghuagaocq@gmail.com





Figure S1. $E_{\text{ext}}(\text{RH})$ fitting Leshan Ji Jiangxian station. The red solid lines are fitting lines of the scattered dots.






Figure S2. $E_{ext}(RH)$ fitting Yibin Xingwenzhong station. The red solid lines are fitting lines of the scattered dots.





Figure S3. $E_{ext}(RH)$ fitting Bazhong Tongjiangzhongxue station. The red solid lines are fitting lines of the scattered dots.






Figure S4. $E_{ext}(RH)$ fitting Chongqing Guanyinqiao station. The red solid lines are fitting lines of the scattered dots.