

Article

Characterization of Sidr (*Ziziphus* spp.) Honey from Different Geographical Origins

Ahmed G. Hegazi ¹, Fayez M. Al Guthami ², Mohamed F. A. Ramadan ³, Ahmed F. M. Al Gethami ³,
A. Morrie Craig ⁴ and Salud Serrano ^{5,*}

¹ Department of Zoonotic Diseases, National Research Centre, Dokki, Giza 12622, Egypt

² Al Guthami Foundation, Saudi Arabia

³ Alnahalajwal Foundation, Makkah 21926, Saudi Arabia

⁴ Department of Biomedical Sciences, College of Veterinary Medicine, Oregon State University, Corvallis, OR 97331, USA

⁵ Department of Food Science and Technology, Faculty of Veterinary medicine, University of Córdoba, 14071 Córdoba, Spain

* Correspondence: bt2sejis@uco.es; Tel.: +34-957-212-000

Featured Application: The current investigation was conducted to characterize the Sidr honey through melissopalynological analysis, its physicochemical, and biochemical properties, antimicrobial, antioxidant activities as well as total phenolic and total flavonoid contents. For this purpose, Sidr honey samples collected from the Saudi market imported from 12 different countries were analyzed.

Abstract: The current investigation was conducted to assess the melissopalynological, physicochemical, and biochemical properties, antimicrobial and antioxidant activities as well as total phenolic and total flavonoid contents of 794 Sidr honey samples collected from the Saudi market that had been imported from 12 different countries. Testing Sidr honey from different countries showed different levels of growth suppression observed against five drug resistant bacterial strains. The pathogenic strains were *Staphylococcus aureus*, *Streptococcus mutans*, *Klebsiella pneumoniae*, *Escherichia coli* and *Pseudomonas aeruginosa*. The antimicrobial activity showed growth suppression levels which varied according to the origin of the honey. The comparative study of Sidr honeys revealed a strong correlation between total polyphenol and flavonoid contents and significant radical scavenging activities in particular Egyptian and Saudi Arabian honeys. The melissopalynological and physicochemical properties of different Sidr honeys complied with the recommendations of the WHO *Codex Alimentarius*, the European Union standards for honey quality, and the Gulf Technical Regulation on honey (GSO 147:2008-Standards Store-GCC Standardization Organization). It was concluded that Sidr honey from different geographical areas has the capacity to suppress the growth of pathogenic bacteria and perform significant radical scavenging activities.

Keywords: melissopalynological and physicochemical analysis; antibacterial activity; antioxidant activity; Sidr honey



Citation: Hegazi, A.G.; Al Guthami, F.M.; Ramadan, M.F.A.; Al Gethami, A.F.M.; Craig, A.M.; Serrano, S. Characterization of Sidr (*Ziziphus* spp.) Honey from Different Geographical Origins. *Appl. Sci.* **2022**, *12*, 9295. <https://doi.org/10.3390/app12189295>

Academic Editor: Dimitris Mossialos

Received: 4 August 2022

Accepted: 9 September 2022

Published: 16 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Apis mellifera worker bees produce honey from collected from plants [1,2]. Honey can be either monofloral or multifloral depending on the pollen plant source [3]. Honey composition is influenced by many factors such as plant species, climate, environmental conditions, harvesting time, beekeeper's handling, processing and storage conditions [4]. Honey quality depends on the chemical composition of the source plants, and on the climatic conditions and soil mineral composition [4]. The main components in honey are fructose and glucose (monosaccharides) (65%), and water (18–20%), and the minor components include free amino acids, aroma compounds, vitamins, organic acids, minerals,

and phenolic acids and flavonoids [5]. The beehive products (honey, propolis, royal jelly, bee venom, and beeswax) contain several [6,7] important medicinal compounds. Major honey bioactive compounds include phenolics, methylglyoxal, royal jelly proteins (MRJPs), and oligosaccharides that have anti-inflammatory, antimicrobial and antioxidant activities [7]. The presence of novel antimicrobial compounds in natural authentic honey is well documented [7–9]. Adulteration of honey either by adding cheap sugar syrup or by feeding the bees on sugar solutions is a major worldwide problem that leads to the loss of many biological properties of honey including its effective antimicrobial activity [10].

Therefore, the potential use of specific types of fresh authentic honey as candidate antibacterial agents is based on their physicochemical characteristics. Although monofloral honey is appreciated and it is also more expensive [11], Devi & Jangir (2018) [12] report that volatile compounds in honey come from its diverse floral origin and could be the source of its biological activity and medical importance.

Sidr (*Ziziphus* spp.) honey is produced from *Ziziphus* trees [13]. In Saudi Arabia, this is the most valuable honey, and customers believe that this type of honey is superior to other honey imported from other countries around the world or produced locally [13].

Several Sidr honeys are produced in different parts of the world. However, the available information on their physical and chemical properties is limited [14]. This honey is often subjected to adulteration due to limited availability and its high price [15]. Honey color and composition are dependent greatly on the geographical and botanical origins [10–16]. In addition, geographical and botanical origins are the two main criteria for general honey authenticity according to different national standards on honey authenticity and the *Codex Alimentarius* Standard [17]. Therefore, the current investigation was conducted to assess the melissopalynological, physicochemical, and biochemical properties, antimicrobial and antioxidant activities as well as total phenolic and total flavonoid contents of Sidr sider honey samples collected from the Saudi market that had been imported from 12 zones in different countries.

2. Materials and Methods

2.1. Materials

All used reagents and chemicals were of analytical grade and were purchased from Sigma (St. Louis, MO, USA).

2.2. Honey Samples

A total of 794 fresh Sidr honey samples (1 kg each) were collected from the Saudi Arabia markets during 2021. They were imported from 12 geographical areas in different countries. Each honey sample was collected in a sterile universal glass container and kept at 2–8 °C until tested. Melissopalynological analysis was used to corroborate the samples' level of authenticity as Sidr honey, which means that the honey must have at least 55% of pollen from a specific floral source [18]. The collected honey samples are shown in Table 1.

2.3. Melissopalynological and Physicochemical Analysis

Melissopalynological and physicochemical analyses were performed [19]. The pollen content was identified by the sedimentation technique as described by [18,20]. Other parameters determined were color, water content [21], insoluble solids [22], pH, acidity, optical rotation, and electrical conductivity [23]. The assessments for sugar content, inverted sugars, glucose, fructose, fructose/glucose ratio, fructose + glucose %, glucose/moisture ratio, and sucrose were performed by HPLC-DAD according to standard methods [24]. Additionally, diastase enzyme activity [16], and hydroxymethylfurfural (HMF) were analyzed [25].

Table 1. Geographical origin of Sidr honey samples.

Honey Samples	Number of Samples
Emirates	72
China	64
Iraq	53
Pakistan	67
Bashawer	75
Panjab	60
Saudi Arabia	65
Kashmir	90
Libya	44
Egypt	66
India	75
Yemen	63

2.4. Detection of Total Phenolic Content (TPC)

TPC was detected using Folin–Ciocalteu reagent [26] following [27,28]. The honey solution (0.5 mL) was mixed with 2.5 mL Folin–Ciocalteu reagent (2N) and incubated for 5 min. Subsequently, 2 mL sodium carbonate solution (75 gr/L) was added and incubated for 2 h at 25 °C. The absorbance of the solution was measured at 765 nm after incubation using a UV-Visible spectrophotometer (Perkin-Elmer Lambda 25, Waltham, MA, USA). For the calibration curve preparation, gallic acid (0–1000 mg/L) was used as a standard. The mean values of triplicate assays of TPC are reported, expressed as milligrams of gallic acid equivalent (GAE) per gram of honey [29].

2.5. Determination of Total Flavonoid Content (TFC)

TFC was determined using a 5 mL sample of diluted honey at 0.1 g/mL concentration. This solution was mixed with 5 mL of 2% aluminum chloride (AlCl_3) for the determination of TFC. The mixture was then incubated for 10 min at 25 °C. The absorbance of the formed complex was measured at 415 nm using a UV-Visible spectrophotometer. Rutin was the standard chemical used for the calibration curve preparation, with a concentration 0–100 mg/L. The mean values of triplicate assays of TFC are reported, expressed as milligrams of rutin equivalent (RE) per gram of honey [28,29].

2.6. Antioxidant Assay to Determine the DPPH Scavenging Activity

An antioxidant assay was used to determine the DPPH scavenging activity of the different honey samples. This test is based on the change in the absorbance that results from reducing the purple DPPH radical using an oxidizing antioxidant. The scavenging effect of vitamin C and caffeic acid as well as the honey samples corresponded to the quenching intensity of 1,1-diphenyl-2-picrylhydrazyl (DPPH) as carried out by [30]. The absorbance resulting from reducing the purple DPPH radical by an oxidizing antioxidant was measured at 520 nm.

To determine the percent inhibition, the antioxidant ability of vitamin C samples was measured as a decrease in absorbance of DPPH solution (purple DPPH reduction) due to the addition of the sample solution. The absorbance value of the DPPH solution measurement results before and after the addition of the sample solution was calculated as percent inhibition. Using the percent inhibition obtained, a linear regression equation with the sample concentration ($\mu\text{g/mL}$) on the x axis and the inhibition value on the y axis was calculated. The antioxidant activity of the test material was assessed by calculating the inhibitory concentration 50% (IC₅₀) using the following formula: $50 = ax + b$. The IC₅₀ value indicates the concentration of the test sample ($\mu\text{g/mL}$) which results in a 50% DPPH reduction (able to inhibit or reduce the oxidation process by 50%). The results of the calculation are entered into the obtained regulatory equation [31].

2.7. Bacterial Strains

The five antibiotic-resistant bacterial strains (Gram-positive and Gram-negative) included *Staphylococcus aureus* (ATCC 25923), *Streptococcus mutans* (1815T), *Escherichia coli* (ATCC 35218), *Klebsiella pneumoniae* (ATCC 27736), *Escherichia coli* (ATCC 35218) and *Pseudomonas aeruginosa* (ATCC 27853). These microorganisms were provided and maintained by the Department of Zoonotic Diseases, National Research Centre, Egypt. Each bacterial strain suspension was prepared by inoculating fresh stock culture into the broth tube containing 10 mL Muller Hinton Broth (Sigma Aldrich company). The inoculated tubes were incubated aerobically at 37 °C for 24 hr. The bacterial suspension was adjusted by comparison with 0.5 Mc Farland turbidity standards (5×10^7 cells/mL). It was then further diluted to obtain a final of 5×10^6 cells/mL. Physiological saline PBS pH 7.2 was used for all dilution steps under aseptic conditions. These bacterial strains were enriched on selective broth for bacterial propagation [32]. A separate tube containing 40 µL of 21.30% honey concentration was mixed with 0.20 µL/10 mL from the enriched broth of each propagated *S. aureus*, *S. mutans*, *K. pneumoniae*, *E. coli*, and *P. aeruginosa* [9,33,34]. These tubes were incubated at 37 °C for 24 h. The growths of the control bacterial strains and the inhibition of the bacterial growth due to mixing with honey were measured using the disc diffusion method. The mean values of inhibition were calculated from triplicate readings in each test. Evaluations of the antibacterial activity of different honey dilutions were performed according to Hegazi et al., 2017; 2020 and 2021) [9,33,34]. The results of antibacterial activity against different examined bacteria were recorded.

2.8. Minimum Inhibitory Concentration (MIC)

The MIC of different samples of Sidr honey were determined by a two-fold serial dilution method [6]. Serial dilution of 100 mg/mL for the rest of the samples were performed separately to achieve 50, 25, 12.50, 6.25, 3.12, 1.56, 0.78 mg/mL, and 390, 195, 97 µg/mL concentrations were used for the MIC determination. Briefly, 100 µL of varying sample concentrations were added separately to the test tubes containing 9 mL of the standardized suspension of the tested bacteria (10^8 CFU/mL). The test tubes were incubated at 37 °C for 24 h. Control tests with the test organisms were performed using distilled water instead of honey. The lowest concentration of these samples with no visible growth was taken as the MIC [6].

2.9. Statistical Analysis

The tests were conducted in triplicate and the statistical analysis then performed using SPSS Ver. 21 (IBM, New York, NY, USA) software. A one-way ANOVA was applied for comparisons between and within the tested groups. The mean \pm standard error (SE) is presented for all data and *p* values less than 0.05 were considered significant.

3. Results

A total of 794 Sidr honey types that had been imported from different countries were collected from the Saudi market. Melissopalynological analysis of Sidr honey from different geographical origins proved that not only the expected pollen type based on the specific source of nectar but different pollen from some other sources were also present depending on the geographical origin (Table 2 and Figure 1). The Sidr honey included pollen from the following species: *Ziziphus jujuba*, *Conocarpus erectus*, and *Eleusine coracana* (Emirates); *Ziziphus jujuba*, and *Brassica napus* (China); *Ziziphus jujuba*, and *Cichorium intybus* (Iraq); *Ziziphus spina-christi*, *Amaranthus blitum*, *Capsella bursa-pastoris*, and *Oryza meyeriana* (Pakistan); *Ziziphus spina-christi*, *Rhanterium epapposum*, *Sesamum indicum*, and *Oryza meyeriana* (Bashawer); *Ziziphus spina-christi*, *Amaranthus blitum*, and *Oryza meyeriana* (Panjab); *Ziziphus jujuba*, *Acacia asak*, and *Blepharis Ciliaris* (Saudi Arabia); *Ziziphus spina-christi*, *Capsella bursa-pastoris*, *Amaranthus blitum*, and *Chrysanthemum leucanthemum* (Kashmir); *Ziziphus jujuba*, and *Cynara aurantica* (Libya); *Ziziphus lotus*, *Oryza meyeriana*, *Zea mays*, and *Brassica*

tournefortii Gouan (Egypt); *Ziziphus jujuba*, *Oryza meyeriana*, and *Acacia asak*, (India); and *Ziziphus jujuba*, and *Acacia asak* (Yemen).

Table 2. Melissopalynological analysis of Sidr honey from different origins.

Honey Origin	Botanical Family	Botanical Species	Pollen Count
Emirates	Rhamnaceae	<i>Ziziphus jujuba</i>	++++
	Combretaceae	<i>Conocarpus erectus</i>	++
	Poaceae	<i>Eleusine coracana</i>	++
China	Rhamnaceae	<i>Ziziphus jujuba</i>	++++
	Brassicaceae	<i>Brassica napus</i>	++
Iraq	Rhamnaceae	<i>Ziziphus jujuba</i>	++++
	Asteraceae	<i>Cichorium intybus</i>	++
	Rhamnaceae	<i>Ziziphus spina-christi</i>	++++
Pakistan	Amaranthaceae	<i>Amaranthus blitum</i>	++
	Brassicaceae	<i>Capsella bursa-pastoris</i>	++
	Poaceae	<i>Oryza meyeriana</i>	+
Bashawer	Rhamnaceae	<i>Ziziphus spina-christi</i>	++++
	Asteraceae	<i>Rhanterium epapposum</i>	++
	Pedaliaceae	<i>Sesamum indicum</i>	+++
	Poaceae	<i>Oryza meyeriana</i>	+
Panjab	Rhamnaceae	<i>Ziziphus spina-christi</i>	++++
	Amaranthaceae	<i>Amaranthus blitum</i>	++
	Poaceae	<i>Oryza meyeriana</i>	+
Saudi Arabia	Rhamnaceae	<i>Ziziphus jujuba</i>	++++
	Mimosaceae	<i>Acacia asak</i>	+
	Acanthaceae	<i>Blepharis Ciliaris</i>	++
	Rhamnaceae	<i>Ziziphus spina-christi</i>	++++
Kashmir	Brassicaceae	<i>Capsella bursa-pastoris</i>	+++
	Amaranthaceae	<i>Amaranthus blitum</i>	++
	Asteraceae	<i>Chrysanthemum leucanthemum</i>	+
	Rhamnaceae	<i>Ziziphus jujuba</i>	++++
Libya	Asteraceae	<i>Cynara auranitica</i>	++
	Rhamnaceae	<i>Ziziphus lotus</i>	++++
Egypt	Poaceae	<i>Oryza meyeriana</i>	++
	Tamaricaceae	<i>Zea mays</i>	++
	Brassicaceae	<i>Brassica tournefortii</i>	+
		<i>Gouan</i>	
India	Rhamnaceae	<i>Ziziphus jujuba</i>	++++
	Poaceae	<i>Oryza meyeriana</i>	++
	Mimosaceae	<i>Acacia asak</i>	++
Yemen	Rhamnaceae	<i>Ziziphus jujuba</i>	++++
	Mimosaceae	<i>Acacia asak</i>	++

++++: more than 70%; +++: 70%; ++: 60%; +: 50%.

The physicochemical properties as shown in Table 3 reveal that the Sidr honey samples were comparable in water content, which ranged from 14.2 ± 0.41 (Libya) to $17.06 \pm 0.47\%$ (Iraq). The optical rotation ranged from -1.13° (Egypt) to -2.42° (China). The colors observed ranged from extra white (China, Kashmir and India) to deep amber (Yemen). The pH varied from 3.6 ± 0.1 (Egypt) to 7.4 ± 0.1 (Yemen). The acidity also varied from 7.7 ± 0.5 (Egypt) to 11.3 ± 1.4 meq/l (Libya). Electrical conductivity ranged from 0.55 ± 0.02 (China, and Libya) to 1.40 ± 0.39 mS/cm (Pakistan). The insoluble solids ranged from 0.05 ± 0.02 and $0.05 \pm 0.03\%$ (Saudi Arabia, and Libya, respectively) to 1.36 ± 0.02 (Pakistan).




Figure 1. Pollen grain images from microscope preparations of Sidr honey from different geographical origins.

Table 3. Physicochemical parameters of Sidr honey samples from different origins.

Geographical Origin	Samples (n)	Water Content (g/100 g)	Optical Rotation (°)	Color	Insoluble Solids (%)	pH	Acidity (meq/l)	Electrical Conductivity (mS/cm)
Emirates	72	16.26 * ± 0.64	−1.38	Light Amber	0.06 ± 0.01	6.5 * ± 0.1	8.4 * ± 0.9	1.18 ** ± 0.11
China	64	15.37 ± 0.57	−2.42 **	Extra white	0.9 ± 0.02	6.2 * ± 0.6	7.0 ± 0.3	0.55 ± 0.02
Iraq	53	17.06 ** ± 0.47	−1.86	Amber	0.08 ± 0.02	5.6 ± 0.1	9.0 ** ± 0.8	0.93 ± 0.01
Pakistan	67	14.84 ± 0.38	−1.92 *	White	1.36 * ± 0.02	6.8 * ± 0.1	10.8 ** ± 1.2	1.40 ** ± 0.39
Bashawer	75	14.29 ± 1.06	−1.79	White	1.02 ± 0.01	6.1 * ± 0.0	8.9 * ± 0.9	0.96 * ± 0.04
Panjab	60	15.7 ± 0.34	−1.42	Amber	1.09 ± 0.09	6.2 * ± 0.0	8.8 * ± 1.5	1.14 * ± 0.09
Saudi Arabia	65	14.6 ± 0.21	−1.36	Light Amber	0.05 ± 0.02	5.4 ± 0.2	10.6 ** ± 0.0	0.84 ± 0.04
Kashmir	90	15.53 ± 0.38	−1.89	Extra White	1.35 * ± 0.24	3.9 ± 1.1	8.8 * ± 0.9	0.79 ± 0.16
Libya	44	14.2 ± 0.41	−1.98 *	Amber	0.05 ± 0.03	5.8 ± 0.1	11.3 ** ± 1.4	0.55 ± 0.04
Egypt	66	14.96 ± 0.22	−1.13	Amber	1.35 * ± 0.02	3.6 ± 0.1	7.7 ± 0.5	0.65 ± 0.17
India	75	15.78 ± 0.52	−1.69	Extra White	1.05 ± 0.02	7.2 ** ± 0.1	8.6 ± 1.4	0.99 * ± 0.07
Yemen	63	16.1 ** ± 0.53	−2.09 **	Deep Amber	1.04 ± 0.01	7.4 ** ± 0.1	12.2 ** ± 0.3	0.99 * ± 0.13

** Highly significant; * Significant.

The glucose, fructose, sucrose, diastase activity and HMF levels in different Sidr honey samples are shown in Table 4. Glucose was detected at the lowest level ($24.77 \pm 0.65\%$) in India Sidr honey, whereas the highest level ($28.89 \pm 0.11\%$) was observed in Iraq Sidr honey. The level of fructose ranged between 32.79 ± 0.64 (India) and 36.01 ± 1.05 (Iraq). The sucrose level ranged from $1.24 \pm 0.37\%$ (Pakistan) to $3.79 \pm 0.27\%$ (Egypt). Diastase activity had a range of 9.75 ± 1.78 D.U. (Yemen) to 17.4 ± 2.16 D.U. (Egypt). The lowest HMF (mg/kg) was observed in Libya Sidr honey (11.07 ± 4.38), whereas the highest level was detected in Yemen Sidr honey (25.11 ± 6.63).

Table 4. Glucose, fructose, sucrose, diastase activity and HMF levels in different Sidr honey samples.

Geographical Origin	Samples (n)	Glucose (g/100 g)	Fructose (g/100 g)	Fructose/Glucose Ratio	Fructose + Glucose %	Glucose/Moisture Ratio	Sucrose (g/100 g)	Diastase Activity (D.U.)	HMF mg/kg
Emirates	72	25.06 ± 1.15	33.26 ± 0.48	1.33 *	58.33 * ± 1.28	1.54	3.01 ± 0.92	13.73 ± 0.95	12.64 ± 1.87
China	64	26.64 ± 0.70	35.02 ** ± 0.62	1.31	61.66 ** ± 1.16	1.73 *	2.33 ± 0.83	17.36 ** ± 1.16	17.55 ** ± 4.47
Iraq	53	28.89 * ± 0.11	36.01 ** ± 1.05	1.25	64.9 ** ± 0.95	1.69	3.5 ± 0.23	12.7 ± 1.53	13.8 ± 1.91
Pakistan	67	25.7 ± 0.83	34.43 * ± 0.48	1.26	60.14 ** ± 1.10	1.73 *	1.24 ± 0.37	11.53 ± 1.28	12.73 ± 1.38
Bashawer	75	27.67 ± 0.78	33.94 * ± 1.13	1.22	61.61 ** ± 1.61	1.93 **	1.63 ± 0.48	13.06 ± 1.35	12.52 ± 1.49
Panjab	60	25.61 ± 0.96	33.48 ± 0.46	1.31	59.1 * ± 1.18	1.631	3.78 * ± 0.84	16.23 * ± 2.58	14.95 ± 0.90
Saudi Arabia	65	27.53 * ± 0.27	34.94 * ± 0.31	1.27	62.47 ** ± 0.52	1.88 **	4.52 ** ± 0.46	15.19 ± 1.31	17.6 ** ± 2.80
Kashmir	90	25.77 * ± 0.89	33.38 ± 1.21	1.29	59.16 * ± 1.93	1.66	2.83 ± 0.89	11.27 ± 1.86	18.94 ** ± 6.06
Libya	44	25.61 ± 1.14	35.99 ** ± 0.66	1.4 *	61.61 ** ± 1.64	1.8 **	0.175 ± 0.06	17.15 ** ± 1.21	11.07 ± 4.38
Egypt	66	26.02 * ± 0.98	33.69 ± 0.91	1.29	59.71 * ± 1.61	1.73 *	3.79 * ± 0.27	17.4 ** ± 2.16	16.6 ± 0.12
India	75	24.77 ± 0.65	32.79 ± 0.64	1.32	57.56 * ± 0.91	1.56	3.36 * ± 0.24	10.97 ± 2.45	19.8 ** ± 0.58
Yemen	63	27.99 * ± 0.48	34.73 * ± 0.52	1.24	62.72 ** ± 0.83	1.73 *	2.16 ± 0.65	9.75 ± 1.78	25.11 ** ± 6.63

** Highly significant; * Significant.

The TPC (mg GAE/100 g honey), TFC (mg RE/100 g honey) and DPPH (mg AAE/100 g honey) content is shown in Table 5. The highest levels for the three parameters were found in Egypt Sidr honey 159.3 ± 15.32, 83.1 ± 18.33 and 177.8 ± 10.51, respectively, whereas the lowest level for total phenolics was detected in Yemen Sidr honey. On the other hand, the lowest levels of TFC and DPPH were detected in China Sidr Honey (35.1 ± 7.10 and 75.1 ± 7.57, respectively).

Table 5. TPC, TFC and DPPH of the Sidr honeys.

Origin	Samples (n)	Total Phenolic (mg GAE/100 g Honey)	Total Flavonoid (mg RE/100 g Honey)	DPPH (mg Ascorbic Acid Equation/100 g Honey)
Emirates	72	138.0 ± 9.16 *	58.0 ± 9.57 *	98.1 ± 10.90
China	64	121.0 ± 10.24	35.1 ± 7.10	75.1 ± 7.57
Iraq	53	139.0 ± 16.45 *	47.0 ± 14.10	167.0 ± 13.87 **
Pakistan	67	136.0 ± 9.15	49.0 ± 7.48	79.0 ± 8.25
Bashawer	75	134.1 ± 10.58	41.3 ± 9.78	98.3 ± 10.57
Panjab	60	125.5 ± 13.95	35.3 ± 7.52	135.3 ± 13.00 *
Saudi Arabia	65	144.9 ± 10.66 *	81.5 ± 13.47 **	131.3 ± 15.18 *
Kashmir	90	136.1 ± 11.38	45.0 ± 13.22	125.0 ± 14.132
Libya	44	138.2 ± 13.30 *	43.6 ± 11.13	111.0 ± 11.82
Egypt	66	159.3 ± 15.32 *	83.1 ± 18.33 **	177.8 ± 10.51 **
India	75	122.0 ± 7.24	68.0 ± 11.59 *	109.0 ± 12.94
Yemen	63	118.9 ± 9.48	38.13 ± 9.71	95.0 ± 10.41

** Highly significant; * Significant.

The antibacterial activity of the Sidr honey from twelve geographical origins was evaluated according to the zone of inhibition. The antibacterial potency of honey was investigated against various Gram-negative and Gram-positive pathogenic bacteria. All geographical honey types showed high antibacterial activity against most of the tested bacterial strains. *Staphylococcus aureus* showed the highest zones of inhibition 25.00 ± 0.58, 23.00 ± 0.58 mm in Egypt and Saudi Arabia honey respectively, followed by Emirates (21.33 ± 0.88 mm) and Punjab (21.00 ± 0.11 mm) honeys, while Egyptian (29.33 ± 0.64 mm), Saudi (23.00 ± 0.22 mm), and Iraq (22.00 ± 0.58 mm) honeys revealed the highest zones of inhibition against *Streptococcus mutans*. The highest zones of inhibition against *Klebsiella pneumoniae* were observed in the Saudi (25.00 ± 0.61 mm), Egyptian (24.00 ± 0.34 mm),

Emirates (22.00 ± 0.68 mm), and Libya (21.00 ± 0.31 mm) honey samples. Additionally, Egyptian (29.16 ± 0.60 mm), Saudi (27.00 ± 0.61 mm) and Libya (20.33 ± 0.88 mm) honey were highly effective against *Escherichia coli*. Significantly higher zones of inhibition against *Pseudomonas aeruginosa* were shown for Saudi (40.67 ± 0.67 mm) and Egyptian (29.00 ± 0.58 mm) honeys (Table 6).

Table 6. The inhibition zone of Sidr honey against various pathogenic microorganisms by well diffusion method.

Antibacterial Activity Honey Origin	Gram-Positive			Gram-Negative	
	<i>Staphylococcus aureus</i>	<i>Streptococcus mutans</i>	<i>Klebsiella pneumoniae</i>	<i>Escherichia coli</i>	<i>Pseudomonas aeruginosa</i>
Emirates	21.33 ± 0.88	18.33 ± 0.88	22.00 ± 0.68	15.33 ± 0.33	16.00 ± 1.15
China	16.00 ± 0.58	16.33 ± 0.88	11.00 ± 0.56	8.00 ± 0.58	9.67 ± 0.33
Iraq	19.00 ± 0.58	22.00 ± 0.58	11.00 ± 0.48	17.00 ± 0.48	15.00 ± 0.10
Pakistan	18.00 ± 0.58	16.00 ± 0.48	11.00 ± 0.44	14.00 ± 0.66	15.00 ± 0.51
Bashawer	8.00 ± 0.58	12.00 ± 0.32	11.00 ± 0.55	17.00 ± 0.38	14.00 ± 0.58
Panjab	21.00 ± 0.11	17.00 ± 0.58	10.33 ± 0.47	8.00 ± 0.64	9.67 ± 0.43
Saudi Arabia	$23.00 \pm 0.58^*$	$23.00 \pm 0.22^*$	$25.00 \pm 0.61^*$	$27.33 \pm 0.45^*$	$40.67 \pm 0.67^*$
Kashmir	11.00 ± 0.58	17.00 ± 0.66	15.00 ± 0.30	19.33 ± 1.85	14.00 ± 0.68
Libya	15.33 ± 0.33	18.33 ± 0.33	21.00 ± 0.31	20.33 ± 0.88	10.00 ± 0.58
Egypt	$25.00 \pm 0.58^*$	$29.33 \pm 0.64^*$	$24.00 \pm 0.34^*$	$29.16 \pm 0.60^*$	$29.00 \pm 0.58^*$
India	11.00 ± 0.58	10.33 ± 0.88	17.00 ± 0.58	15.00 ± 0.10	11.33 ± 0.88
Yemen	8.33 ± 0.33	9.67 ± 0.33	8.00 ± 0.44	11.00 ± 0.58	9.00 ± 0.58

* Significant.

The MIC of the Sidr honey from different geographical origins against the pathogenic bacteria *Staphylococcus aureus*, *Streptococcus mutans*, *Klebsiella pneumoniae*, *Escherichia coli*, and *Pseudomonas aeruginosa* is shown in Table 7. The results of the antibacterial activity assay showed significant differences in MIC values when antibiotic-resistant strains were tested. Analyzing MICs, the strongest antibacterial potential against all tested bacteria was observed for Saudi Arabian and Egyptian (MIC = 0.4 g/mL) honeys. Iraq, Panjab, Kashmir, Libya, and India honeys showed the strongest antibacterial potential against *Staphylococcus aureus*, *Streptococcus mutans*, and *Klebsiella pneumoniae* (MIC = 0.4 g/mL). In contrast, the weakest overall antibacterial activity was shown in honey samples from Emirates, China, Pakistan, and Bashawer. Yemen honey showed the lowest antibacterial potential against *Escherichia coli* (MIC = 0.05 g/mL).

Table 7. Minimal inhibitory concentration (MIC) of Sidr honey against antibiotic resistant bacterial strains.

MIC	MIC of Bacterial Strains (g/mL)				
	Gram-Positive			Gram-Negative	
Honey Origin	<i>Staphylococcus aureus</i>	<i>Streptococcus mutans</i>	<i>Klebsiella pneumoniae</i>	<i>Escherichia coli</i>	<i>Pseudomonas aeruginosa</i>
Emirates	0.1	0.2	0.2	0.2	0.2
China	0.1	0.2	0.2	0.2	0.2
Iraq	0.4 *	0.4 *	0.4 *	0.1	0.2
Pakistan	0.2	0.2	0.2	0.1	0.1
Bashawer	0.2	0.2	0.2	0.1	0.1

Table 7. Cont.

MIC of Bacterial Strains (g/mL)					
MIC	Gram-Positive			Gram-Negative	
Honey Origin	<i>Staphylococcus aureus</i>	<i>Streptococcus mutans</i>	<i>Klebsiella pneumoniae</i>	<i>Escherichia coli</i>	<i>Pseudomonas aeruginosa</i>
Panjab	0.4 *	0.4 *	0.4 *	0.2	0.4 *
Saudi Arabia	0.4 *	0.4 *	0.4 *	0.4 *	0.4 *
Kashmir	0.4 *	0.4 *	0.4 *	0.1	0.2
Libya	0.4 *	0.4 *	0.4 *	0.1	0.1
Egypt	0.4 *	0.4 *	0.4 *	0.4 *	0.4 *
India	0.4 *	0.4 *	0.4 *	0.1	0.1
Yemen	0.2	0.2	0.2	0.05	0.1
Drugs for positive control for growth inhibition					
Vancomycin	0.000001	0.000001	0.00000004	0.000001	0.000000016
Meropenem	0.000000064	0.00000004	0.000000064	0.000000064	0.000000006

* Significant.

4. Discussion

This research revealed that monofloral honey obtained from the *Ziziphus* spp. nectar of different geographical origins contained more than 70% of a specific pollen. The most important physicochemical parameter of honey is the water content. Low water content increases honey shelf life while high levels promote its fermentation during storage. The water content of the investigated honey samples was within the accepted range which, in general, should not exceed 20% [12,35]. During honey production by bees, the water content is affected by the relative humidity and temperature. Water content results for honey samples of different geographical origins have been documented previously for their water content [34,36], being 18.32 ± 0.67 g/100 g for Egyptian, 16.28 ± 0.22 g/100 g for Yemeni, 15.64 ± 0.30 g/100 g for Saudi and 14.73 ± 0.3 g/100 g for Kashmiri honey.

The sugar content of honey is widely used to assess the authenticity and the overall quality of honey [37]. Honey adulteration by mixing the honey with other cheaper sugar syrups is a frequent problem in the worldwide market [38]. For this reason, the sugar analysis of honey is an indicator of whether the honeybees were fed naturally with flower nectar or were fed with sugar solution. The use of artificial feeding is evident when the glucose content of honey is much higher than its fructose content [39].

The results of this investigation revealed that the sum of glucose and fructose (content of reducing sugars in honey) was within the accepted range and is consistent with the standardization and authenticity of honey as observed by Aljohar et al., (2018) [40]. It has been demonstrated that the most dominant sugar in honey is fructose [39] and Szczesna et al., (2021), in their study of the winter feeding of honeybees, observed the fructose to glucose ratio (F/G) was higher than 1.00 indicating the natural feeding of honeybees [41]. In all the honey samples in this study, sucrose content did not exceed 5%, which is the accepted level for honey to be considered authentic as observed by Kazeminia et al., (2021) [42]. A higher content of sucrose may indicate the artificial feeding of bees with some types of sugar syrup or adulteration of the honey [38]. Two important parameters used to prove the freshness of honey are hydroxymethylfurfural content and diastase activity [43,44]. There are many factors affecting diastase activity, including the physiological period of the colony, age of the bees, the quantity of nectar and its sugar content as well as the nectar collection period [45]. The WHO *Codex Alimentarius*, the European Union and the Gulf Technical Regulation on honey (GSO 147:2008-Standards Store-GCC Standardization Organization) [46] recommend that the maximum level for HMF content in honey does not

exceed 40 mg/kg but in countries with tropical temperatures, the HMF content of honey should not exceed 80 mg/kg [34,43,44]. In the present study, all the samples were within the allowed range for HMF content and diastase number. These parameters indicate the freshness of the samples which was maintained by preventing their exposure to heat and shortening the storage time before the experiment. In addition, the acidity of all the honey samples was found to be within the accepted range. The acidity in honey is due to the presence of organic acids, in particular, gluconic acid, which has been found to affect honey flavor, texture, shelf life, and stability [47].

The TPC ranged from 118.9 to 159.3 mg GAE/100 g honey, which is higher than the results found in previous studies in honey from India [48] (47–98 mg GAE/100 g honey), Poland [49] (71.7 to 202.6 µg/g honey), Argentina [29] (18.730–107.213 mg GAE/100 g honey), Burkina Faso [50] (32.59–114.75 mg GAE/100 g honey) Portugal [51] (30.87 to 87.27 mg GAE/100 g), or from Romania [52] (2–125 mg GAE/100 g honey).

Variable levels of TPC were reported in different honey types as observed by [53] who observed that the TPC of forest honey (806.10 mg GAE/kg honey) was significantly higher ($p < 0.05$) in comparison with acacia and polyfloral samples (68.48 and 87.46 mg GAE/kg honey, respectively). In addition, Roby et al., (2020) [36] determined the phenolic compounds of Egyptian honeys, with TPC amounting to 338.5 and 536.4 mg GAE kg⁻¹ in clover and citrus honeys, respectively. This variability was associated with the botanic origin of the honey, and multi-floral honey was found to have higher phenolic contents than monofloral honey [54]. The antioxidant, antiviral, antimicrobial, antifungal, and anti-inflammatory activities of honey are noteworthy due to phenolic compounds, especially flavonoids [55], with the quality of polyphenols being more important than their quantity [56].

The TPC and TFC of honey depend mainly on its botanical and geographical origins [57,58]. Considering the TFC results, it can be concluded that dark honey contains the highest concentration ($p < 0.05$). Flavonoids in Egyptian and Saudi Arabia Sidr honey (83.1 and 81.5 mg RE/100 g honey, respectively) are present in high amounts compared with other tested origins. Similar results were observed by analysis of three types of monofloral honey from Portugal which showed that dark honey was richer in phenolics content [59].

A DPPH radical scavenging method was used to determine the antioxidant activity of the honey samples. The Sidr honey from Egypt (177.8), Panjab (135.3), and Saudi Arabia (131.3) showed the highest level of DPPH radical scavenging activities ($p < 0.05$), compared with antioxidant activities of other tested honey. These results are in accordance with the results reported by van den Berg et al., (2008) about Buckwheat honey [60], Bueno-Costa et al., (2016) in Brazilian honeys [61] and Boussaid et al., (2018) in Tunisian honeys [62]. Alves et al., (2014) [63] demonstrated a positive relationship between phenolic concentration, antioxidant capacity, and the color of honey.

In our study, the antibacterial activity of all honeys may be attributed to the narrow ranges of their TFC, TPC, and sugar contents. There is a positive correlation between TPC and the antibacterial activity of honey [64], which is attributed to the inhibition of virulence factors in the pathogen [65].

The antibacterial activity of honey also results from the low pH and high osmolarity along with the hydrogen peroxide activity produced by the glucose oxidase enzyme [8,9,34,66,67]. The antibacterial activity of honey may also be attributed to the presence of lysozyme, methylglyoxal, and bee peptides as well as its high sugar content [9,34,68,69]. The presented results were in accordance with the research of Mandal & Mandal (2011) [70], Szweda (2017) [71], Libonatti et al., (2014) [72], Irish (2011) et al., [73], Morroni et al., (2018) [74], and Al Masaudi (2021) [69], which also confirmed that Gram-positive bacteria were more sensitive to the honey samples than Gram-negative ones [75,76]. Vancomycin and meropenem were used as reference antibiotics, with MIC values much lower than those obtained for honey, ranging from 0.1–64 µg/mL [77].

The use of honey to treat microbial infections has been investigated previously [34,51,67,76]. The honey antimicrobial properties are due to a high sugar osmolarity or the presence

of other biologically active compounds [33,77–80]. The osmotic stress of honey is due to the high content of various sugars in combination with its low moisture content, which prevents the spoilage of honey by microorganisms. Hydrogen peroxide (H₂O₂) is one of the antibacterial compounds of honey [81], whose presence and origin were confirmed by White et al., (1963) [82]. In the hydrogen peroxide formation, a glucose oxidase enzyme is added by the worker bee during the collection of the nectar, allowing the conversion of glucose to gluconic acid, whereby hydrogen peroxide is produced as a side product [82]. The antimicrobial activity of honey is directly related to its botanical origin [63] and is associated with the different content of phenolic compounds, flavonoids, and phenolic acids [68,83].

5. Conclusions

The physicochemical characteristics of honey determine its biological activity and at the same time they serve as tools for authentication. This research examines the authenticity of monofloral Sidr honey (*Ziziphus* spp.) from 12 different countries based on melissopalynological, physicochemical and bioactive compounds analyses. Additionally, the antimicrobial activity of each honey was determined, providing relevant information about its efficacy and clarifying its mechanisms of biological activity. Geographical differences were evident in the pollen profile of the samples. The physicochemical parameters were assessed according to the criteria from the different honey quality standards, and the biological activity revealed the Saudi and Egyptian Sidr honeys have the highest antibacterial activity as well as total phenolics and flavonoids contents. Our results identify Sidr honey as a promising natural product that can be potentially used as an alternative to synthetic antibiotics; however, further studies are also needed to identify and standardize protocols for the use of Sidr honey either in the protection against microbial infections or their treatments.

Author Contributions: Conceptualization, A.G.H.; methodology, M.F.A.R.; validation, M.F.A.R. and S.S.; formal analysis, M.F.A.R.; investigation, A.G.H., M.F.A.R. and A.F.M.A.G.; resources, F.M.A.G.; data curation, A.F.M.A.G. and A.M.C.; writing—original draft preparation, A.G.H. and A.M.C.; writing—review and editing, A.G.H., A.M.C. and S.S.; visualization, A.G.H. and A.M.C.; supervision, A.G.H. and S.S.; project administration, F.M.A.G.; funding acquisition, F.M.A.G. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Research Center of Egypt under registration number 12/5/1 and the Al Guthami Foundation (Saudi Arabia).

Institutional Review Board Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Silva, L.R.; Videira, R.; Monteiro, A.P.; Valentão, P.; Andrade, P.B. Honey from Luso region (Portugal): Physicochemical characteristics and mineral contents. *Microchem. J.* **2009**, *93*, 73–77. [[CrossRef](#)]
2. Habib, H.M.; Meqbal, F.T.; Kamal, H.; Souka, U.D.; Ibrahim, W.H. Physicochemical and biochemical properties of honeys from arid regions. *Food Chem.* **2014**, *153*, 35–43. [[CrossRef](#)]
3. Mustafa, G.; Iqbal, A.; Javid, A.; Hussain, A.; Bukhari, S.M.; Ali, W.; Saleem, M.; Azam, S.M.; Sughra, F.; Ali, A.; et al. Variations in nutritional profile of honey produced by various species of genus *Apis*. *Braz. J. Biol. Rev. Bras. Biol.* **2021**, *83*, e246651. [[CrossRef](#)]
4. Yeli, A.; Fabiola, M.-T.; Lobato-Rosales, G.; Landa, M.; Giovanni, Z.; Luis, L.-C.; García-Santamaría, E.; Fernández-Lambert, G. Influence variables for honey production using bees *Apis mellifera* in the Misantla region. *Rev. Mex. Cienc. Agrícolas* **2019**, *10*, 1353–1365. [[CrossRef](#)]
5. Bogdanov, S.; Jurendic, T.; Sieber, R.; Gallmann, P. Honey for Nutrition and Health: A Review. *J. Am. Coll. Nutr.* **2008**, *27*, 677–689. [[CrossRef](#)]

6. Gobin, I.; Crnković, G.; Magdalenić, M.; Begić, G.; Babić, A.; Lušić, D.; Vučković, D. Antibacterial potential of Croatian honey against antibiotic resistant pathogenic bacteria. *Med. Glas.* **2018**, *15*, 139–144. [[CrossRef](#)]
7. Fyfe, L.; Okoro, P.; Paterson, E.; Coyle, S.; McDougall, G.J. Compositional analysis of Scottish honeys with antimicrobial activity against antibiotic-resistant bacteria reveals novel antimicrobial components. *LWT—Food Sci. Technol.* **2017**, *79*, 52–59. [[CrossRef](#)]
8. Hegazi, A.G. Antimicrobial activity of different Egyptian honeys as comparison of Saudi Arabia honey. *Res. J. Microbiol.* **2011**, *6*, 488–495. [[CrossRef](#)]
9. Hegazi, A.; Guthami, F.; Al Ghetami, A.; Abd Allah, F.M.; Saleh, A.A.; Fouad, E.A. Potential antibacterial activity of some Saudi Arabia honey. *Vet. World* **2017**, *10*, 233. [[CrossRef](#)]
10. Dezmirean, D.S.; Mărghițaș, L.A.; Chirilă, F.; Copaciu, F.; Simonca, V.; Bobiș, O.; Erler, S. Influence of geographic origin, plant source and polyphenolic substances on antimicrobial properties of propolis against human and honeybee pathogens. *J. Apic. Res.* **2017**, *56*, 588–597. [[CrossRef](#)]
11. di Rosa, A.R.; Marino, A.M.F.; Leone, F.; Corpina, G.G.; Giunta, R.P.; Chiofalo, V. Characterization of Sicilian Honeys Pollen Profiles Using a Commercial E-Tongue and Melissopalynological Analysis for Rapid Screening: A Pilot Study. *Sensors* **2018**, *18*, 4065. [[CrossRef](#)] [[PubMed](#)]
12. Devi, A.; Jangir, J. Chemical characterization complemented with chemometrics for the botanical origin identification of unifloral and multifloral honeys from India. *Food Res. Int.* **2018**, *107*, 216–226. [[CrossRef](#)] [[PubMed](#)]
13. Taha, E.K.A.; Al-Kahtani, S.; Taha, R. Comparison of the physicochemical characteristics of sidr (*Ziziphus* spp.) honey produced by *Apis florea* F. and *Apis mellifera* L. *J. Apic. Res.* **2021**, *60*, 470–477. [[CrossRef](#)]
14. Azonewade, F.E.; Paraiso, A.; Agbangnan Dossa, C.P.; Dougnon, V.T.; N'Tcha, C.; Mousse, W.; Baba-Moussa, L. Physicochemical Characteristics and Microbiological Quality of Honey Produced in Benin. *J. Food Qual.* **2018**, *2018*. [[CrossRef](#)]
15. Crăciun, M.E.; Pârvolescu, O.C.; Donise, A.C.; Dobre, T.; Stanciu, D.R. Characterization and classification of Romanian acacia honey based on its physicochemical parameters and chemometrics. *Sci. Rep.* **2020**, *10*, 20690. [[CrossRef](#)]
16. Miłek, M.; Bocian, A.; Kleczyńska, E.; Sowa, P.; Dżugan, M. The Comparison of Physicochemical Parameters, Antioxidant Activity and Proteins for the Raw Local Polish Honeys and Imported Honey Blends. *Molecules* **2021**, *26*, 2423. [[CrossRef](#)]
17. Adamchuk, L.; Sukhenko, V.; Akulonok, O.; Bilotserkivets, T.; Vyshniak, V.; Lisohurska, D.; Lisohurska, O.; Slobodyanyuk, N.; Shanina, O.; Galyasnyj, I. Methods for determining the botanical origin of honey. *Potravin. Slovak J. Food Sci.* **2020**, *14*, 483–493. [[CrossRef](#)]
18. Louveaux, J.; Maurizio, A.; Vorwohl, G. Methods of Melissopalynology. *Bee World* **2015**, *59*, 139–157. [[CrossRef](#)]
19. Kirs, E.; Pall, R.; Martverk, K.; Laos, K. Physicochemical and melissopalynological characterization of Estonian summer honeys. *Procedia Food Sci.* **2011**, *1*, 616–624. [[CrossRef](#)]
20. Chukwuemeka, R.; Simeonov, V. Evaluation of Pollen and Chemical Composition of Honey Samples Sourced from Open Markets in Anambra State, Nigeria to Ascertain their Authenticity. *J. Appl. Life Sci. Int.* **2019**, *22*, 1–12. [[CrossRef](#)]
21. Chataway, H.D. The determination of moisture in honey. *Can. J. Res.* **2011**, *6*, 532–547. [[CrossRef](#)]
22. Nyau, V.; Mwanza, E.; Moonga, H. Physico-chemical qualities of honey harvested from different beehive types in Zambia. *Afr. J. Food 2013 Agric. Nutr. Dev.* **2013**, *13*, 7415. [[CrossRef](#)]
23. Yadata, D. Detection of the Electrical Conductivity and Acidity of Honey from Different Areas of Tepi. *Food Sci. Technol.* **2014**, *2*, 59–63. [[CrossRef](#)]
24. Zhou, J.; Yao, L.; Li, Y.; Chen, L.; Wu, L.; Zhao, J. Floral classification of honey using liquid chromatography–diode array detection–tandem mass spectrometry and chemometric analysis. *Food Chem.* **2014**, *145*, 941–949. [[CrossRef](#)] [[PubMed](#)]
25. Shapla, U.M.; Solyaman, M.; Alam, N.; Khalil, M.I.; Gan, S.H. 5-Hydroxymethylfurfural (HMF) levels in honey and other food products: Effects on bees and human health. *Chem. Cent. J.* **2018**, *12*, 1–18. [[CrossRef](#)] [[PubMed](#)]
26. Beretta, G.; Granata, P.; Ferrero, M.; Orioli, M.; Facino, R.M. Standardization of antioxidant properties of honey by a combination of spectrophotometric/fluorimetric assays and chemometrics. *Anal. Chim. Acta* **2005**, *533*, 185–191. [[CrossRef](#)]
27. Bertoneclj, J.; Doberšek, U.; Jamnik, M.; Golob, T. Evaluation of the phenolic content, antioxidant activity and color of Slovenian honey. *Food Chem.* **2007**, *105*, 822–828. [[CrossRef](#)]
28. Chua, L.S.; Rahaman, N.L.A.; Adnan, N.A.; Eddie Tan, T.T. Antioxidant activity of three honey samples in relation with their biochemical components. *J. Anal. Methods Chem.* **2013**, *2013*. [[CrossRef](#)]
29. Isla, M.I.; Craig, A.; Ordoñez, R.; Zampini, C.; Sayago, J.; Bedascarrasbure, E.; Alvarez, A.; Salomón, V.; Maldonado, L. Physicochemical and bioactive properties of honeys from Northwestern Argentina. *LWT—Food Sci. Technol.* **2011**, *44*, 1922–1930. [[CrossRef](#)]
30. Ruiz-Ciau, D.; Cuevas-Glory, L.; Quijano, L.; Sauri-Duch, E. Chemical Composition and Antioxidant DPPH Activity of the Floral and Leaves Essential Oils of *Montanoa speciosa* DC. *Am. J. Plant Sci.* **2017**, *8*, 745–753. [[CrossRef](#)]
31. Daulay, A.S.; Syahputra, R.A.; Nafitri, A. Antioxidant Activity Test of Chayote (*Sechium edule* (Jacq.) Swartz) Ethanol Extract using DPPH Method. *J. Phys. Conf. Ser.* **2021**, *1819*, 12035. [[CrossRef](#)]
32. Rahman, M.M.; Gan, S.H.; Khalil, M.I. Neurological effects of honey: Current and future prospects. In *Evidence-Based Complementary and Alternative Medicine*; Oxford University Press: Oxford, UK, 2014. [[CrossRef](#)]
33. Hegazi, A.G.; Guthami, F.M.A.; Gethami, A.F.M.A.; Fouad, E.A. Antibacterial and Antioxidant Activities of Some Saudi Arabia Honey Products. *Iran. J. Med. Microbiol.* **2020**, *14*, 490–500. [[CrossRef](#)]

34. Hegazi, A.G.; Guthami, F.M.; Al Gethami, A.F.M.; Fouad, E.A.; Abdou, A.M.; Hegazi, S.; Al Guthami, F.M.; Abdou, A.M. Antibacterial activity and characterisation of some Egyptian honey of different floral origin. *Bulg. J. Vet. Med.* **2021**, *24*, 278–290. [[CrossRef](#)]
35. Wei Se, K.; Abdul Wahab, R.; Nuratiqah Syed Yaacob, S.; Krishna Ghoshal, S. Study review Detection techniques for adulterants in honey: Challenges and recent trends. *J. Food Compos. Anal.* **2019**, *80*, 16–32. [[CrossRef](#)]
36. Roby, M.H.H.; Abdelaliem, Y.F.; Esmail, A.H.M.; Mohdaly, A.A.A.; Ramadan, M.F. Evaluation of Egyptian honeys and their floral origins: Phenolic compounds, antioxidant activities, and antimicrobial characteristics. *Environ. Sci. Pollut. Res.* **2020**, *27*, 20748–20756. [[CrossRef](#)]
37. Geană, E.I.; Ciucure, C.T.; Costinel, D.; Ionete, R.E. Evaluation of honey in terms of quality and authenticity based on the general physicochemical pattern, major sugar composition and $\delta^{13}\text{C}$ signature. *Food Control* **2020**, *109*, 106919. [[CrossRef](#)]
38. Salvador, L.; Guijarro, M.; Rubio, D.; Aucatoma, B.; Guillén, T.; Jentsch, P.V.; Ciobotă, V.; Stolker, L.; Ulic, S.; Vásquez, L.; et al. Exploratory Monitoring of the Quality and Authenticity of Commercial Honey in Ecuador. *Foods* **2019**, *8*, 105. [[CrossRef](#)]
39. El Sohaimy, S.A.; Masry, S.H.D.; Shehata, M.G. Physicochemical characteristics of honey from different origins. *Ann. Agric. Sci.* **2015**, *60*, 279–287. [[CrossRef](#)]
40. Aljohar, H.I.; Maher, H.M.; Albaqami, J.; Al-Mehaizie, M.; Orfali, R.; Alrubia, S. Physical and chemical screening of honey samples available in the Saudi market: An important aspect in the authentication process and quality assessment. *Saudi Pharm. J.* **2018**, *26*, 932–942. [[CrossRef](#)]
41. Szczesna, T.; Waś, E.; Semkiw, P.; Skubida, P.; Jaśkiewicz, K.; Witek, M. Changes in the Physicochemical Properties of Starch Syrups after Processing by Honeybees. *Agriculture* **2021**, *11*, 335. [[CrossRef](#)]
42. Kazemina, M.; Mahmoudi, R.; Aali, E.; Ghajarbygi, P. Evaluation of Authenticity in Honey Samples from Qazvin, Iran. *J. Chem. Health Risks* **2021**. [[CrossRef](#)]
43. Bentabol Manzanares, A.; Hernández García, Z.; Rodríguez Galdón, B.; Rodríguez, E.; Díaz Romero, C. Physicochemical characteristics of minor monofloral honeys from Tenerife, Spain. *LWT—Food Sci. Technol.* **2014**, *55*, 572–578. [[CrossRef](#)]
44. Pasiás, I.N.; Kiriakou, I.K.; Proestos, C. HMF and diastase activity in honeys: A fully validated approach and a chemometric analysis for identification of honey freshness and adulteration. *Food Chem.* **2017**, *229*, 425–431. [[CrossRef](#)] [[PubMed](#)]
45. Belay, A.; Haki, G.D.; Birringer, M.; Borck, H.; Lee, Y.C.; Kim, K.T.; Baye, K.; Melaku, S. Enzyme activity, amino acid profiles and hydroxymethylfurfural content in Ethiopian monofloral honey. *J. Food Sci. Technol.* **2017**, *54*, 2769. [[CrossRef](#)]
46. GSO 147:2008-Standards Store-GCC Standardization Organization. Gulf Technical Regulation. Available online: <https://www.gso.org.sa/store/standards/GSO:518550?lang=en> (accessed on 17 February 2022).
47. Warui, M.W.; Hansted, L.; Gikungu, M.; Mburu, J.; Kironchi, G.; Bosselmann, A.S. Characterization of Kenyan Honeys Based on Their Physicochemical Properties, Botanical and Geographical Origin. *Int. J. Food Sci.* **2019**, *2019*. [[CrossRef](#)]
48. Saxena, S.; Gautam, S.; Sharma, A. Physical, biochemical and antioxidant properties of some Indian honeys. *Food Chem.* **2010**, *118*, 391–397. [[CrossRef](#)]
49. Kaškonienė, V.; Maruska, A.; Kornysova, O.; Ligor, M.; Kaškonienė, V.; Maruška, A.; Kornysova, O.; Charczun, N.; Ligor, M.; Buszewski, B. Quantitative and qualitative determination of phenolic compounds in honey Plantarum project View project Impact of clear cuttings on transformation of biodiversity in forest ecosystems View project Quantitative and qualitative determination of phenolic compounds in honey. *Cheminé Technol.* **2009**, *3*, 74–80.
50. Meda, A.; Lamien, C.E.; Romito, M.; Millogo, J.; Nacoulma, O.G. Determination of the total phenolic, flavonoid and proline contents in Burkina Fasan honey, as well as their radical scavenging activity. *Food Chem.* **2005**, *91*, 571–577. [[CrossRef](#)]
51. Gonçalves, J.; Ribeiro, I.; Marçalo, J.; Rijo, P.; Faustino, C.; Pinheiro, L. Physicochemical, Antioxidant and Antimicrobial Properties of selected Portuguese Commercial Monofloral Honeys. *J. Food Nutr. Res.* **2018**, *6*, 645–654. [[CrossRef](#)]
52. Al, M.L.; Daniel, D.; Moise, A.; Bobis, O.; Laslo, L.; Bogdanov, S. Physico-chemical and bioactive properties of different floral origin honeys from Romania. *Food Chem.* **2009**, *112*, 863–867. [[CrossRef](#)]
53. Srečković, N.Z.; Mihailović, V.B.; Katanić Stanković, J.S. Physico-chemical, antioxidant and antimicrobial properties of three different types of honey from central Serbia. *Kragujev. J. Sci.* **2019**, *41*, 53–68. [[CrossRef](#)]
54. Perna, A.; Intaglietta, I.; Simonetti, A.; Gambacorta, E. A comparative study on phenolic profile, vitamin C content and antioxidant activity of Italian honeys of different botanical origin. *Int. J. Food Sci. Technol.* **2013**, *48*, 1899–1908. [[CrossRef](#)]
55. Ahmed, S.; Sulaiman, S.A.; Baig, A.A.; Ibrahim, M.; Liaqat, S.; Fatima, S.; Jabeen, S.; Shamim, N.; Othman, N.H. Honey as a Potential Natural Antioxidant Medicine: An Insight into Its Molecular Mechanisms of Action. *Oxidative Med. Cell. Longev.* **2018**, *2018*. [[CrossRef](#)] [[PubMed](#)]
56. Cianciosi, D.; Forbes-Hernández, T.Y.; Afrin, S.; Gasparrini, M.; Reboledo-Rodríguez, P.; Manna, P.P.; Zhang, J.; Lamas, L.B.; Flórez, S.M.; Toyos, P.A.; et al. Phenolic Compounds in Honey and Their Associated Health Benefits: A Review. *Molecules* **2018**, *23*, 2322. [[CrossRef](#)]
57. Alvarez-Suarez, J.M.; Tulipani, S.; Díaz, D.; Estevez, Y.; Romandini, S.; Giampieri, F.; Damiani, E.; Astolfi, P.; Bompadre, S.; Battino, M. Antioxidant and antimicrobial capacity of several monofloral Cuban honeys and their correlation with color, polyphenol content and other chemical compounds. *Food Chem. Toxicol.* **2010**, *48*, 2490–2499. [[CrossRef](#)]
58. Silici, S.; Sagdic, O.; Ekici, L. Total phenolic content, antiradical, antioxidant and antimicrobial activities of Rhododendron honeys. *Food Chem.* **2010**, *121*, 238–243. [[CrossRef](#)]

59. Ferreira, I.C.F.R.; Aires, E.; Barreira, J.C.M.; Estevinho, L.M. Antioxidant activity of Portuguese honey samples: Different contributions of the entire honey and phenolic extract. *Food Chem.* **2009**, *114*, 1438–1443. [[CrossRef](#)]
60. van den Berg, A.J.; van den Worm, E.; van Ufford, H.C.Q.; Halkes, S.B.; Hoekstra, M.J.; Beukelman, C.J. An in vitro examination of the antioxidant and anti-inflammatory properties of buckwheat honey. *J. Wound Care* **2008**, *17*, 172–178. [[CrossRef](#)]
61. Bueno-Costa, F.M.; Zambiasi, R.C.; Bohmer, B.W.; Chaves, F.C.; Silva, W.P.; da Zanusso, J.T.; Dutra, I. Antibacterial and antioxidant activity of honeys from the state of Rio Grande do Sul, Brazil. *LWT—Food Sci. Technol.* **2016**, *65*, 333–340. [[CrossRef](#)]
62. Boussaid, A.; Chouaibi, M.; Rezig, L.; Hellal, R.; Donsi, F.; Ferrari, G.; Hamdi, S. Physicochemical and bioactive properties of six honey samples from various floral origins from Tunisia. *Arab. J. Chem.* **2018**, *11*, 265–274. [[CrossRef](#)]
63. Pontis, J.A.; da Costa, L.A.M.A.; da Silva, S.J.R.; Flach, A. Color, phenolic and flavonoid content, and antioxidant activity of honey from Roraima, Brazil. *Food Sci. Technol.* **2014**, *34*, 69–73. [[CrossRef](#)]
64. Nayaka, N.M.D.M.W.; Fidrianny, I.; Sukrasno Hartati, R.; Singgih, M. Antioxidant and antibacterial activities of multiflora honey extracts from the Indonesian Apis cerana bee. *J. Taibah Univ. Med. Sci.* **2020**, *15*, 211–217. [[CrossRef](#)] [[PubMed](#)]
65. Stagos, D.; Soultisiotis, N.; Tsadila, C.; Papaeconomou, S.; Arvanitis, C.; Ntontos, A.; Karkanta, F.; Adamou-Androulaki, S.; Petrotos, K.; Spandidos, D.A.; et al. Antibacterial and antioxidant activity of different types of honey derived from Mount Olympus in Greece. *Int. J. Mol. Med.* **2018**, *42*, 726. [[CrossRef](#)] [[PubMed](#)]
66. Hegazi, A.G.; Abd Allah, F.M. Antimicrobial Activity of Different Saudi Arabia Honeys. *Global Veterinaria* **2012**, *9*, 53–59.
67. Al Masaudi, S. The antibacterial activities of honey. *Saudi J. Biol. Sci.* **2021**, *28*, 2188–2196. [[CrossRef](#)]
68. Mandal, M.D.; Mandal, S. Honey: Its medicinal property and antibacterial activity. *Asian Pac. J. Trop. Biomed.* **2011**, *1*, 154. [[CrossRef](#)]
69. Israili, Z.H. Antimicrobial properties of honey. *Am. J. Ther.* **2014**, *21*, 304–323. [[CrossRef](#)]
70. Libonatti, C.; Varela, S.; Basualdo, M. Antibacterial activity of honey: A review of honey around the world. *J. Microbiol. Antimicrob.* **2014**, *6*, 51–56. [[CrossRef](#)]
71. Szweda, P. Antimicrobial Activity of Honey. *Honey Anal.* **2017**, *3*, 15. [[CrossRef](#)]
72. Morroni, G.; Alvarez-Suarez, J.M.; Brenciani, A.; Simoni, S.; Fioriti, S.; Pugnaloni, A.; Giampieri, F.; Mazzoni, L.; Gasparrini, M.; Marini, E.; et al. Comparison of the antimicrobial activities of four honeys from three countries (New Zealand, Cuba, and Kenya). *Front. Microbiol.* **2018**, *9*, 1378. [[CrossRef](#)]
73. Irish, J.; Blair, S.; Carter, D.A. The antibacterial activity of honey derived from Australian flora. *PLoS ONE* **2011**, *6*, e18229. [[CrossRef](#)] [[PubMed](#)]
74. Mohapatra, D.P.; Thakur, V.; Brar, S.K. Antibacterial Efficacy of Raw and Processed Honey. *Biotechnol. Res. Int.* **2011**, *2011*, 1–6. [[CrossRef](#)] [[PubMed](#)]
75. Matzen, R.D.; Leth-Espensen, J.Z.; Jansson, T.; Nielsen, D.S.; Lund, M.N.; Matzen, S. The Antibacterial Effect in Vitro of Honey Derived from Various Danish Flora. *Dermatol. Res. Pract.* **2018**, *2018*. [[CrossRef](#)] [[PubMed](#)]
76. Molan, P.C. The antibacterial activity of honey 2. Variation in the potency of the antibacterial activity. *Bee World* **1992**, *73*, 59–76. [[CrossRef](#)]
77. Lee, H.; Churey, J.J.; Worobo, R.W. Antimicrobial activity of bacterial isolates from different floral sources of honey. *Int. J. Food Microbiol.* **2008**, *126*, 240–244. [[CrossRef](#)]
78. Anthimidou, E.; Mossialos, D. Antibacterial activity of Greek and Cypriot honeys against staphylococcus aureus and pseudomonas aeruginosa in comparison to manuka honey. *J. Med. Food* **2013**, *16*, 42–47. [[CrossRef](#)]
79. Kuś, P.M.; Szweda, P.; Jerković, I.; Tuberoso, C.I.G. Activity of Polish uniflora honeys against pathogenic bacteria and its correlation with colour, phenolic content, antioxidant capacity and other parameters. *Lett. Appl. Microbiol.* **2016**, *62*, 269–276. [[CrossRef](#)]
80. Leyva-Jiménez, F.J.; Lozano-Sánchez, J.; Borrás-Linares, I.; Cadiz-Gurrea, M.d.l.L.; Mahmoodi-Khaledi, E. Potential antimicrobial activity of honey phenolic compounds against Gram positive and Gram-negative bacteria. *LWT* **2019**, *101*, 236–245. [[CrossRef](#)]
81. Brudzynski, K. Effect of hydrogen peroxide on antibacterial activities of Canadian honeys. *Can. J. Microbiol.* **2006**, *52*, 1228–1237. [[CrossRef](#)]
82. White, J.W.; Subers, M.H.; Schepartz, A.I. The identification of inhibine, the antibacterial factor in honey, as hydrogen peroxide and its origin in a honey glucose-oxidase system. *Biochim. Biophys. Acta (BBA)—Spec. Sect. Enzymol. Subj.* **1963**, *73*, 57–70. [[CrossRef](#)]
83. Bhat, R.R.; Pai, M.v.; Ram, H.S.; Reddy, S.; Rai, S.; Thejeswi, P.; Prabhu, S. Comparison of Sugar and Honey Dressings in Healing of Chronic Wounds. *IOSR J. Dent. Med. Sci.* **2014**, *13*, 82–88. [[CrossRef](#)]