

Supplementary materials

Physicochemical and Antimicrobial Properties of Lactic Acid-Based Natural Deep Eutectic Solvents as a Function of Water Content

Olga N. Pozharitskaya ¹, Ekaterina D. Obluchinskaya ^{1,2}, Veronika A. Shikova ³, Elena V. Flisyuk ³, Evgeny V. Vishnyakov ⁴, Elena V. Makarevich ² and Alexander N. Shikov ^{1,3,*}

¹ Murmansk Marine Biological Institute of the Russian Academy of Sciences (MMBI RAS), Vladimirskaia Str., 17, 183038 Murmansk, Russia; olgapozhar@mail.ru (O.N.P.), okaterine@yandex.ru (E.D.O.)

² Department of Microbiology and Biochemistry, Murmansk Arctic University, Institute of Natural Sciences and Technology, Kapitana Egorova Str., 15, 183038 Murmansk, Russia; makarevichev@mstu.edu.ru

³ Department of Technology of Pharmaceutical Formulations, St. Petersburg State Chemical Pharmaceutical University, Prof. Popov Str., 14, 197376 Saint-Petersburg, Russia; veronika.shikova@spcpu.ru (V.A.S.); elena.flisyuk@pharminnotech.com (E.V.F.)

⁴ Department of Pharmaceutical Chemistry, St. Petersburg State Chemical Pharmaceutical University, Prof. Popov Str., 14, 197376 Saint-Petersburg, Russia; evgeniy.vishnyakov@pharminnotech.com

* Correspondence: spb.pharmacy@gmail.com




Figure S1. FTIR spectra of NADES1. Water content in NADES: 0% - black line; 10% - red line; 20% - green line; 30% - violet line; 40% - blue line.




Figure S2. FTIR spectra of NADES2. Water content in NADES: 0% - black line; 10% - red line; 20% - green line; 30% - violet line; 40% - blue line




Figure S3. FTIR spectra of NADES3. Water added in NADES: 0% - black line; 10% - red line; 20% - green line; 30% - violet line; 40% - blue line.