

**Table S5.** Prevalence of neuropsychiatric symptoms, anxiety, and depression by gender at each time point.

Measure	Baseline, N=97			Follow-Up, N=48		
	Men, N=26 <sup>1</sup>	Women, N=71 <sup>1</sup>	p-value	Men, N=16 <sup>1</sup>	Women, N=32 <sup>1</sup>	p-value
Since COVID, do you now have more difficulty...						
Remembering conversations a few days later?						
Yes	3 (14%)	22 (35%)	0.067 <sup>2</sup>	5 (38%)	9 (32%)	0.7 <sup>3</sup>
No	18 (86%)	40 (65%)		8 (62%)	19 (68%)	
Remembering placement of familiar objects?						
Yes	2 (10%)	22 (36%)	0.021 <sup>2</sup>	4 (27%)	10 (37%)	0.7 <sup>3</sup>
No	19 (90%)	39 (71%)		11 (73%)	17 (63%)	
Finding the right words when speaking?						
Yes	8 (40%)	22 (39%)	0.5 <sup>2</sup>	8 (50%)	20 (80%)	0.048 <sup>3</sup>
No	12 (60%)	35 (61%)		8 (50%)	5 (20%)	
Thinking clearly?						
Yes	6 (30%)	32 (39%)	0.5 <sup>2</sup>	5 (32%)	11 (42%)	0.5 <sup>3</sup>
No	14 (70%)	23 (61%)		11 (68%)	15 (58%)	
Staying Focused?						
Yes	7 (37%)	31 (51%)	0.3 <sup>2</sup>	9 (56%)	12 (43%)	0.4 <sup>2</sup>
No	12 (63%)	30 (49%)		7 (44%)	16 (57%)	
Since COVID, have you experienced...						
New or more frequent/severe headaches?						
Yes	2 (10%)	23 (34%)	0.022 <sup>2</sup>	3 (20%)	11 (38%)	0.3 <sup>3</sup>
No	18 (90%)	44 (66%)		12 (80%)	18 (62%)	
Weakness in one or more parts of your body?						
Yes	5 (22%)	18 (26%)	0.7 <sup>2</sup>	2 (14%)	7 (24%)	0.7 <sup>3</sup>
No	18 (78%)	50 (74%)		12 (86%)	22 (76%)	
Numbness or tingling?						
Yes	3 (13%)	15 (22%)	0.3 <sup>2</sup>	3 (20%)	9 (31%)	0.5 <sup>3</sup>
No	20 (87%)	53 (78%)		12 (80%)	20 (69%)	
Changes in vision?						
Yes	2 (10%)	17 (25%)	0.096 <sup>2</sup>	2 (14%)	5 (18%)	>0.9 <sup>3</sup>
No	21 (90%)	51 (75%)		12 (86%)	23 (82%)	
Sense of dizziness, imbalance, or vertigo?						
Yes	3 (13%)	23 (35%)	0.039 <sup>2</sup>	4 (25%)	8 (30%)	>0.9 <sup>3</sup>
No	21 (81%)	43 (65%)		12 (75%)	19 (70%)	
Seizures?						
Yes	0 (0%)	1 (2%)	0.5 <sup>2</sup>	0 (0%)	1 (4%)	>0.9 <sup>3</sup>
No	25 (100%)	66 (98%)		15 (100%)	27 (96%)	
Overall Median (IQR) Scores						
BAI	5 (1, 8)	4 (1, 9.75)	>0.9 <sup>4</sup>	5 (3, 2)	5 (1, 8.5)	0.7
PHQ-9	3 (1, 8)	4 (1, 9.75)	0.6 <sup>4</sup>	4.5 (2.75, 8.75)	3 (1, 6)	0.2

<sup>1</sup> n (%); median (IQR).

<sup>2</sup> Pearson's chi-squared test.

<sup>3</sup> Fisher's Exact test.

<sup>4</sup> Wilcoxon rank sum test.

\*total participants reported for each symptom will not equate to "N" due to some participants choosing: (1) to not reply; or (2) "unsure".