

Article

# Inhibitory Control under Threat: The Role of Spontaneous Eye Blinks in Post-Traumatic Stress Disorder

Mikael Rubin <sup>1,2</sup>, Denise A. Hien <sup>1,3</sup>, Dipanjana Das <sup>1,4</sup> and Robert D. Melara <sup>1,\*</sup>

<sup>1</sup> Department of Psychology, City College, City University of New York, New York, NY 10017, USA; mikael.r13@gmail.com (M.R.); dr.denise.hien@gmail.com (D.A.H.); ddas4425@gmail.com (D.D.)

<sup>2</sup> Department of Psychology, University of Texas at Austin, Austin, TX 78705, USA

<sup>3</sup> Derner Institute of Advanced Psychological Studies, Adelphi University, South Avenue, Garden City, NY 11530, USA

<sup>4</sup> Department of Psychological & Brain Sciences, University of California at Santa Barbara, Santa Barbara, CA 93106, USA

\* Correspondence: bobmelara@gmail.com

Academic Editor: Zhewu Wang

Received: 25 July 2016; Accepted: 24 January 2017; Published: 4 February 2017

**Abstract:** This study is the first to explore spontaneous eye blink rate (sEBR) in individuals with post-traumatic stress disorder (PTSD). We investigated the connection between the magnitude of flanker interference in PTSD participants and sEBR during performance on a modified version of the Eriksen flanker task. As a peripheral measure of cognitive control and dopaminergic function, sEBR may illuminate the relationship between PTSD and executive function. Findings revealed a positive relationship between sEBR and flanker interference in participants diagnosed with PTSD, to both threat-related and neutral stimuli, whereas this relationship was negative in participants exposed to trauma but without PTSD and in healthy controls. Although our results are suggestive of sEBR as a potential physiological index of emotional management in PTSD, most of the correlations were not significant, indicating that further research with a larger sample is needed.

**Keywords:** post-traumatic stress disorder; spontaneous eye blink rate; trauma exposure; flanker interference; cognitive control; inhibitory control

---

## 1. Introduction

Post-traumatic stress disorder (PTSD), affecting between 3.6% and 9.7% of the general population [1], is a traumatic stress-related disorder characterized by intrusive thoughts and memories of traumatic events, avoidance of thoughts and feelings associated with the trauma, emotional numbing, and an amplified vigilance and alertness to environmental triggers following exposure to a life-threatening stressor. Results of neuropsychological tests reveal compromise to executive control functions in PTSD patients, particularly working memory (ability to maintain recent experiences in memory) and inhibitory control (ability to suppress task-irrelevant memories and experiences) [2–5]. The aim of the current study was to investigate the relationship between working memory (as indexed by spontaneous eye blink rate; sEBR) and inhibitory control of attentional interference (as measured by behavioral performance on the temporal flanker paradigm) in (a) individuals with PTSD; (b) individuals with trauma exposure but no PTSD (TE); and (c) healthy controls (HCs) with neither trauma exposure nor PTSD. The relationships between sEBR and attentional interference in PTSD and TE participants may help clarify underlying attentional processes that are thought to result from trauma exposure.

### 1.1. Spontaneous Eye Blink Rate (sEBR)

Spontaneous eye blinks represent a broad index of information processing functions spanning attention and working memory and are distinct from both reflexive and voluntary eye blinks [6]. Neuroimaging analyses reveal the role of spontaneous eye blinks in modulating activity in oculomotor regions—including the frontal eye-fields and the supplementary eye-fields [7]—and also in higher-level cortical regions, including the parahippocampal gyrus and the medial frontal gyrus [8]. The parahippocampal gyrus is intimately involved in the updating of working memory to task-relevant stimuli [9–11], whereas the medial frontal cortex has been linked to executive control of task-irrelevant stimuli, most specifically during the monitoring and resolution of stimulus conflict in Stroop-like tasks [12]. Both working memory and executive control are compromised in PTSD [2–5]. sEBR also has been strongly associated with the dopaminergic system [13–15]: Pharmacological interventions reveal close ties between sEBR and D2 dopamine receptor functioning [13–15]. Researchers hence have used the eye blink rate as a non-invasive measure of dopamine functioning. Patients with disorders that involve hyperdopaminergic states—including schizophrenia, Tourette’s syndrome, and Huntington’s disease—show higher blink rates than healthy controls, whereas those with disorders that have hypodopaminergic states—such as Parkinson’s disease—show lower blink rates [16]. In non-disease states, sEBR serves as a gauge of cognitive load [14–16] independent of motor function.

In cognitive research, investigators have consistently found that high rates of spontaneous eye blinks during cognitive task performance are associated with enhanced perceptual sensitivity [17–20]. One speculative hypothesis is that sEBR gauges the buildup [21] or release [22] from working memory of task-related stimulus representations. In this view, enhanced perceptual sensitivity from high sEBRs might yield better inhibitory control of distractors once working memory is freed up [23]. This interpretation is supported by findings of reduced flanker [24] or Stroop [25] interference (i.e., greater inhibitory control of distractors) in healthy participants with greater sEBR. Inhibitory control entails increased resistance to distracting information during the performance of a cognitive task [26]. This includes both inhibitory action control and inhibitory interference control. Inhibitory control can be described by the ability to ignore the conflict between targets and distractors (flankers), by concentrating only on the target. In a standard flanker task this represents inhibitory interference control. Thus, decreased flanker interference indicates greater inhibitory control [27].

In the context of exposure to trauma, sEBR may offer insight into distinct cognitive strategies for handling threats, possibly leading to differential outcomes in inhibitory control to threats observed in PTSD and TE individuals. Functional neuroimaging of PTSD participants during the performance of cognitive control tasks reveals abnormal activity in the ventromedial prefrontal cortex, anterior cingulate cortex, amygdala, and insula [28–32]. Inefficient prefrontal control mechanisms, combined with hyperactive amygdala sensitivity, may result in the reduced inhibitory control to threats observed in PTSD [28,30,33].

The relatively greater sensitivity to threats seen in individuals with PTSD (e.g., [34]) may be expressed as greater monitoring of threat-related cues during task performance. Here, sEBR could gauge the accumulation of threat material in working memory (i.e., the filling up of working memory), which would effectively undermine the inhibitory control of conflict between targets and distractors in PTSD. In this case, higher sEBR would predict poorer inhibitory control (i.e., more interference) with exposure to task-irrelevant threatening stimuli. In contrast, the sEBR of TE individuals to threat-related stimuli may reflect the adaptive unburdening of emotional material from working memory (i.e., the emptying out of working memory), thus enabling greater cognitive control of stimulus conflict. In this case, higher sEBR would predict enhanced inhibitory control (i.e., less interference) with exposure to task-irrelevant threatening stimuli (i.e., threat suppression) in TE.

### 1.2. The Present Study

We addressed two gaps in the literature concerning the relationship between the control of working memory and the psychopathology of stress and PTSD. First, we evaluated in patients with

PTSD predictive associations between the monitoring or control of working memory, measured here as sEBR, and the magnitude of flanker interference, as an inverse measure of inhibitory control to distractors. Second, we evaluated whether traumatic stress exposure per se influences inhibitory control by examining whether participants with PTSD use control processes differently from those in each of the two comparison groups (TE and healthy controls).

The current study was part of a larger study to examine inhibitory control in PTSD [35]. We used a modified flanker conflict task [36] in which targets and distractors were separated temporally. Participants were asked to make judgments of line orientation while ignoring task-irrelevant lines and threatening or non-threatening images. We employed two sets of distracting images—a set of faces and a set of International Affective Picture System (IAPS) scenes—to probe the generality of task-irrelevant images in PTSD. We recorded behavioral performance and sEBRs across blocks of threatening or non-threatening trials. Although the relationship between sEBR and symptomology has been explored in other clinical populations, to the best of our knowledge our study is the first to examine the link between sEBR and inhibitory control in the context of PTSD.

## 2. Methods

### 2.1. Participants


Fifty individuals were recruited via online and print advertisements and paid to participate in the study. The nature of the procedures was explained fully, and informed consent was obtained from each participant; the Institutional Review Board of the City University of New York approved the protocol #298190. Following written consent, participants were given a breathalyzer/urine toxicology screen and completed a demographics survey and a life events checklist (LEC [37]). Trained clinicians administered the Structured Clinical Interview for the Diagnostic and Statistical Manual (SCID-IV [38]; DSM-IV). Inclusion criteria included: (1) physically healthy with no serious medical problems; (2) normal or corrected normal visual acuity; (3) aged 18–65; and (4) able to provide informed consent. Exclusion criteria included: (1) substance use disorder; (2) a major depressive episode; (3) suicidality; (4) current or history of psychosis or bipolar disorder, and (5) lack of fluency in English. Participants also were administered the Clinician-Administered PTSD Scale (CAPS [37]), and completed the Multiscale Dissociation Inventory (MDI [39]), the Test of Nonverbal Intelligence (TONI-4 [40]), and the Beck Depression Inventory<sup>®</sup>-II [41]. Three groups were created: (1) PTSD ( $n = 19$ , 10 males, mean age = 35.3), participants who met all DSM-IV-TR [42] criteria for PTSD; (2) TE ( $n = 16$ , eight males, mean age = 34.3), participants without PTSD or an Axis I diagnosis who had experienced a traumatic event meeting Criterion A for PTSD; and (3) HC ( $n = 15$ , five males, mean age = 33.4), participants without PTSD, an Axis I diagnosis, or Criterion A trauma exposure.

### 2.2. Stimulus, Apparatus, and Procedure

Each participant completed 24 blocks of experimental trials (and one or more blocks of practice trials) in a modified version of the visual flanker task [36] called the temporal flanker paradigm. The study was carried out in an electrically and acoustically shielded Industrial Acoustics Company (New York, NY, USA) chamber. Stimuli were created in Presentation<sup>®</sup> version 16.0 software (Neurobehavioral Systems, Berkeley, CA, USA) on a Dimension 5150 Dell desktop computer and appeared to participants as they sat at a distance of 60 cm from a 17-inch Dell Model P1130 RGB computer monitor with a refresh rate of 75 Hz. Each task in the paradigm comprised a block of 80 trials. Each trial consisted of a fixation square ( $0.67^\circ$ ) followed by three stimulus displays presented sequentially: (1) First Flanker; (2) Target; and (3) Second Flanker (see Figure 1). Each display appeared for 150 ms separated by an inter-stimulus interval varying between 153 ms and 390 ms in random distribution. The first and second flankers were identical on each trial: a vertical line, a horizontal line, or a cross. Each target was a vertical or horizontal line superimposed on an image, either a grayscale threatening (i.e., fearful) or non-threatening face from the NimStim set [43] or a colored threatening

or non-threatening picture from the International Affective Picture System (IAPS [44]). Line stimuli, subtending a visual angle of  $0.57^\circ$ , appeared in gray on a black background; face and IAPS stimuli subtended  $9.93^\circ$  (V)  $\times$   $7.13^\circ$  (H) and  $10.85^\circ$  (V)  $\times$   $9.74^\circ$  (H) of visual angle, respectively. We included two types of threat cues in the current study—faces and IAPS images—to evaluate the generality of the relationship between sEBR and flanker interference in PTSD when confronted with threatening versus non-threatening material.

On each trial of each task participants were asked to respond by mouse key as quickly and accurately as possible to the orientation of the target line, ignoring the flanker lines and the surrounding image. To mitigate eye movement artifacts during EEG recording participants were asked to blink only before or after, but not during, stimulus presentation, as needed. The duration of stimuli was 1265 ms on average within a trial that lasted approximately 3665 ms. Assignment of line orientation to response keys was counterbalanced across participants. Tasks were divided into baseline tasks (distractor constant) and filtering tasks (distractor random) [45]; for purposes of the present study, only performance on filtering tasks was analyzed. In each filtering task distractors appeared randomly on each trial: On 40% of trials (32 of 80) target and flanker lines matched in orientation (congruent trials), on 40% they mismatched (incongruent trials), and on 20% (16 of 80) flankers were crosses (neutral trials). All eight images from one of the four sets (threat faces, non-threat faces, threat IAPS, non-threat IAPS) appeared randomly an equal number of times with the target line, creating four different filtering tasks. Each task set was repeated three times in a session; task order was balanced across participants. The entire experiment, including EEG preparation, lasted approximately 3 h.



**Figure 1.** The modified flanker task. This example is of a fearful face incongruent trial.

### 2.3. Data Recording and Analysis

Reaction times (RTs) to correct response trials of each participant were averaged for each of two trial types (i.e., congruent, incongruent) in each of four conditions (i.e., threat faces, non-threat faces, threat IAPS, non-threat IAPS). Our measure of inhibitory control (i.e., flanker interference) was defined as the difference between congruent and incongruent trials in each condition.

This study was part of a larger EEG study using a BioSemi Active-Two system in a high-density (160 electrodes) montage arranged in an elastic cap with blinks and other eye movements monitored by electrooculogram (EOG) at a sampling rate of 512 Hz from two electrode montages, one on the infra- and supra-orbital ridges of the right eye (VEOG), the other on the outer canthi of each eye (HEOG) (see [35]). For purposes of the present study, only VEOG recordings were analyzed. An automated Matlab function called Peakfinder [46] was used to identify and count blinks in the VEOG channel. Peakfinder implements a user-defined magnitude threshold to compare temporally adjacent peaks within a noisy unidimensional vector. Blinks were defined as peaks with a magnitude greater than  $100 \mu\text{V}$  [47]. Blink frequency was summed across the three repetitions of each condition and divided

by condition duration (in minutes) to determine average blink rate per minute (EBR) for each of the four conditions. The automated blink detector was checked for accuracy by two trained raters independently scoring a random selection of 30 blocks. No blinks were detected at a threshold lower than 100  $\mu$ V (no false positives). The raters revealed 98% agreement in identifying blinks; the manual counts were 92% similar to the automated ones.

Two-tailed correlations were carried out within each of the three groups (i.e., PTSD, TE, HC) to examine the relationship between sEBR and inhibitory control in each condition. Additionally, we conducted exploratory analyses to examine whether specific clinical domains are related to sEBR. Correlational analyses were performed on clinical measures by separately regressing each subscale of the CAPS, BDI, and MDI on sEBR in the PTSD and TE groups. Fisher r-to-z transformations were performed to evaluate differences in the magnitude of correlations between groups ( $p < 0.05$  for all tests).

### 3. Results

#### 3.1. Demographic, Clinical, Physiological, and Behavioral Data

Table 1 contains a tally and statistical analysis of demographic and clinical data. PTSD and TE showed significantly higher rates of trauma than HC. PTSD revealed higher scores on the BDI than TE or HC. The three groups did not differ significantly in gender, age, education or race/ethnicity. The three groups did not differ significantly in sEBR or flanker interference.

**Table 1.** Demographic, clinical, physiological, and behavioral information and comparisons among participants.

|                             | PTSD ( $n = 19$ ) | TE ( $n = 16$ ) | HC ( $n = 15$ ) | Test Statistic             | $p$  |
|-----------------------------|-------------------|-----------------|-----------------|----------------------------|------|
| Gender                      |                   |                 |                 |                            |      |
| Male                        | 10                | 8               | 5               | $\chi^2(2, N = 50) = 1.38$ | 0.50 |
| Female                      | 9                 | 8               | 10              |                            |      |
| Age M (SD)                  | 35.3 (10.1)       | 34.3 (12.0)     | 33.4 (9.7)      | $F(2, 47) = 0.129$         | 0.88 |
| Education (years) M (SD)    | 14.7 (1.9)        | 14.3 (1.8)      | 14.1 (4.7)      | $F(2, 47) = 0.151$         | 0.86 |
| Race/Ethnicity              |                   |                 |                 |                            |      |
| White                       | 3                 | 3               | 3               | $\chi^2(2, N = 50) = 2.61$ | 0.27 |
| Black                       | 8                 | 5               | 6               |                            |      |
| Latino                      | 6                 | 3               | 4               |                            |      |
| Asian                       | 1                 | 4               | 1               |                            |      |
| Other                       | 1                 | 1               | 1               |                            |      |
| Trauma type (LEC)           |                   |                 |                 |                            |      |
| Physical Abuse/Assault      | 16                | 11              | 3               | $\chi^2(2, N = 50) = 5.93$ | 0.00 |
| Sexual Trauma               | 17                | 11              | 1               |                            |      |
| Natural Disaster            | 6                 | 6               | 7               |                            |      |
| War/Combat                  | 2                 | 2               | 2               |                            |      |
| Other                       | 36                | 44              | 24              |                            |      |
| BDI                         | 11.3 (7.9)        | 3.8 (5.4)       | 0.47 (0.92)     | $F(2, 47) = 15.8$          | 0.00 |
| Blink rate (mean) M (SD)    |                   |                 |                 |                            |      |
| Non-threat faces            | 15.9 (8.5)        | 20.8 (9.2)      | 19.4 (15.4)     | $F(2, 47) = 0.898$         | 0.41 |
| Threat faces                | 15.2 (9.1)        | 20.1 (8.5)      | 19.6 (12.2)     | $F(2, 47) = 1.292$         | 0.28 |
| Non-threat IAPS             | 17.3 (9.4)        | 19.4 (11.3)     | 17.6 (11.1)     | $F(2, 47) = 0.200$         | 0.82 |
| Threat IAPS                 | 16.3 (8.4)        | 17.5 (12.2)     | 17.3 (10.1)     | $F(2, 47) = 0.066$         | 0.94 |
| Overall                     | 16.3 (8.4)        | 19.2 (9.5)      | 18.2 (11.3)     | $F(2, 47) = 0.408$         | 0.67 |
| Flanker interference M (SD) |                   |                 |                 |                            |      |
| Non-threat faces            | 112.1 (100.0)     | 133.1 (102.0)   | 111.8 (89.8)    | $F(2, 47) = 0.923$         | 0.41 |
| Threat faces                | 98.7 (86.3)       | 114.9 (110.4)   | 119.2 (108.3)   | $F(2, 47) = 0.999$         | 0.38 |
| Non-threat IAPS             | 119.4 (97.3)      | 142.5 (105.9)   | 97.8 (63.9)     | $F(2, 47) = 0.200$         | 0.82 |
| Threat IAPS                 | 94.0 (83.7)       | 140.2 (129.7)   | 109.2 (69.2)    | $F(2, 47) = 0.254$         | 0.78 |
| Overall                     | 106.0 (84.2)      | 132.7 (104.0)   | 109.4 (78.2)    | $F(2, 47) = 0.436$         | 0.65 |

PTSD = Post-traumatic Stress Disorder, TE = Trauma Exposed, HC = Healthy Controls, LEC = Life Events Checklist, BDI = Beck Depression Inventory, IAPS = International Affective Picture System.

### 3.2. Inhibitory Control

Table 2 summarizes correlation coefficients between sEBR and flanker interference scores (incongruent RT minus congruent RT) for each group in each of the four conditions. A pattern of correlations emerged in which sEBR and flanker interference consistently correlated positively in the PTSD group and consistently correlated negatively in the TE and HC groups. At an overall level (across conditions), the correlation coefficients were not statistically significant in two-tailed tests:  $r = 0.43$ ,  $p = 0.067$  (PTSD);  $r = -0.25$ ,  $p = 0.37$  (TE);  $r = -0.28$ ,  $p = 0.30$  (HC). However, there was a significant difference in the correlation coefficients between the PTSD and TE groups,  $z = -1.98$ ,  $p < 0.05$ . Moreover, the PTSD group showed a significant positive association between sEBR and flanker interference in the threat IAPS condition,  $r = 0.51$ ,  $p < 0.05$ , which differed significantly from the (nonsignificant) negative correlations between sEBR and flanker interference in the TE ( $r = -0.20$ ;  $z = -2.29$ ,  $p < 0.05$ ) and HC ( $r = -0.29$ ;  $z = -2.01$ ,  $p < 0.05$ ) groups.

**Table 2.** Correlation coefficients between sEBR and inhibitory control scores (incongruent RT minus congruent RT) for PTSD, TE, and healthy control participants in each of four conditions: threat faces, non-threat faces, threat IAPS, and non-threat IAPS.

| Interference Condition | PTSD Blink Rate ( $n = 19$ ) |       |                    |                    | TE Blink Rate ( $n = 16$ ) |        |        |        | HC Blink Rate ( $n = 15$ ) |        |        |        |
|------------------------|------------------------------|-------|--------------------|--------------------|----------------------------|--------|--------|--------|----------------------------|--------|--------|--------|
|                        | NF                           | TF    | NI                 | TI                 | NF                         | TF     | NI     | TI     | NF                         | TF     | NI     | TI     |
| Non-threat Faces       | 0.363                        |       |                    |                    | -0.225                     |        |        |        | -0.295                     |        |        |        |
| Threat Faces           |                              | 0.284 |                    |                    |                            | -0.201 |        |        |                            | -0.196 |        |        |
| Non-threat IAPS        |                              |       | 0.415 <sup>†</sup> |                    |                            |        | -0.219 |        |                            |        | -0.158 |        |
| Threat IAPS            |                              |       |                    | 0.510 <sup>*</sup> |                            |        |        | -0.201 |                            |        |        | -0.286 |

PTSD = Post-traumatic Stress Disorder, TE = Trauma Exposed, HC = Healthy Controls, NF = Non-threat Faces, TF = Threat Faces, NI = Non-threat IAPS, TI = Threat IAPS; <sup>†</sup>  $p < 0.1$ , <sup>\*</sup>  $p < 0.05$ .

### 3.3. Clinical Subscales

Exploratory two-tailed correlations were carried out in the PTSD and TE groups between sEBR and clinical subscales of the CAPS (four subscales) and MDI (six subscales), and the overall BDI. The blink rate correlated negatively only with the emotional constriction subscale of the MDI in the PTSD group,  $r = -0.48$ ,  $p < 0.05$ . (The five items from the emotional constriction subscale are: “Not being able to feel emotions”; “Feeling frozen inside without feelings”; “Knowing you should be mad or sad about something, but not having any feelings”; “Not having emotions or feelings at a time when you should have been upset”; “Knowing you must be upset, but not being able to feel it”).

## 4. Discussion

The current study investigated links between the magnitude of inhibitory control in PTSD participants and their rate of eye blinks during performance of a temporal flanker task. We found that sEBR in participants diagnosed with PTSD was associated positively with flanker interference: Participants evinced a higher blink rate as the difference in RT between congruent and incongruent stimuli increased in the face of both threatening and non-threatening IAPS images. Thus, higher sEBR was associated with poorer attentional performance in PTSD. In contrast, in healthy control participants, as well as in participants who had been exposed during their lifetime to traumatic stressors but did not have PTSD, there was no significant association between blink rates and inhibitory control; at most, they exhibited a trend towards the usual relationship of sEBR with better attentional performance. Importantly, the correlations between PTSD and TE differed significantly from each other, suggesting that sEBR may help elucidate differences in underlying cognitive processes between the two groups.

In the PTSD group the IAPS stimuli yielded numerically higher correlation coefficients than the face stimuli (see Table 2). Interestingly, the difference between stimulus sets was particularly robust to threat stimuli. The effect cannot be understood merely as a weaker response in PTSD to facial processing because faces also appeared in many of the IAPS images. Perhaps IAPS images are more potent stimuli for eliciting threat because, in depicting threatening scenes, they are more directly

threatening to PTSD participants, whereas, in expressing fear, NimStim faces depict only the aftermath of (i.e., the response to) threat. We believe this tentative explanation merits further investigation (see Shin and Liberzon [48]). In any case, we conclude that in our cue reactivity paradigm, the PTSD participants only weakly experienced the fearful faces as trauma cues compared with the IAPS images.

Hester and Garavan [23] hypothesized that sEBR predicts the success of cognitive processes responsible for inhibiting the influence of task-irrelevant stimuli on performance. At high rates of spontaneous blinking, the contents of working memory are dispersed [22], thus enabling frontal executive mechanisms to exert top-down control over distractors. Here, blinks are linked closely to cognitive control [24,25]. This perspective fits with our findings in the TE and HC groups. The negative correlations between sEBR and flanker interference in these groups suggest that greater sEBR is related to greater cognitive control. We speculate that there may be an emotional component to this process: Individuals in these two groups may be able to release threatening and neutral material equally well from working memory, enabling executive control of stimulus conflict across all incongruent trials.

A different understanding of the role of sEBR is desirable in explaining the performance of our participants with PTSD. Optimal levels of dopamine during the performance of conflict tasks [49] correspond to the peak of an inverted U-shaped function [50]. PTSD has been linked to abnormally elevated release of presynaptic dopamine [51,52] in both the amygdala [50] and dorsal anterior cingulate cortex (dACC) [53], implying inefficient inhibitory control of hypersensitive limbic regions [54]. To the extent that sEBR gauges hyperactivation in these regions, the positive correlation of the blink rate with flanker interference found in the current study may represent poor overall inhibitory control in PTSD.

What cognitive or clinical functions might be served by a positive correlation between sEBR and interference in PTSD? It is conceivable that sEBR acts as a check on the flow of information entering working memory [21]. A diagnostic feature of PTSD is heightened sensitivity to environmental threat [55,56]. On this account, blinks in individuals with PTSD may serve as a proxy for the threat monitoring processes that these patients engage in automatically to complete everyday tasks [57]. In particular, sEBR in this population could gauge the buildup of threat material within working memory, a process at cross-purposes with the cognitive control needed to ameliorate stimulus conflict. sEBR in those with PTSD may reflect the output of a hypersensitive emotion management system, rather than a cognitive control system. Emotional management involves both emotion monitoring, observed here in the positive correlation in PTSD between sEBR and flanker interference, and emotion control, observed in our exploratory findings indicating a negative correlation in PTSD between sEBR and the dissociative symptom of emotional constriction. Importantly, we found correlations of nearly equal magnitude to both threat and non-threat images. Responsivity under conditions of low threat in PTSD has also been reported elsewhere [58,59]. An explanation favored by previous investigators, and in concert with our emotional management account, is that in assessing risk patients with PTSD, vigilantly monitor both safety and danger. Still, at present our explanations are speculative and our results are in need of replication.

## 5. Conclusions

The current study is the first to explore sEBR in individuals with PTSD. We showed that whereas increasing rates of spontaneous eye blinks are associated negatively with flanker interference in TE and HC, sEBR and flanker interference are actually associated positively in PTSD. One implication is that sEBR represents the engagement of differential functions for individuals with and without PTSD, enabling either improved cognitive control (TE) or more effective emotion management (PTSD). To probe these differential uses, in future research sEBR could serve as the dependent measure in manipulations of working memory load with threatening or non-threatening material in participants from PTSD and TE populations.

**Acknowledgments:** The study was supported by a City Seeds grant from the City College, City University of New York (PI: Denise A. Hien).

**Author Contributions:** Robert D. Melara had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Study concept and design: Denise A. Hien, Robert D. Melara; Acquisition, analysis, or interpretation of data: Mikael Rubin, Dipanjana Das; Drafting of the manuscript: Mikael Rubin, Robert D. Melara; Critical revision of the manuscript for important intellectual content: Robert D. Melara, Denise A. Hien, Dipanjana Das; Statistical analysis: Mikael Rubin; Obtained funding: Denise A. Hien; Administrative, technical, or material support: Denise A. Hien, Robert D. Melara; Study supervision: Denise A. Hien, Robert D. Melara.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Disclaimer:** The views and opinions expressed in this report are those of the authors and should not be construed to represent the views of sponsoring organizations, agencies, or the U.S. government.

## References

1. Kessler, R.C.; Chiu, W.T.; Demler, O.; Merikangas, K.R.; Walters, E.E. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch. Gen. Psychiatry* **2005**, *62*, 617–627. [[CrossRef](#)] [[PubMed](#)]
2. Harrington, K.M.; Miller, M.W.; Wolf, E.J.; Reardon, A.F.; Ryabchenko, K.A.; Ofrat, S. Attention-deficit/hyperactivity disorder comorbidity in a sample of veterans with posttraumatic stress disorder. *Compr. Psychiatry* **2012**, *53*, 679–690. [[CrossRef](#)] [[PubMed](#)]
3. Leskin, L.P.; White, P.M. Attentional networks reveal executive function deficits in posttraumatic stress disorder. *Neuropsychology* **2007**, *21*, 275–284. [[CrossRef](#)] [[PubMed](#)]
4. Swick, D.; Honzel, N.; Larsen, J.; Ashley, V.; Justus, T. Impaired Response Inhibition in Veterans with Post-Traumatic Stress Disorder and Mild Traumatic Brain Injury. *J. Int. Neuropsychol. Soc.* **2012**, *18*, 917–926. [[CrossRef](#)] [[PubMed](#)]
5. Clark, C.R.; Mcfarlane, A.C.; Morris, P.; Weber, D.L.; Sonkilla, C.; Shaw, M.; Marcina, J.; Egan, G.F. Cerebral Function in Posttraumatic Stress Disorder during Verbal Working Memory Updating: A Positron Emission Tomography Study. *Biol. Psychiatry* **2003**, *53*, 474–481. [[CrossRef](#)]
6. Walrath, L.C.; Goldstein, R.; Stern, J.A. The Endogenous Eyeblink. *Psychophysiology* **1984**, *21*, 22–34.
7. Bristow, D.; Haynes, J.-D.; Sylvester, R.; Frith, C.D.; Rees, G. Blinking suppresses the neural response to unchanging retinal stimulation. *Curr. Biol.* **2005**, *15*, 1296–1300. [[CrossRef](#)] [[PubMed](#)]
8. Yoon, H.W.; Chung, J.-Y.; Song, M.-S.; Park, H. Neural correlates of eye blinking; improved by simultaneous fMRI and EOG measurement. *Neurosci. Lett.* **2005**, *381*, 26–30. [[CrossRef](#)] [[PubMed](#)]
9. Halgren, E.; Baudena, P.; Clarke, J.M.; Heit, G.; Liégeois, C.; Chauvel, P.; Musolino, A. Intracerebral potentials to rare target and distractor auditory and visual stimuli. I. Superior temporal plane and parietal lobe. *Electroencephalogr. Clin. Neurophysiol.* **1995**, *94*, 191–220. [[CrossRef](#)]
10. Knight, R.T.; Scabini, D.; Woods, D.L.; Clayworth, C.C. Contributions of temporal-parietal junction to the human auditory P3. *Brain Res.* **1989**, *502*, 109–116. [[CrossRef](#)]
11. Halgren, E.; Marinkovic, K.; Chauvel, P. Generators of the late cognitive potentials in auditory and visual oddball tasks. *Electroencephalogr. Clin. Neurophysiol.* **1998**, *106*, 156–164. [[CrossRef](#)]
12. Ridderinkhof, K.R.; Ullsperger, M.; Crone, E.A.; Nieuwenhuis, S. The role of the medial frontal cortex in cognitive control. *Science* **2004**, *306*, 443–447. [[CrossRef](#)] [[PubMed](#)]
13. Karson, C.N.; Kleinman, J.E.; Berman, K.F.; Phelps, B.H. An inverse correlation between spontaneous eye-blink rate and platelet monoamine oxidase activity. *Br. J. Psychiatry* **1983**, *142*, 43–46. [[CrossRef](#)] [[PubMed](#)]
14. Lawrence, M.S.; Redmond, D.E. MPTP lesions and dopaminergic drugs alter eye blink rate in African green monkeys. *Pharmacol. Biochem. Behav.* **1991**, *38*, 869–874. [[CrossRef](#)]
15. Elsworth, J.D.; Lawrence, M.S.; Roth, R.H.; Taylor, J.R.; Mailman, R.B.; Nichols, D.E.; Lewis, M.H.; Redmond, D.E., Jr. D1 and D2 dopamine receptors independently regulate spontaneous blink rate in the vervet monkey. *J. Pharmacol. Exp. Ther.* **1991**, *259*, 595–600. [[PubMed](#)]
16. Caplan, R.; Caplan, R.; Guthrie, D.; Guthrie, D.; Komo, S.; Komo, S. Blink rate in children with attention-deficit-hyperactivity disorder. *Biol. Psychiatry* **1996**, *39*, 1032–1038. [[CrossRef](#)]
17. Pivik, R.T.; Dykman, R.A. Endogenous eye blinks in preadolescents: Relationship to information processing and performance. *Biol. Psychol.* **2004**, *66*, 191–219. [[CrossRef](#)] [[PubMed](#)]



18. Tanaka, Y.; Yamaoka, K. Blink activity and task difficulty. *Percept. Mot. Skills* **1993**, *77*, 55–66. [[CrossRef](#)] [[PubMed](#)]
19. Goldstein, R.; Walrath, L.C.; Stern, J.A. Blink activity in a discrimination task as a function of stimulus modality and schedule of presentation. *Psychophysiology* **1985**, *22*, 629–635. [[CrossRef](#)] [[PubMed](#)]
20. Ohira, H. Eyeblink activity in a word-naming task as a function of semantic priming and cognitive load. *Percept. Mot. Skills* **1996**, *82*, 835–842. [[CrossRef](#)] [[PubMed](#)]
21. Siegle, G.J.; Ichikawa, N.; Steinhauer, S. Blink before and after you think: Blinks occur prior to and following cognitive load indexed by pupillary responses. *Psychophysiology* **2008**, *45*, 679–687. [[CrossRef](#)] [[PubMed](#)]
22. Ichikawa, N.; Ohira, H. Eyeblink activity as an index of cognitive processing: Temporal distribution of eyeblinks as an indicator of expectancy in semantic priming. *Percept. Motor Skills* **2004**, *98*, 131–140. [[CrossRef](#)] [[PubMed](#)]
23. Hester, R.; Garavan, H. Working memory and executive function: The influence of content and load on the control of attention. *Mem. Cogn.* **2005**, *33*, 221–233. [[CrossRef](#)]
24. Van Bochove, M.E.; Van der Haegen, L.; Notebaert, W.; Verguts, T. Blinking predicts enhanced cognitive control. *Cogn. Affect. Behav. Neurosci.* **2013**, *13*, 346–354. [[CrossRef](#)] [[PubMed](#)]
25. Zhang, T.; Mou, D.; Wang, C.; Tan, F.; Jiang, Y.; Lijun, Z.; Li, H. Dopamine and executive function: Increased spontaneous eye blink rates correlate with better set-shifting and inhibition, but poorer updating. *Int. J. Psychophysiol.* **2015**, *96*, 155–161. [[CrossRef](#)] [[PubMed](#)]
26. Melara, R.D.; Algom, D. Driven by information: A tectonic theory of Stroop effects. *Psychol. Rev.* **2003**, *110*, 422–471. [[CrossRef](#)] [[PubMed](#)]
27. Melara, R.D.; Rao, A.; Tong, Y. The duality of selection: Excitatory and inhibitory processes in auditory selective attention. *J. Exp. Psychol. Hum. Percept. Perform.* **2002**, *28*, 279–306. [[CrossRef](#)] [[PubMed](#)]
28. Dickie, E.W.; Brunet, A.; Akerib, V.; Armony, J.L. Neural correlates of recovery from post-traumatic stress disorder: A longitudinal fMRI investigation of memory encoding. *Neuropsychologia* **2011**, *49*, 1771–1778. [[CrossRef](#)] [[PubMed](#)]
29. Dickie, E.W.; Brunet, A.; Akerib, V.; Armony, J.L. An fMRI investigation of memory encoding in PTSD: Influence of symptom severity. *Neuropsychologia* **2008**, *46*, 1522–1531. [[CrossRef](#)] [[PubMed](#)]
30. Mazza, M.; Tempesta, D.; Pino, M.C.; Catalucci, A.; Gallucci, M.; Ferrara, M. Regional cerebral changes and functional connectivity during the observation of negative emotional stimuli in subjects with post-traumatic stress disorder. *Eur. Arch. Psychiatry Clin. Neurosci.* **2013**, *263*, 575–583. [[CrossRef](#)] [[PubMed](#)]
31. Garfinkel, S.N.; Garfinkel, S.N.; Liberzon, I.; Liberzon, I. Neurobiology of PTSD: A Review of Neuroimaging Findings. *Psychiatr. Ann.* **2009**, *39*, 370–381. [[CrossRef](#)]
32. Koenigs, M.; Huey, E.D.; Raymond, V.; Cheon, B.; Solomon, J.; Wassermann, E.M.; Grafman, J. Focal brain damage protects against post-traumatic stress disorder in combat veterans. *Nat. Neurosci.* **2008**, *11*, 232–237. [[CrossRef](#)] [[PubMed](#)]
33. Robinson, O.J.; Charney, D.R.; Overstreet, C.; Vytal, K.; Grillon, C. The adaptive threat bias in anxiety: Amygdala-dorsomedial prefrontal cortex coupling and aversive amplification. *NeuroImage* **2012**, *60*, 523–529. [[CrossRef](#)] [[PubMed](#)]
34. Liberzon, I.; Sripada, C.S. The functional neuroanatomy of PTSD: A critical review. *Prog. Brain Res.* **2008**, *167*, 151–169. [[PubMed](#)]
35. Melara, R.D.; Ruglass, L.; Fertuck, E.; Hien, D.A. Self-Regulation of Threat in Post-traumatic Stress Disorder: Associations between Inhibitory Control and Dissociative Symptomatology. *BMC Public Health* **2017**, in preparation.
36. Eriksen, B.A.; Eriksen, C.W. Effects of noise letters upon the identification of a target letter in a nonsearch task. *Percept. Psychophys.* **1974**, *16*, 143–149. [[CrossRef](#)]
37. Blake, D.D.; Weathers, F.W.; Nagy, L.M.; Kaloupek, D.G.; Gusman, F.D.; Charney, D.S.; Keane, T.M. The development of a Clinician-Administered PTSD Scale. *J. Traum. Stress* **1995**, *8*, 75–90. [[CrossRef](#)]
38. First, M.B.; Gibbon, M. *The Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I) and the Structured Clinical Interview for DSM-IV Axis II Disorders (SCID-II)*; John Wiley & Sons Inc.: Hoboken, NJ, USA, 2004.
39. Briere, J. *Multiscale Dissociation Inventory (MDI)*; Psychological Assessment Resources: Odessa, FL, USA, 2002.
40. Brown, L.; Sherbenou, R.J.; Johnsen, S.K. *Test of Non-Verbal Intelligence (TONI-3)*, 3rd ed.; Pro-Ed: Austin, TX, USA, 1997.

41. Beck, A.T.; Steer, R.A.; Brown, G.K. *Manual for the Beck Depression Inventory-II*; Psychological Corporation: San Antonio, TX, USA, 1996.
42. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR Fourth Edition (Text Revision)*, 4th ed.; American Psychiatric Association: Washington, DC, USA, 2000.
43. Tottenham, N.; Tanaka, J.W.; Leon, A.C.; McCarry, T.; Nurse, M.; Hare, T.A.; Marcus, D.J.; Westerlund, A.; Casey, B.J.; Nelson, C. The NimStim set of facial expressions: Judgments from untrained research participants. *Psychiatry Res.* **2009**, *168*, 242–249. [[CrossRef](#)] [[PubMed](#)]
44. Lang, P.J.; Bradley, M.M.; Cuthbert, B.N. International Affective Picture System (IAPS): Technical Manual and Affective Ratings. Available online: [http://www2.unifesp.br/dpsicobio/Nova\\_versao\\_pagina\\_psicobio/adap/instructions.pdf](http://www2.unifesp.br/dpsicobio/Nova_versao_pagina_psicobio/adap/instructions.pdf) (accessed on 25 January 2017).
45. Garner, W.R. *The Processing of Information and Structure*; Erlbaum: Potomac, MD, USA, 1974.
46. Yoder, N. PeakFinder. Available online: <https://www.mathworks.com/matlabcentral/fileexchange/25500-peakfinder-x0--sel--thresh--extrema--includeendpoints--interpolate-?requestedDomain=www.mathworks.com> (accessed on 25 January 2017).
47. Barbato, G.; della Monica, C.; Costanzo, A.; De Padova, V. Dopamine activation in Neuroticism as measured by spontaneous eye blink rate. *Physiol. Behav.* **2012**, *105*, 332–336. [[CrossRef](#)] [[PubMed](#)]
48. Shin, L.M.; Liberzon, I. The Neurocircuitry of Fear, Stress, and Anxiety Disorders. *Neuropsychopharmacology* **2010**, *35*, 169–191. [[CrossRef](#)] [[PubMed](#)]
49. Badgaiyan, R.D.; Wack, D.; Wack, D. Evidence of Dopaminergic Processing of Executive Inhibition. *PLoS ONE* **2011**, *6*, e28075. [[CrossRef](#)] [[PubMed](#)]
50. Cools, R.; D'Esposito, M. Inverted-U-Shaped Dopamine Actions on Human Working Memory and Cognitive Control. *Biol. Psychiatry* **2011**, *69*, e113–e125. [[CrossRef](#)] [[PubMed](#)]
51. Hamner, M.B.; Diamond, B.I. Elevated plasma dopamine in posttraumatic stress disorder: A preliminary report. *Biol. Psychiatry* **1993**, *33*, 304–306. [[CrossRef](#)]
52. Glover, D.A.; Powers, M.B.; Bergman, L.; Smits, J.A.J.; Telch, M.J.; Stuber, M. Urinary dopamine and turn bias in traumatized women with and without PTSD symptoms. *Behav. Brain Res.* **2003**, *144*, 137–141. [[CrossRef](#)]
53. Thomaes, K.; Dorrepaal, E.; Draijer, N.; de Ruiter, M.B.; Elzinga, B.M.; van Balkom, A.J.; Smit, J.H.; Veltman, D.J. Treatment effects on insular and anterior cingulate cortex activation during classic and emotional Stroop interference in child abuse-related complex post-traumatic stress disorder. *Psychol. Med.* **2012**, *42*, 2337–2349. [[CrossRef](#)] [[PubMed](#)]
54. Assadi, S.M.; Yücel, M.; Pantelis, C. Dopamine modulates neural networks involved in effort-based decision-making. *Neurosci. Biobehav. Rev.* **2009**, *33*, 383–393. [[CrossRef](#)] [[PubMed](#)]
55. Cisler, J.M.; Wolitzky-Taylor, K.B.; Adams, T.G., Jr.; Babson, K.A.; Badour, C.L.; Willems, J.L. The emotional Stroop task and posttraumatic stress disorder: A meta-analysis. *Clin. Psychol. Rev.* **2011**, *31*, 817–828. [[CrossRef](#)] [[PubMed](#)]
56. Mogg, K.; Bradley, B.; Miles, F.; Dixon, R. BRIEF REPORT Time course of attentional bias for threat scenes: Testing the vigilance-avoidance hypothesis. *Cogn. Emot.* **2004**, *18*, 689–700. [[CrossRef](#)]
57. Lanius, R.A.; Frewen, P.A.; Vermetten, E.; Yehuda, R. Fear conditioning and early life vulnerabilities: Two distinct pathways of emotional dysregulation and brain dysfunction in PTSD. *Eur. J. Psychotraumatol.* **2010**, *1*, 585. [[CrossRef](#)] [[PubMed](#)]
58. Grillon, C.; Morgan, C. Fear-Potentiated Startle Conditioning to Explicit and Contextual Cues in Gulf War Veterans With Posttraumatic Stress Disorder. *J. Abnorm. Psychol.* **1999**, *108*, 134–142. [[CrossRef](#)] [[PubMed](#)]
59. Pole, N.; Neylan, T.C.; Best, S.R.; Orr, S.P.; Orr, S.P.; Marmar, C.R.; Marmar, C.R. Fear-potentiated startle and posttraumatic stress symptoms in urban police officers. *J. Traum. Stress* **2003**, *16*, 471–479. [[CrossRef](#)] [[PubMed](#)]

