

Table S2: Mean, standard error (SE), median, interquartile range (IQR) and *p* value of the reproductive parameters and litter characteristics and comparison between the groups.

Reproductive indicator	Group	N	Mean	SE	Median	IQR	<i>p</i> value
Total born	T1	16	16.56	0.86	16.50	13.00- 20.00	0.95
	T2	16	15.88	0.38	15.50	15.00- 17.00	
	T3	16	16.56	0.79	15.50	15.00- 17.50	
Born alive	T1	16	15.62	0.67	16.50	13.00- 18.00	0.99
	T2	16	15.43	0.32	15.00	14.50- 16.50	
	T3	16	15.88	0.64	15.00	14.00- 16.50	
Stillborn	T1	16	2.13	0.38	2.00 ^a	1.00- 3.00	0.03
	T2	16	0.94	0.27	1.00 ^b	0.00- 1.50	
	T3	16	1.06	0.23	1.00 ^b	0.50- 1.00	
Mummies	T1	16	0.75	0.23	0.50	0.00- 1.00	0.97
	T2	16	0.56	0.13	1.00	0.00- 1.00	
	T3	16	0.63	0.18	0.50	0.00- 1.00	
Alive >24h	T1	16	13.50	0.48	13.00 ^b	12.00- 15.00	0.04
	T2	16	14.94	0.31	15.00 ^a	14.00- 15.00	
	T3	16	15.19	0.59	14.00 ^a	14.00- 16.00	
Weaning	T1	16	12.56	0.24	13.00 ^b	12.00- 13.00	<0.01
	T2	16	14.19	0.14	14.00 ^a	14.00- 14.50	
	T3	16	14.06	0.21	14.00 ^a	14.00- 15.00	
Mean BW of weaning	T1	16	7.46	0.13	7.50	6.90- 7.90	0.32
	T2	16	7.64	0.16	7.50	7.20- 8.20	
	T3	16	7.75	0.12	7.65	7.35- 8.20	
Weaning to oestrus interval	T1	16	6.19	0.23	6.00 ^a	5.00- 7.00	0.03
	T2	16	5.63	0.15	6.00 ^{a,b}	5.00- 6.00	
	T3	16	5.38	0.18	5.00 ^b	5.00- 6.00	

*Figures with different superscripts are indicative of a statistically significant difference (*p* value < 0.05).