

Innovative Cosmeceutical Ingredients: Harnessing Selenosugar-Linked Hydroxycinnamic Acids for Antioxidant and Wound-Healing Properties

Giovanna Cimmino ^{1,2}, Mauro De Nisco ³, Simona Piccolella ^{1,*}, Claudia Gravina ¹, Silvana Pedatella ² and Severina Pacifico ¹


¹ Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "Luigi Vanvitelli", Via Vivaldi 43, 81100 Caserta, Italy; giovanna.cimmino@unicampania.it (G.C.); claudia.gravina@unicampania.it (C.G.); severina.pacifico@unicampania.it (S.P.)

² Department of Chemical Sciences, University of Napoli Federico II, Via Cinthia 4, 80126 Napoli, Italy; silvana.pedatella@unina.it

³ Department of Sciences, University of Basilicata, Via dell'Ateneo Lucano 10, 85100 Potenza, Italy; mauro.denisco@unibas.it

* Correspondence: simona.piccolella@unicampania.it

(A)



(B)




Figure S1. ¹H-NMR (A) and ¹³C-NMR (B) spectra of compound **3a** in CDCl₃.




Figure S2. ^1H -NMR (A) and ^{13}C -NMR (B) spectra of compound **3b** in CD_3OD .