

Supplementary Material

Effect of Alkylresorcinols Isolated from Wheat Bran on the Oxidative Stability of Minced-Meat Models as Related to Storage

Carolina Cantele ¹, Giulia Potenziani ¹, Ambra Bonciolini ¹, Marta Bertolino ^{1,†} and Vladimiro Cardenia ^{1,2,*,†}


¹ Department of Agricultural, Forest and Food Sciences (DISAFA), University of Turin,
10095 Grugliasco, TO, Italy; carolina.cantele@unito.it (C.C.); giulia.potenziani@unito.it (G.P.);
ambra.bonciolini@unito.it (A.B.); marta.bertolino@unito.it (M.B.)

² AgriForFood Chromatography and Mass Spectrometry Open Access Laboratory, University of Turin,
10095 Grugliasco, TO, Italy


* Correspondence: vladimiro.cardenia@unito.it

† These authors contributed equally to this work.


Figure S1. Molecular structure of alkylresorcinols.




AR15:0




AR17:0




AR19:0




AR21:0



AR23:0




AR25:0




AR27:0

Figure S2. GC/MS spectra of the alkylresorcinols (TMS derivatives) extracted from wheat bran.


AR15:0 TMS derivative




AR17:0 TMS derivative




AR19:0 TMS derivative




AR21:0 TMS derivative



AR23:0 TMS derivative



AR25:0 TMS derivative



AR27:0 TMS derivative

