

Supplemental files

Immune Responses and Protection Profiles in Mice Induced by Subunit Vaccine Candidates Based on the Extracellular Domain Antigen of Respiratory Syncytial Virus G Protein Combined with Different Adjuvants

Ruiwen Han ¹, Tangqi Wang ^{1†}, Xueting Cheng ¹, Jialuo Bing ², Jia Li ³, Yao Deng ¹, Xuchang Shan ², Xuejie Zhang ², Donghong Wang ¹, Shucaï Sun ⁴ and Wenjie Tan ^{1,2,3,4*}

¹ Key Laboratory of Biosafety, National Health Commissions, National Institute for Viral Disease Control and Prevention, China CDC, 155 Changbai Road, Beijing 102206, China; 18267839995@163.com


² School of Public Health, Xinxiang Medical University, Xinxiang 453003, China

³ Zhejiang Provincial Key Laboratory of Medical Genetics, School of Laboratory Medicine and Life Sciences, Wenzhou Medical University, Wenzhou 325035, China


² Department of Nuclear Medicine, The Second Hospital of Hebei Medical University, Shijiazhuang 050000, China

* Correspondence: tanwj@ivdc.chinacdc.cn

† These authors contributed equally to this work.



Supplementary Figure S1. Detection of intracellular cytokines IFN- γ , IL-2, TNF, and IL-4 in CD8⁺T cells from Gecto-immunized mice.



Supplementary Figure S2. Gating strategy for detection of CD4⁺ T cells intracellular cytokines. Exemplary gating strategy of Splenocytes sample from an G_{ect}o-immunized mouse. Splenocytes were collected 14 days later and the expression of IL-2, IFN- γ , TNF, and IL-4 in CD4⁺T cells.