

Supplementary Table 2. Microbiology of Infective Endocarditis in liver transplant recipients. Values show cases among patients with available data.

Characteristic	Value
Gram-positives, n (%)	43/62 (69.4%)
<i>Enterococcus</i> , n (%)	16/62 (25.8%)
<i>Enterococcus faecalis</i> , n (%)	9/16 (56.3%)
<i>Enterococcus faecium</i> , n (%)	5/16 (31.3%)
<i>Staphylococcus aureus</i> , n (%)	14/62 (22.6%)
Coagulase-negative <i>Staphylococcus</i> , n (%)	8/62 (12.9%)
<i>Streptococcus</i> spp., n (%)	2/62 (3.2%)
Gram-negatives, n (%)	6/62 (9.7%)
<i>Klebsiella</i> spp., n (%)*	3/62 (4.8%)
<i>Escherichia coli</i> , n (%)*	2/62 (3.2%)
<i>Proteus mirabilis</i> , n (%)	1/62 (1.6%)
<i>Enterobacter</i> spp., n (%)	1/62 (1.6%)
Fungi, n (%)	16/62 (25.8%)
<i>Aspergillus</i> spp., n (%)	13/62 (21%)
<i>Candida albicans</i> , n (%)	1/62 (1.6%)
<i>Pseudallescheria boydii</i> , n (%)	1/62 (1.6%)
<i>Phaeoacremonium parasiticum</i> , n (%)	1/62 (1.6%)
Polymicrobial, n (%)	9/62 (14.5%)
Culture negative, n (%)	5/54 (9.3%)

* One patient had a polymicrobial infection by *Klebsiella* spp. and *E. coli*.