

Table S1. Statistics of survival analyses in the group of NET patients.

rs8005354					
	Mean age of patients at time of admission [years ± std.dev]	P value	Median survival time [years]	HR (95% CI)	Log-rank p value
Without minor allele	63.7 ± 11.8	0.9926	12	1.24 (0.60–2.56)	0.5822
Minor allele carriers	63.7 ± 11.4		13		
rs2069762					
	Mean age of patients at time of admission [years ± std.dev]	P value	Median survival time [years]	HR (95% CI)	Log-rank p value
Without minor allele	64.9 ± 10.6	0.3329	12	0.93 (0.48–1.83)	0.8381
Minor allele carriers	62.9 ± 11.9		12		
rs3731198					
	Mean age of patients at time of admission [years ± std.dev]	P value	Median survival time [years]	HR (95% CI)	Log-rank p value
Without minor allele	63.6 ± 11.0	0.4330	13	0.86 (0.43–1.74)	0.6550
Minor allele carriers	65.5 ± 12.4		12		
rs1800872					
	Mean age of patients at time of admission [years ± std.dev]	P value	Median survival time [years]	HR (95% CI)	Log-rank p value
Without minor allele	63.0 ± 11.0	0.6052	12	1.41 (0.69–2.86)	0.3310
Minor allele carriers	64.1 ± 11.9		13		

Table S2. Comparison of the risk of locoregional or distant metastasis presence depending on the presence of the rarer allele (to confirm statistical significance expected p value after the implementation of the Bonferroni correction <0.0125).

variant	p value
rs8005354; T>C	p=0.80637
rs2069762; A>C	p=0.04369
rs3731198; T>C	p=0.07644
rs1800872; G>T	P=0.54011