

Supplementary Table S1: Most frequent symptoms and main biological parameters of Residents With COVID-19 During 11 Nursing Home Outbreaks in Sevilla, Spain

Clinical Features	Global (N = 587)	Patients Diagnosed With COVID-19 Before MP (N = 295)	Patients Diagnosed With COVID-19 During MP (N = 292)	p
Mean (SD) / Median [Q1-Q3] / No. (%)				
Most frequent symptoms				
Fatigue and global deterioration	229 (39%)	125 (42.5%)	104 (35.6%)	0.087
Dyspnea	203 (34.6%)	128 (43.5%)	75 (25.8%)	<0.001
Cough	167 (28.4%)	85 (29%)	82 (28.1%)	0.804
Low grade fever (37-37.9° C)	158 (27%)	91 (31%)	67 (22.9%)	0.063
Anorexia	129 (22%)	62 (21.1%)	67 (23%)	0.572
Delirium	108 (18.4%)	62 (21.2%)	46 (15.8%)	0.092
Diarrhea	76 (12.9%)	36 (12.2%)	40 (13.7%)	0.600
High grade fever (≥38° C)	73 (12.4%)	38 (12.9%)	35 (12.1%)	0.051
Nausea/Vomiting	42 (7.2%) / 32 (5.5%)	20 (6.8%) / 13 (4.4%)	22 (7.5%) / 19 (6.5%)	0.731/0.267
Rhinorrhea	21 (3.6%)	13 (4.4%)	8 (2.7%)	0.273
Odynophagia	19 (3.2%)	11 (3.7%)	8 (2.7%)	0.494
Falls	19 (3.2%)	10 (3.4%)	9 (3.1%)	0.827
Headache	18 (3.1%)	8 (2.7%)	10 (3.4%)	0.622
Ageusia	14 (2.4%)	11 (3.9%)	3 (1.1%)	0.031
Anosmia	10 (1.7%)	8 (2.8%)	2 (0.7%)	0.056
Asymptomatic	173 (29.5%)	73 (24.7%)	100 (34.2%)	0.012
Main biological parameters				
Hemoglobin (g/dL)	12.2 (2.2)	12.3 (2.3)	12.0 (2.1)	0.955
Leukocytes (no./µL)	7582 (4181)	8431 (4532)	6849 (3716)	0.003
Lymphocytes (no./µL)	1271 (684)	1189 (691)	1343 (673)	0.071
Platelets (no./µL)	231 341 (103 320)	236 757 (112 042)	226 705 (95 391)	0.443
D Dimer	2571 (6292)	2638 (5833)	2515 (6675)	0.886
Creatinine (mg/dL)	1.14 (1.0)	1.23 (1.0)	1.06 (1.0)	0.187
ASAT	30 (24.9)	34 (27)	27 (22)	0.051
ALAT	23 (21.8)	27 (28)	20 (14)	0.021
Creatinine kinase	135 (323)	158 (250)	117 (374)	0.329
CRP	64 (88)	83 (92)	48 (81)	0.002
Ferritin (ng/mL)	430 (623)	597 (809)	294 (363)	<0.001




Figure S1: Distribution of residents in all the nursing homes.