



Article

Do Surgeons Anticipate Women's Hopes and Fears Associated with Prolapse Repair? A Qualitative Analysis in the PROSPERE Trial

Xavier Fritel ^{1,*} , Marion Ravit ², Anne-Cécile Pizzoferrato ¹, Sandrine Campagne-Loiseau ³, Georges Bader ⁴, Perrine Capmas ^{5,6}, Michel Cosson ⁷, Philippe Debodinance ⁸, Xavier Deffieux ⁹, Hervé Fernandez ^{5,6}, Philippe Ferry ¹⁰, Olivier Garbin ¹¹, Bernard Jacquetin ³, Guillaume Legendre ¹² , Christian Saussine ¹³, Renaud de Tayrac ¹⁴ , Laurent Wagner ¹⁵, Jean-Philippe Lucot ¹⁶, Arnaud Fauconnier ¹⁷ and the PROSPERE team [†]

- ¹ Faculté de Médecine et Pharmacie, Université de Poitiers, INSERM CIC 1402, Service de Gynécologie-Obstétrique et Médecine de la Reproduction, CHU de Poitiers, 86000 Poitiers, France
 - ² Emergency Obstetric and Quality of Care Unit, Liverpool School of Tropical Medicine, Liverpool L3 5QA, UK
 - ³ Service de Gynécologie, CHU Estaing, 63000 Clermont-Ferrand, France
 - ⁴ Medical and Surgical Center Ambroise Paré, Hartmann, Pierre Cherest, 92200 Neuilly-sur-Seine, France
 - ⁵ INSERM Centre of research in Epidemiology and population health (CESP), U1018, 94276 Villejuif, France
 - ⁶ Service de Gynécologie-Obstétrique, Hôpital Bicêtre, GHU Sud, AP-HP, 94270 Le Kremlin-Bicêtre, France
 - ⁷ Faculté de Médecine, Université de Lille, Service de Gynécologie, CHRU Jeanne de Flandre, 59000 Lille, France
 - ⁸ Dunkerque Hospital Center, 59240 Dunkerque, France
 - ⁹ Faculté de Médecine, Université Paris-Sud, Service de Gynécologie-Obstétrique, Hôpital Antoine-Béclère APHP, 92140 Clamart, France
 - ¹⁰ La Rochelle Hospital Center, 17000 La Rochelle, France
 - ¹¹ Faculté de Médecine, Université de Strasbourg, Service de Gynécologie, CHRU de Strasbourg, 67000 Strasbourg, France
 - ¹² Faculté de Médecine, Université d'Angers, UMR S1085, Service de Gynécologie-Obstétrique, CHU d'Angers, 49000 Angers, France
 - ¹³ Faculté de Médecine, Université de Strasbourg, Service d'Urologie, CHRU de Strasbourg, 67000 Strasbourg, France
 - ¹⁴ Faculté de Médecine, Université de Montpellier, Service de Gynécologie-Obstétrique, CHU Carémeau, 30000 Nîmes, France
 - ¹⁵ Service d'Urologie, CHU Carémeau, 30000 Nîmes, France
 - ¹⁶ Saint-Vincent-de-Paul Hospital Center, 59000 Lille, France
 - ¹⁷ Faculté de Médecine, Université Paris-Saclay, Service de Gynécologie-Obstétrique, CHI Poissy-Saint-Germain-en-Laye, 78000 Poissy, France
- * Correspondence: xavier.fritel@univ-poitiers.fr; Tel.: +33-549-443-360
† See Acknowledgements.



Citation: Fritel, X.; Ravit, M.; Pizzoferrato, A.-C.; Campagne-Loiseau, S.; Bader, G.; Capmas, P.; Cosson, M.; Debodinance, P.; Deffieux, X.; Fernandez, H.; et al. Do Surgeons Anticipate Women's Hopes and Fears Associated with Prolapse Repair? A Qualitative Analysis in the PROSPERE Trial. *J. Clin. Med.* **2023**, *12*, 1332. <https://doi.org/10.3390/jcm12041332>

Academic Editor: Emilio Sacco

Received: 10 October 2022

Revised: 31 January 2023

Accepted: 2 February 2023

Published: 7 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Women's preoperative perceptions of pelvic-floor disorders may differ from those of their physicians. Our objective was to specify women's hopes and fears before cystocele repair, and to compare them to those that surgeons anticipate. We performed a secondary qualitative analysis of data from the PROSPERE trial. Among the 265 women included, 98% reported at least one hope and 86% one fear before surgery. Sixteen surgeons also completed the free expectations-questionnaire as a typical patient would. Women's hopes covered seven themes, and women's fears eleven. Women's hopes were concerning prolapse repair (60%), improvement of urinary function (39%), capacity for physical activities (28%), sexual function (27%), well-being (25%), and end of pain or heaviness (19%). Women's fears were concerning prolapse relapse (38%), perioperative concerns (28%), urinary disorders (26%), pain (19%), sexual problems (10%), and physical impairment (6%). Surgeons anticipated typical hopes and fears which were very similar to those the majority of women reported. However, only 60% of the women reported prolapse repair as an expectation. Women's expectations appear reasonable and consistent with the scientific literature on the improvement and the risk of relapse or complication related to cystocele repair. Our analysis encourages surgeons to consider individual woman's expectations before pelvic-floor repair.

Keywords: pelvic organ prolapse; surgery; expectations; hope; fear

1. Introduction

Pelvic organ prolapse (POP) is a frequent but not life-threatening disorder. It is an important cause of morbidity, and has a well-documented negative impact on quality of life (QoL) [1]. Some women affected by this condition may choose surgical repair, while others may not. A woman's lifetime risk of prolapse surgery is estimated at 19% [2].

Little is known about why women choose surgical treatment and their expectations of surgical repair. The severity of symptoms probably contributes to seeking this surgery [3]. Data on women's fears before prolapse surgery are even sparser. Complications reported with the use of mesh may induce specific fears.

Women's preoperative perceptions of the problems associated with this disorder appear to differ from those of their physicians [4]. It may be that women have greater expectations—higher hopes—for the results of surgical repair. We do not know the extent to which surgeons may anticipate women's expectations.

Surgeons habitually assess the results of their interventions based on the extent of the anatomical correction [5], the improvement of symptoms such as protrusion, and functional urinary or rectal signs [6]. More and more authors also analyze the results of prolapse surgery with a QoL scale. Nonetheless, these scales do not include women's expectations of the surgery—expectations themselves influenced by their past experiences. Understanding and identifying these expectations better before the intervention could enable an increase in women's satisfaction.

Today, when surgery with mesh implants is increasingly called into question and access to health information massively facilitated by the diversity and variability of its sources, it is necessary to tailor the management, information, and advice provided as part of shared medical decision making, which is a major advance in the management of functional disorders [7].

Our aim was to perform a secondary qualitative analysis of expectations (hopes and fears) reported by women before surgery scheduled for pelvic organ prolapse (POP) and included in the PROSPERE (PROSthetic PELvic floor REpair) [8] randomized-controlled trial, and to compare the hopes and fears reported by the women to those that surgeons anticipate their patients will have.

2. Materials and Methods

The PROSPERE multicenter randomized trial was designed to compare complications associated with cystocele repair with mesh by laparoscopy versus transvaginally [8]. The study of women's hopes and fears before surgery was planned as an ancillary study before the trial began.

2.1. Inclusion

From October 2012 through April 2014, participating surgeons invited patients aged 45 to 75 years who had a symptomatic prolapse of the anterior vaginal wall stage 2 or higher (according to the Pelvic Organ Prolapse Quantification (POP-Q) system) to participate in this study. Exclusion criteria were previous surgical POP repair, the impossibility of, or a contraindication to, either surgical route, pelvic-organ cancer, contraindication to the use of mesh, inability to read French, lack of health insurance, pregnancy, or a desire to become pregnant in the future.

The study-information leaflet given to each woman before requesting her written consent explained that complications were the principal outcome measure of the PROSPERE study—both intraoperative and postoperative, during the year after surgery. The intervention principles were described. The document listed as possible complications of the surgery its potential failure, vaginal, gastrointestinal, or bladder exposure of the mesh

implant, bladder injury, infection, intestinal injury, hemorrhage, chronic pain, dyspareunia, constipation, occlusion, incontinence, or urinary symptoms.

2.2. Questionnaires

Before surgery, we asked each woman included to report 5 hopes and 5 fears (ranked in descending order of importance) that she had concerning the planned operation [9]. Women who mentioned at least one hope or one fear on the inclusion questionnaire were included in our analysis. We also asked the surgeons (blinded to the women’s responses) involved in the RCT to report and rank what they presumed would be the preoperative hopes and fears of a typical patient.

2.3. Data Analysis

Categorization of women’s hopes and fears was based on qualitative analysis of their responses to the questionnaire. The initial development of themes began with the analysis of the first responses; this thematic classification was progressively revised and completed as we read and reread (XF and MR) all the questionnaires received. For this first step of the thematic elaboration, similar answers worded similarly were brought together in the same theme; on the other hand, we took care to respect the distinction made by each woman between different themes of hopes and themes of fears. Women’s responses were reread until the point of data saturation (no more emerging themes) [9]. A last reading enabled us to classify each participant’s responses according to the classification developed in the previous stage.

The hopes and fears expected by the surgeons were categorized with the same thematic classification developed for the women.

The Comité de Protection des Personnes Nord-Ouest IV (Institutional Review Board, Lille, France) and the Agence Nationale de Sécurité du Médicament et des Produits de Santé (the French National Agency for Medicines and Health Products Safety, #B111368-30) approved the PROSPERE study, which was registered at ClinicalTrials.gov (#NCT01637441). All patients provided written informed consent.

3. Results

Among the 265 women included in the PROSPERE trial, 261 (98%) reported at least one hope (mean, 2.3 hopes; 72 women reported only one hope, 79 two, 70 three, 35 four, and 5 five), and 229 (86%) at least one fear (mean, 1.5 fears; 114 women reported only one fear, 77 two, 20 three, 12 four, and 6 five). Among the 39 surgeons participating in the PROSPERE trial, 16 (41%) answered the expectations questionnaire from the viewpoint of a typical patient scheduled for prolapse repair. Characteristics of the 265 women are reported in Table 1.

Table 1. Description of the population before cystocele repair.

Women Characteristics		N = 265
Age	median (range)	64 (45–75)
Number of deliveries	median (range)	2 (0–16)
POP-Q Stage—no. of patients (%)	2	58 (21.9%)
	3	185 (69.8%)
	4	21 (7.9%)
PFDI 20	means (±SD)	94.32 (±50.67)
	POPDI-6	43.81 (±22.24)
	DDI-8	16.25 (±16.18)
	UDI-6	34.25 (±25.98)
EQ5D	means (±SD)	0.78 (±0.20)

The main fears ranked first by women concerned a prolapse relapse (28%), concerns about the perioperative period (21%), and urinary disorders (17%) (see Table 4 and Figure 4). The leading fears ranked second were again prolapse relapse (23%) and the perioperative period (17%), as well as pain (17%).

Table 2. Hope and Fears thematic classification for women and surgeons (examples of responses).

Hope Themes	Women	Surgeons
Prolapse repair	"no longer bulge out of my vagina", "do not see this ball anymore", "that my bladder is put back in place", "stop having to push up my uterus", "avoid this organ descent sensation", "do not push back the ball in the vagina"	"no more vaginal ball"
Improved urinary function	"no more leakage", "not have to go urinate every 5 min", "fewer runs to the bathroom", "urinate normally"	"reduction of urgency-pollakiuria", "urinate easily"
Capacity for physical activities	"to return to normal physical activity", "to walk and move normally", "to regain my mobility"	"recovery of physical activity"
Improved sexual function	"resume my sex life", "regain my femininity"	"more harmonious sexuality"
Well-being or comfort	"feel better", "have a normal life", "not have this embarrassment anymore"	"feel better"
End of pain or heaviness	"no more feeling of heaviness", "no more pain in the lower abdomen"	"no longer feeling of heaviness"
Improved bowel function	"move my bowels correctly", "no longer sitting on the toilet waiting for a bowel movement"	"move my bowels easily"
No more pads or other expectation	"not wearing pads anymore", "no more irritation or burn", "that the operation goes well"	-
Fears themes		
Failure or prolapse relapse	"fear of relapse", "afraid that the net does not hold"	"risk of recurrence", "failure"
Perioperative concerns	"fear of anesthesia", "afraid of not waking up", "afraid of nausea on waking", "postoperative problem", "surgical complications"	"complication during the operation"
Urinary disorder	"being incontinent", "continuing to smell the odor of pee!"	"risk of urinary leakage"
Pain	"fear of pain"	"fear of pain"
Sexual issues	"decreased sexual desire", "won't recover sexual desire"	"pain during sex"
Other fear	"that my belly will be even bigger", "infection", "health disorder", "gynecologic problem", "sequelae"	"infection"
Physical impairment	"Fear of not being able to resume my physical activities", "stay physically diminished"	"duration of convalescence"
Mesh-related complications	"mesh exposure", "fear the implant will be rejected"	"Serious adverse effect due to the mesh"
Worsened condition	"have to pay attention to everything", "being worse after than before", "fear of not recovering my general fairly satisfactory condition"	"worse than before"
Bowel problems	"constipation", "afraid of more severe constipation"	"Improvement in anorectal disorders"
No fears	"I'm not afraid, he knows what to do", "I am confident that this operation will make me more comfortable"	-

Overall, 87 (38%) women reported a fear of prolapse relapse, 64 (28%) a fear of the perioperative period, 63 (26%) a fear of a urinary disorder, and 40 women (17%) no

particular fear. The main fear among women worried about the perioperative period ($n = 64$) concerned anesthesia ($n = 40$).

The surgeons anticipated hopes and fears very similar to those the women reported (Tables 3 and 4).

Table 3. Preoperative hopes reported by women and by surgeons (for a “typical” patient).

Sample	Women (w)/Surgeons (s)						
	1st N _w /N _s n _w /n _s	2nd N _w /N _s n _w /n _s	3rd N _w /N _s n _w /n _s	4th N _w /N _s n _w /n _s	5th N _w /N _s n _w /n _s	Overall * Women N _w = 261 n _w (%)	Overall * Surgeons N _s = 16 n _s (%)
Hopes							
Prolapse repair	132/11	20/1	9/1	1/.	./.	156 (60)	13 (81)
Improved urinary function	37/.	58/6	21/4	6/6	./.	103 (39)	15 (94)
Capacity for physical activities	21/.	36/4	23/3	11/2	3/3	74 (28)	11 (69)
Improved sexual function	13/.	30/1	29/4	4/2	1/2	70 (27)	9 (56)
Well-being or comfort	36/5	18/2	16/1	7/1	2/2	66 (25)	10 (62)
End of pain or heaviness	17/.	19/2	7/2	7/1	./1	49 (19)	5 (31)
Improved bowel function	3/.	7/.	4/1	3/3	./3	16 (6)	7 (44)
No more pads or other	1/.	1/.	2/.	./1	./.	3 (1)	1 (6)

* Redundant answers in the same theme were not counted; as it was possible to declare several hopes or fears, the sum of the percentages is greater than 100%. Women’s responses in bold, N_w/n_w for number of women responding (N_w) in each theme (n_w), N_s/n_s for surgeons.

Table 4. Preoperative fears reported by women and surgeon (for a “typical” patient).

Sample	Women (w)/Surgeons (s)						
	1st N _w /N _s n _w /n _s	2nd N _w /N _s n _w /n _s	3rd N _w /N _s n _w /n _s	4th N _w /N _s n _w /n _s	5th N _w /N _s n _w /n _s	Summary * Women N _w = 229 n _w (%)	Summary * Surgeons N _s = 16 n _s (%)
Fears							
Failure or prolapse relapse	63/8	27/2	4/1	./2	2/3	87 (38)	15 (94)
Perioperative concerns	49/5	20/5	7/4	3/.	1/.	64 (28)	14 (87)
Urinary disorders	40/1	14/4	7/8	./2	./.	60 (26)	14 (87)
Pain	16/1	20/3	5/2	4/4	./4	44 (19)	10 (62)
Sexual problems	7/1	10/.	2/.	2/3	1/3	22 (10)	7 (44)
Other fear	2/.	7/.	7/1	6/1	2/2	17 (7)	3 (19)
Physical impairment	2/.	7/.	4/.	./1	./1	13 (6)	1 (6)
Mesh-related	5/.	4/1	1/.	1/1	./.	11 (5)	2 (12)
Worsened condition	4/.	2/1	1/.	1/.	./.	7 (3)	2 (12)
Bowel problems	1/.	3/.	./.	1/1	./1	5 (2)	1 (6)
No fears	40/.	1/.	./.	./.	./.	40 (17)	.

* Redundant answers in the same domain were not counted; as it was possible to declare several hopes or fears, the sum of the percentages is greater than 100%. Women’s responses in bold, N_w/n_w for number of women responding (N_w) in each theme (n_w), N_s/n_s for surgeons.

The surgeons’ responses about the preoperative hopes and fears of a typical patient were very similar to the women’s responses (Tables 3 and 4). From the surgeons’ perspective, the first hope of a typical patient was correction of the prolapse (11 of 16 surgeons put this hope in first place) and the second was improved urinary function (6 of 16 put this hope in second place; Table 3). As the first fear of a typical patient, 8 surgeons chose the risk of failure or relapse, and as the second fear, concerns about the operation (Table 4).




Figure 3. Women’s preoperative hopes.




Figure 4. Women’s preoperative fears.

4. Discussion

Women awaiting a cystocele repair reported as their main hopes the repair of their prolapse, improvement in their urinary symptoms, capacity for physical activities, sexual function, well-being, and end of pain or heaviness. Their main fears were related to prolapse relapse, perioperative concerns, urinary disorders, pain, and sexual problems. These reasonable expectations were consistent with the information the surgeons had given them. Our results show that surgeons were able to anticipate typical hopes and fears.

The prospective collection of these hopes and fears—before the surgery—is one of the strong points of our work, for it enabled us to measure the quality of the information as the

women perceived it. The other strong point was that the information was standardized and formalized by its delivery in a leaflet and by the woman's signature consenting to participate in the study.

One of the limitations of our work is the surgeons' low response rate to this questionnaire. It is nonetheless consistent with the response rate expected in surveying these specialists [10]. We assume that this low response rate does not constitute a substantial bias, given that the members of the PROSPERE group habitually work together and regularly exchange information about their surgical practices [11]. Moreover, the PROSPERE investigators were involved in drafting the information leaflet and participated in conversations about every element of this information. However, it is possible that some surgeons did not respond because they did not feel sufficiently aware of each patients' expectations to be able to respond appropriately. Another limitation is that we asked the surgeons about what they expected a typical patient would hope and fear; it would have been preferable to ask their opinion for each woman they included.

Our preliminary hypothesis was that some women might have hopes or fears that were exaggerated and did not correspond to the medical reality of these operations. On the contrary, we observed that they had reasonable expectations of the prolapse surgery: only 60% expected the complete disappearance of the prolapse, and one-third (38%) feared the risk of failure or relapse. These results are consistent with the high failure rate related to prolapse surgery [12], and suggest that surgeons do take the time to explain the risk of failure.

Alongside this focus on the anatomical results of prolapse surgery, many of the women's expectations concerned functional results, with a quarter or more hoping for improvement in their urinary symptoms, physical capacity, sexual function, and comfort. Although the participants were aware of mesh-related complications (the study of which was the PROSPERE trial's main objective, and which were clearly mentioned in the information sheet), less than 5% listed this specific fear.

Although studies show that POP has a negative impact on women's sexual function [13] and that surgical management significantly improves it [14], we found that expectations of its improvement were not the women's main priority. This prudence of the expectations of the women in our study is consistent with the results of a study showing that after surgery for POP, women's expectations concerning physical symptoms and activity were met more often than those concerning their sexual function [15].

The French women included in our study reported expectations similar to those of the Dutch women questioned by Lawndy et al., although there were some differences in rankings [16]. While prolapse repair was the main hope (60%) in our study, Lawndy's participants (70%) ranked the improvement of urinary symptoms first. The main fear in our study was failure or relapse (38%), while becoming incontinent was the main fear (52%) among the Dutch women. These variations may reflect differences in patient information or in the social and cultural environment between French and Dutch practices.

Surgeons who participated in the study successfully anticipated the hopes and fears reported by most women, probably because they practice this type of surgery on a regular basis. However, our study shows that the hopes and fears of women scheduled for POP surgery may vary broadly. A study analyzing interviews after surgery shows that women's expectations focus principally on the disappearance of their physical symptoms [17]. Another work shows that women's expectations can be multiple and concern domains as varied as urinary or gastrointestinal symptoms, physical activity, sexual function, and self-image [18].

The PROSPERE trial shows that the women who expressed preoperative expectations related to improvement of bulge symptoms were those most likely to be satisfied after the surgery [19]. We believe that our report on women's typical expectations before prolapse repair is helpful information for physicians, but our results cannot replace an individual assessment of expectations before surgery. Anticipating typical expectations should help physicians in their individual assessment before surgery (and not replace it). As part of a

patient-centered approach and individualized care pathways, identifying some expectations and fears could improve treatment adherence and patient satisfaction. Improving the doctor–patient relationship could therefore improve overall postoperative results. The quality of the preoperative information could also be integrated into this approach.

Our results support an individual patient’s hope and fear assessment before each surgical decision. Further work in this area is welcome to promote understanding and compliance with our patients’ expectations in routine clinical practice.

5. Conclusions

Our study may help surgeons to consider women’s expectations before pelvic-floor reconstructive surgery.

Author Contributions: Conceptualization, X.F.; methodology, X.F.; validation, A.F.; formal analysis, X.F. and M.R.; investigation, X.F., A.-C.P., S.C.-L., G.B., P.C., M.C., P.D., X.D., H.F., P.F., O.G., B.J., G.L., C.S., R.d.T., L.W., J.-P.L., A.F.; data curation, M.R.; writing—original draft preparation, X.F.; writing—review and editing, A.F. and A.-C.P. All authors have read and agreed to the published version of the manuscript.

Funding: Our ancillary qualitative analysis in the PROSPERE trial received no external funding.

Institutional Review Board Statement: The Comité de Protection des Personnes Nord-Ouest IV (Institutional Review Board, Lille, France) approved the PROSPERE study.

Informed Consent Statement: Written informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data from the analysis are available upon request from the authors.

Acknowledgments: We thank all women included in the PROSPERE trial. We thank the team of PROSPERE surgeons not listed as authors who helped make this work possible: Cherif Akladios, Jean-Sebastien Aucouturier, Emmanuelle Baulon-Thaveau, Sébastien Blanc, Pierre Collinet, Christine Dehghani, Patrick Delporte, Géraldine Demoulin, Erika Faivre, Brigitte Fatton, Sandrine Fournet, Catherine Frey, Pierre Gadonneix, Geraldine Giraudet, Amirata Kane, Magali Lesavre, Cédric Nadeau, André Nazac, Delphine Salet-Lizée, Denis Savary, Véronique Thoma, Didier Tardif, Yannick Thirouard, Caroline Trichot, Richard Villet, Etienne Vincens, Arnaud Wattiez.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Fritel, X.; Varnoux, N.; Zins, M.; Bréart, G.; Ringa, V. Symptomatic pelvic organ prolapse at midlife, quality of life, and risk factors. *Obstet. Gynecol.* **2009**, *113*, 609–616. [[CrossRef](#)] [[PubMed](#)]
2. Smith, F.J.; Holman, C.D.; Moorin, R.E.; Tsokos, N. Lifetime Risk of Undergoing Surgery for Pelvic Organ Prolapse. *Obstet. Gynecol.* **2010**, *116*, 1096–1100. [[CrossRef](#)] [[PubMed](#)]
3. Fritel, X.; Panjo, H.; Varnoux, N.; Ringa, V. The individual determinants of care-seeking among middle-aged women reporting urinary incontinence: Analysis of a 2273-woman cohort. *Neurourol. Urodyn.* **2014**, *33*, 1116–1122. [[CrossRef](#)] [[PubMed](#)]
4. Fauconnier, A.; Staraci, S.; Huchon, C.; Roman, H.; Panel, P.; Descamps, P. Comparison of patient- and physician-based descriptions of symptoms of endometriosis: A qualitative study. *Hum. Reprod.* **2013**, *28*, 2686–2694. [[CrossRef](#)] [[PubMed](#)]
5. Barber, M.D.; Brubaker, L.; Nygaard, I.; Wheeler, T.L., 2nd; Schaffer, J.; Chen, Z.; Spino, C. Defining Success After Surgery for Pelvic Organ Prolapse. *Obstet. Gynecol.* **2009**, *114*, 600–609. [[CrossRef](#)] [[PubMed](#)]
6. Digesu, A.; Chaliha, C.; Salvatore, S.; Hutchings, A.; Khullar, V. The relationship of vaginal prolapse severity to symptoms and quality of life. *BJOG* **2005**, *112*, 971–976. [[CrossRef](#)] [[PubMed](#)]
7. Braddock, C.H., 3rd; Edwards, K.A.; Hasenberg, N.M.; Laidley, T.L.; Levinson, W. Informed decision making in outpatient practice: Time to get back to basics. *JAMA* **1999**, *282*, 2313–2320. [[CrossRef](#)] [[PubMed](#)]
8. Lucot, J.P.; Cosson, M.; Bader, G.; Debodinance, P.; Akladios, C.; Salet-Lizée, D.; Delporte, P.; Savary, D.; Ferry, P.; Deffieux, X.; et al. Safety of Vaginal Mesh Surgery Versus Laparoscopic Mesh Sacropepy for Cystocele Repair: Results of the Prosthetic Pelvic Floor Repair Randomized Controlled Trial. *Eur. Urol.* **2018**, *74*, 167–176. [[CrossRef](#)] [[PubMed](#)]
9. Srikrishna, S.; Robinson, D.; Cardozo, L.; Cartwright, R. Experiences and expectations of women with urogenital prolapse: A quantitative and qualitative exploration. *BJOG* **2008**, *115*, 1362–1368. [[CrossRef](#)] [[PubMed](#)]
10. Stöss, C.; Steffani, M.; Pergolini, I.; Hartmann, D.; Radenkovic, D.; Novotny, A.; Friess, H.; Müller, M.W. Impact of the COVID-19 Pandemic on Surgical Oncology in Europe: Results of a European Survey. *Dig. Surg.* **2021**, *38*, 259–265. [[CrossRef](#)] [[PubMed](#)]

11. Jacquetin, B.; Hinoul, P.; Gauld, J.; Fatton, B.; Rosenthal, C.; Clave, H.; Garbin, O.; Berrocal, J.; Villet, R.; Salet-Lizée, D.; et al. Total transvaginal mesh (TVM) technique for treatment of pelvic organ prolapse: A 5-year prospective follow-up study. *Int. Urogynecol. J.* **2013**, *24*, 1679–1686. [[CrossRef](#)] [[PubMed](#)]
12. Olsen, A.; Smith, V.J.; Berstrom, J.O.; Colling, J.C.; Clark, A.L. Epidemiology of surgically managed pelvic organ prolapse and incontinence. *Obstet. Gynecol.* **1997**, *89*, 501–506. [[CrossRef](#)] [[PubMed](#)]
13. Verbeek, M.; Hayward, L. Pelvic floor dysfunction and its effect on quality of sexual life. *Sex. Med. Rev.* **2019**, *7*, 559–564. [[CrossRef](#)] [[PubMed](#)]
14. Jha, S.; Gray, T. A systematic review and meta-analysis of the impact of native tissue repair for pelvic organ prolapse on sexual function. *Int. Urogynecol. J.* **2015**, *26*, 321–327. [[CrossRef](#)] [[PubMed](#)]
15. Srikrishna, S.; Robinson, D.; Cardozo, L.; Thiagamoorthy, G. Patient and surgeon goal achievement 10 years following surgery for pelvic organ prolapse and urinary incontinence. *Int. Urogynecol. J.* **2015**, *26*, 1679–1686. [[CrossRef](#)] [[PubMed](#)]
16. Lawndy, S.S.; Withagen, M.I.; Kluivers, K.B.; Vierhout, M.E. Between hope and fear: Patient's expectations prior to pelvic organ prolapse surgery. *Int. Urogynecol. J.* **2011**, *22*, 1159–1163. [[CrossRef](#)] [[PubMed](#)]
17. Baskayne, K.; Willars, J.; Pitchforth, E.; Tincello, D.G. Women's Expectations of Prolapse Surgery: A Retrospective Qualitative Study. *Neurourol. Urodyn.* **2014**, *33*, 85–89. [[CrossRef](#)] [[PubMed](#)]
18. Hullfish, K.L.; Bovbjerg, V.E.; Gibson, J.; Steers, W.D. Patient-centered goals for pelvic floor dysfunction surgery: What is success, and is it achieved? *Am. J. Obstet. Gynecol.* **2002**, *187*, 88–92. [[CrossRef](#)] [[PubMed](#)]
19. Chattot, C.; Deffieux, X.; Lucot, J.P.; Fritel, X.; Fauconnier, A. Preoperative predictors and a prediction score for perception of improvement after mesh prolapse surgery. *Int. Urogynecol. J.* **2020**, *31*, 1393–1400. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.