

Table S1. Baseline characteristics

| Variable | All (n=95) |
|--------------------------------------|---------------|
| Age | 74.1±10.8 |
| Female | 79 (83.2%) |
| Body mass index (kg/m ²) | 20.9±3.7 |
| Hypertension | 47 (49.5%) |
| Diabetes | 14 (14.7%) |
| Dyslipidemia | 26 (27.4%) |
| Current smoker | 14 (14.7%) |
| Symptom at admission | |
| Chest pain | 35 (36.8%) |
| Dyspnea | 25 (26.3%) |
| Triggers | |
| Emotional stress | 16 (16.8%) |
| Physical stress | 60 (63.2%) |
| No apparent trigger | 24 (25.3%) |
| ECG findings at presentation | |
| ST elevation | 42(44.2%) |
| QTC, msec | 464.6±56.3 |
| Maximal CK myocardial band (U/L) | 19.1±20.5 |
| LV ejection fraction (TTE), % | 50.7±12.6 |
| Ballooning pattern | |
| Apical ballooning type | 60 (63.2%) |
| Midventricular type | 27 (28.4%) |
| Basal type | 2 (2.1%) |
| Focal type | 5 (5.2%) |

Values are shown as mean ± SD, median(IQR), or n(%), CK = creatine kinase, ECG = electrocardiography; QTc = corrected QT interval; LV = left ventricular;