

Supplementary Materials: Calcium Kidney Stones are Associated with Increased Risk of Carotid Atherosclerosis: the Link between Urinary Stone Risks, Carotid Intima-Media Thickness, and Oxidative Stress Markers

Ho Shiang Huang, Pao Chi Liao and Chan Jung Liu *

Table 1. Stone composition in patients with kidney stone group.

| Variables | Kidney Stone Group (n = 114) |
|------------------------------------|------------------------------|
| Mean ± SD% stone composition | |
| CaOx | 53.52 ± 39.78 |
| CaP | 28.79 ± 33.66 |
| Ammonium magnesium | 9.56 ± 29.40 |
| UA | 4.47 ± 20.57 |
| Sodium urate | 2.07 ± 13.40 |
| No. containing any composition (%) | |
| CaOx | 87 (76.3) |
| CaP | 91 (79.8) |
| Ammonium magnesium | 11 (9.6) |
| UA | 6 (5.3) |
| Sodium urate | 3 (2.6) |
| No. pure (%) | |
| CaOx | 6 (20.0) |
| CaP | 7 (23.3) |
| Ammonium magnesium | 10 (33.3) |
| UA | 5 (16.7) |
| Sodium urate | 2 (6.7) |
| No. mixed (%) | |
| No. greater than 70% mixed (%) | 84 (73.7) |
| CaOx | 55 (57.9) |
| CaP | 22 (23.2) |
| Ammonium magnesium | 11 (11.6) |
| UA | 5 (5.3) |
| Sodium urate | 2 (2.1) |

SD: Standard deviation; No.: Number; CaOx: Calcium oxalate; CaP: Calcium phosphate; UA: Uric acid.