

Supplementary Files




Figure S1- List of the ten most significant terms in each topic calculated with the β method. Topics were arbitrarily classified as environmental dynamics (1, 7 and 8), weed management (4 and 10), quantification methodologies (3), omics (5 and 9), and generic (2 and 6).




Figure S2- Ordinary axis graph showing the selection of included/excluded records by topic. Topics were arbitrarily classified as environmental dynamics (1, 7 and 8), weed management (4 and 10), quantification methodologies (3), omics (5 and 9), and generic (2 and 6).




Figure S3 – Color guides of the bibliographic records of the data used throughout the text.

CHEN, X.; HU, J.; LI, J. Determination of imazaquin residues in soil by solid-phase extraction and high-performance liquid chromatography. *Journal of AOAC International*, 90, n. 2, p. 568-574, 2007.

COBUCCI, T.; PRATES, H. T.; FALCAO, C. L. M.; REZENDE, M. M. V. Effect of imazamox, fomesafen, and acifluorfen soil residue on rotational crops. *Weed Science*, 46, n. 2, p. 258-263, 1998.

CURRAN, W. S.; LIEBL, R. A.; SIMMONS, F. W. Effects of tillage and application method on clomazone, imazaquin, and imazethapyr persistence. *Weed Science*, 40, n. 3, p. 482-489, 1992.

DUMAS, J. A.; SEMIDEY, N.; CACHO, C.; CASANOVA, P. Field dissipation of prometryn, oxyfluorfen and imazethapyr in Mollisol and Vertisol soils. *Journal of Agriculture of the University of Puerto Rico*, 92, n. 3-4, p. 183-195, 2008.

FLINT, J. L.; WITT, W. W. Microbial degradation of imazaquin and imazethapyr. *Weed Science*, 45, n. 4, p. 586-591, 1997.

GÁMIZ, B.; VELARDE, P.; SPOKAS, K. A.; CELIS, R. et al. Changes in sorption and bioavailability of herbicides in soil amended with fresh and aged biochar. *Geoderma*, 337, p. 341-349, 2019.

GIANELLI, V.; BEDMAR, F.; COSTA, J. Persistence and sorption of imazapyr in three Argentinean soils. *Environmental Toxicology and Chemistry*, 33, n. 1, p. 29-34, 2014.

GOETZ, A. J.; LAVY, T. L.; GBUR, E. E. Degradation and field persistence of imazethapyr. *Weed Science*, 38, n. 4-5, p. 421-428, 1990.

HU, M.; LIU, K.; QIU, J.; ZHANG, H. et al. Behavior of imidazolinone herbicide enantiomers in earthworm-soil microcosms: Degradation and bioaccumulation. *Science of the Total Environment*, 707, 2020.

ISMAIL, B. S.; AHMAD, A. R. Attenuation of the herbicidal activities of glufosinate-ammonium and imazapyr in 2 soils. *Agriculture Ecosystems & Environment*, 47, n. 4, p. 279-285, 1994.

JENSEN, K. I. N.; IVANY, J. A.; KING, R. R. Persistence of imazethapyr in two Atlantic Canada soils. *Canadian Journal of Soil Science*, 75, n. 4, p. 525-527, 1995.

JOVANOVIC-RADOVANOV, K. Imazethapyr persistence in sandy loam detected using white mustard bioassay. *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes*, 52, n. 10, p. 711-718, 2017.

LOPEZ-CABEZA, R.; GAMIZ, B.; CORNEJO, J.; CELIS, R. Behavior of the enantiomers of the herbicide imazaquin in agricultural soils under different application regimes. *Geoderma*, 293, p. 64-72, 2017.

MCDOWELL, R. W.; CONDRON, L. M.; MAIN, B. E.; DASTGHEIB, F. Dissipation of imazapyr, flumetsulam and thifensulfuron in soil. *Weed Research*, 37, n. 6, p. 381-389, 1997.

MILLS, J. A.; WITT, W. W. Efficacy, phytotoxicity, and persistence of imazaquin, imazethapyr, and clomazone in no-till double-crop soybeans (*Glycine max*). *Weed Science*, 37, n. 3, p. 353-359, 1989.

RAMEZANI, M.; OLIVER, D. P.; KOOKANA, R. S.; LAO, W. et al. Faster degradation of herbicidally-active enantiomer of imidazolinones in soils. *Chemosphere*, 79, n. 11, p. 1040-1045, 2010.

RANI, D.; DUHAN, A.; PUNIA, S. S.; YADAV, D. B. et al. Behavior of pre-mix formulation of imazethapyr and imazamox herbicides in two different soils. *Environmental Monitoring and Assessment*, 191, n. 1, 2019.

SCHROEDER, J. Chlorimuron and Imazaquin Persistence in Selected Southern Soils under Controlled Conditions. *Weed Science*, 42, n. 4, p. 635-640, 1994.

SU, W.; HAO, H.; DING, M.; WU, R. et al. Adsorption and degradation of imazapic in soils under different environmental conditions. *Plos One*, 14, n. 7, 2019.

VENCILL, W. K.; BANKS, P. A.; BARRETT, M.; BRECKE, B. et al. Dissipation of imazaquin in southern soils. *Journal of Environmental Science and Health Part B-Pesticides Food Contaminants and Agricultural Wastes*, 30, n. 5, p. 621-635, 1995.

VISCETTI, C. Measured and simulated persistence of imazethapyr in soil. *Bulletin of Environmental Contamination and Toxicology*, 54, n. 3, p. 420-427, 1995.

VISCETTI, C.; CASUCCI, C.; PERUCCI, P. Relationship between changes of soil microbial biomass content and imazamox and benfluralin degradation. *Biology and Fertility of Soils*, 35, n. 1, p. 13-17, 2002.

WANG, H.-I.; LI, Y.-y.; WEI, G.-h.; WANG, X.-d. Imazaquin degradation and metabolism in a sandy loam soil amended with farm litters. *Journal of Environmental Sciences*, 19, n. 9, p. 1108-1113, 2007.

WANG, X.; WANG, H.; FAN, D. Degradation and metabolism of imazapyr in soils under aerobic and anaerobic conditions. *International Journal of Environmental Analytical Chemistry*, 86, n. 8, p. 541-551, 2006.

WANG, X.; ZHOU, S. M.; WANG, H. L.; FAN, D. F. Biodegradation of imazapyr in typical soils in Zhejiang Province, China. *Journal of Environmental Sciences*, 17, n. 4, p. 593-597, 2005. Article.

WU, H.; HE, X.; DONG, H.; ZHOU, Q. et al. Impact of microorganisms, humidity, and temperature on the enantioselective degradation of imazethapyr in two soils. *Chirality*, 29, n. 7, p. 348-357, 2017.

YAVARI, S.; SAPARI, N. B.; MALAKAHMAD, A.; YAVARI, S. Degradation of imazapic and imazapyr herbicides in the presence of optimized oil palm empty fruit bunch and rice husk biochars in soil. *Journal of Hazardous Materials*, 366, p. 636-642, 2019.