

## Response of Winter Wheat (*Triticum aestivum* L.) to Selected Biostimulants under Drought Conditions


Dominika Radzikowska-Kujawska <sup>1,\*</sup>, Paula John <sup>1</sup>, Tomasz Piechota <sup>1</sup>, Marcin Nowicki <sup>2</sup> and Przemysław Łukasz Kowalczewski <sup>3</sup>

<sup>1</sup>Department of Agronomy, Poznań University of Life Sciences, 11 Dojazd St., 60-632 Poznań, Poland


<sup>2</sup>Department of Entomology and Plant Pathology, Institute of Agriculture, University of Tennessee, 363 Plant Biotechnology Building, 2505 EJ Chapman Drive, Knoxville, TN 37996-4560, USA

<sup>3</sup>Department of Food Technology of Plant Origin, Poznań University of Life Sciences, 31 Wojska Polskiego St., 60-624 Poznań, Poland

\*Correspondence: dominika.radzikowska@up.poznan.pl



**Figure S1.** Photographs of wheat grown under optimal water conditions. From the left: 'Control', 'BaktoKompleks', 'Naturvital®-Plus', 'Raiza-Mix', 'Bacillus velezensis'.



**Figure S2.** Photographs of wheat grown under drought stress. From the left: 'Control', 'BaktoKompleks', 'Naturvital®-Plus', 'Raiza-Mix', 'Bacillus velezensis'.