

Review

Impacts of Climate Change and Agricultural Practices on Nitrogen Processes, Genes, and Soil Nitrous Oxide Emissions: A Quantitative Review of Meta-Analyses

Dafeng Hui , Avedananda Ray, Lovish Kasrija and Jaekedah Christian

Department of Biological Sciences, Tennessee State University, Nashville, TN 37209, USA; aray28@tnstate.edu (A.R.); lkasrija@tnstate.edu (L.K.); jchris10@tnstate.edu (J.C.)

* Correspondence: dhui@tnstate.edu; Tel.: +1-(615)-963-5777

Abstract: Microbial-driven processes, including nitrification and denitrification closely related to soil nitrous oxide (N₂O) production, are orchestrated by a network of enzymes and genes such as *amoA* genes from ammonia-oxidizing bacteria (AOB) and archaea (AOA), *narG* (nitrate reductase), *nirS* and *nirK* (nitrite reductase), and *nosZ* (N₂O reductase). However, how climatic factors and agricultural practices could influence these genes and processes and, consequently, soil N₂O emissions remain unclear. In this comprehensive review, we quantitatively assessed the effects of these factors on nitrogen processes and soil N₂O emissions using mega-analysis (i.e., meta-meta-analysis). The results showed that global warming increased soil nitrification and denitrification rates, leading to an overall increase in soil N₂O emissions by 159.7%. Elevated CO₂ stimulated both *nirK* and *nirS* with a substantial increase in soil N₂O emission by 40.6%. Nitrogen fertilization amplified NH₄⁺-N and NO₃⁻-N contents, promoting *AOB*, *nirS*, and *nirK*, and caused a 153.2% increase in soil N₂O emission. The application of biochar enhanced *AOA*, *nirS*, and *nosZ*, ultimately reducing soil N₂O emission by 15.8%. Exposure to microplastics mostly stimulated the denitrification process and increased soil N₂O emissions by 140.4%. These findings provide valuable insights into the mechanistic underpinnings of nitrogen processes and the microbial regulation of soil N₂O emissions.



Citation: Hui, D.; Ray, A.; Kasrija, L.; Christian, J. Impacts of Climate Change and Agricultural Practices on Nitrogen Processes, Genes, and Soil Nitrous Oxide Emissions: A Quantitative Review of Meta-Analyses. *Agriculture* **2024**, *14*, 240. <https://doi.org/10.3390/agriculture14020240>

Academic Editors: Qin Peng, Xingren Liu and Wanxia Peng

Received: 2 January 2024

Revised: 29 January 2024

Accepted: 30 January 2024

Published: 1 February 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: denitrification; global warming; greenhouse gas emission; mega-analysis; nitrogen fertilizer; N₂O; precipitation

1. Introduction

In the face of a growing global population, a paramount challenge is to increase production levels of food, feed, fiber, and fuel crops while simultaneously mitigating associated environmental impacts [1–3]. To meet the ever-increasing demands for food and energy, substantial quantities of chemical fertilizers, notably inorganic nitrogen (N) fertilizers, are routinely applied to agricultural lands each year. Although essential for production, this practice has created a serious problem: the release of soil greenhouse gases, most notably nitrous oxide (N₂O), into the atmosphere [4,5]. The repeated and excessive use of N fertilizers, coupled with N deposition and climate change, has amplified challenges related to nitrate leaching and N₂O emissions. Agricultural soils contribute up to 80% of anthropogenic N₂O emissions [6–8]. Remarkably, N₂O is a potent, long-lived powerful greenhouse gas with a global warming potential 265 times greater than CO₂ [2,9]. In addition to its impact on global warming, N₂O plays a significant role in stratospheric O₃ depletion [10]. Through photolysis and oxidation to nitric oxide, N₂O can contribute to O₃ depletion in the stratosphere, further accelerating global climate change with diverse effects on human health [11,12].

In recent years, the atmospheric N₂O concentration has risen from 270 ppb during the preindustrial era to 330 ppb, with an average increase of 0.73 ppb year⁻¹ [2,13]. Global N₂O emissions stemming from N inputs have surged by more than 30% in the past four

decades [7,14]. Projections suggest that by 2030, N₂O emissions from croplands could make up 59% of global N₂O emissions [5,15]. This heightened N₂O emission disrupts greenhouse gas balances, offsetting the climate benefits gained from CO₂ removal and other climate mitigation strategies [2,16]. To address this escalating N₂O issue, a comprehensive understanding of the mechanisms and mitigation strategies for soil N₂O emission is not just valuable but indeed imperative.

Soil N₂O production and soil N cycling are intricately influenced by a diverse array of functional soil microorganisms [5,17,18]. Key players in this context include *amoA* genes of ammonia-oxidizing archaea (AOA) and ammonia-oxidizing bacteria (AOB), along with crucial functional genes such as *narG* (encoding nitrate reductase), *nirK* and *nirS* (encoding nitrite reductase), and *norB* and *nosZ* (encoding nitrous oxide reductase) genes [5,8,19]. These genes are significant actors in soil nitrification and denitrification processes. It is important to note that these nitrifying and denitrifying genes can be influenced by many factors, including climate change and various agricultural practices, leading to modifications in soil N transformation rates [5,20,21]. Consequently, evaluating the impacts of global climate change and agricultural practices on N cycling, especially concerning nitrification and denitrification, holds significant importance, as their effects on these microbial processes can induce positive feedback on climate change [7,22].

Numerous investigations have been undertaken in recent decades to explore the repercussions of climate change and agricultural practices on soil N₂O emissions in terrestrial ecosystems [2,23,24]. Due to the inconsistency among different individual studies, meta-analysis has been utilized to synthesize the results from these studies. Recently, a surge in meta-analyses has sought to quantify the impacts of these factors and practices [5,8,14,22]. However, these meta-analyses have, at times, produced varying results, promoting the need for a comprehensive evaluation [25]. This review aims to fill this gap by synthesizing the results of these meta-analyses on the impacts of climate change and agricultural practices on soil N₂O emissions. It also delves into potential mechanisms underlying these effects on enzyme activities and genes associated with nitrification and denitrification processes. This review begins with a brief overview of N cycling and the role of N₂O emissions in climate change, laying the groundwork for an in-depth discussion on the process and key genes governing soil emissions. Special emphasis is placed on recent meta-analysis studies, quantifying the impacts of climate change and agricultural practices, such as global warming, elevated CO₂, precipitation changes, N fertilization, and biochar application on soil N₂O emissions, utilizing mega-analysis (i.e., meta-meta-analysis) techniques. The primary goal of this review is to shed light on existing knowledge while identifying domains deserving of further investigation, emphasizing the significance of ongoing research in this crucial field.

2. Nitrogen Processes, Enzymes, Genes, and Soil N₂O Emissions

Soil N cycling and transforming processes in terrestrial ecosystems are predominantly regulated by soil microorganisms, with their functional genes and their extracellular enzymes playing a central role in these processes [26–29]. Soil N₂O is produced from microbial activities, involving archaea, bacteria, and fungi, engaged in the conversion of inorganic N through a series of processes. Typically, these processes encompass nitrification, which is the aerobic oxidation of NH₄⁺ to NO₃⁻ via NO₂⁻, and denitrification, the anaerobic reduction of NO₃⁻ to N₂O and N₂. This intricate cycle is tightly regulated by a spectrum of enzymes and multiple functional genes [8,30] (Figure 1).

As soil N progresses through the biogeochemical cycle in terrestrial ecosystems, it starts with crucial processes: N fixation and mineralization [31]. These processes are driven by the associated microbial communities [31]. During soil biological N fixation, molecular N is reduced to ammonia through specific biological enzymes [31,32]. Non-symbiotic N fixation, driven by free-living diazotrophs, emerges as a primary source of N in terrestrial ecosystems [33]. The key marker gene in this process is *nifH*, which encodes the nitrogenase reductase subunit responsible for reducing nitrogen gas (N₂) to ammonium (NH₄⁺) [34,35].

While most N-fixation occurs within the root nodules of legumes via symbiotic bacteria, free-living N fixation serves as a potential source for biological N inputs in non-leguminous crops [27,33]. Regarding mineralization, the N-cycling enzymes in soil microbes regulate inorganic N availability via mineralization and hydrolysis [29,36]. Key enzymes (and marker genes) involved in N mineralization include protease (*npr* and *sub*), chitinase (*chiA*), urease (*ureC*), and arginase (*rocF*) [37].




Figure 1. Nitrogen processes, genes, enzymes, soil N₂O emissions, and impact factors investigated in this study [13].

Nitrification is a fundamental process in which inorganic reduced N undergoes oxidation to become nitrate N under the action of ammonia-oxidizing microorganisms (Figure 1) [31,38]. Key genes involved in this process include *amoA*, which codes for a subunit of ammonia monooxygenase. This enzyme is pivotal for the transformation of ammonia (NH₃) or NH₄⁺ to hydroxylamine [31,35]. In particular, this first and rate-limiting step of nitrification, namely ammonia oxidation, is facilitated by either AOA (ammonia-oxidizing archaea) or AOB (ammonia-oxidizing bacteria). Both groups carry the *amoA* gene [8,12,39,40].

Denitrification is a critical process involving the reduction of nitrate (NO₃⁻) to nitrite (NO₂⁻) (Figure 1). Under anaerobic conditions, denitrifying bacteria reduce NO₃⁻ or NO₂⁻ to gaseous N, such as nitric oxide (NO) and N₂O [31,41]. This pathway is a major source of soil N₂O production. Following the reduction of NO₃⁻ to NO₂⁻, two potential pathways can be taken. It can be transformed into NH₄⁺ through dissimilatory reduction or further reduced to N₂O [35,42]. Various denitrification genes play important roles in regulating N₂O production. The denitrification process involves genes such as *narG* and *napA*, which encode subunits of two distinct nitrate reductases: membrane-bound nitrate reductase (NAR) and periplasmic nitrate reductase (NAP) [40]. *napA* is also considered a part of the assimilatory N reduction pathway. These enzymes mediate the reduction of NO₃⁻ to NO₂⁻ in both denitrification and dissimilatory NO₃⁻ reduction to NH₄⁺ [40]. Subsequently, the *nirK* and *nirS* genes, encoding nitrite reductase, are indicators of denitrifiers that facilitate the reduction of NO₂⁻ to NO, which is a rate-limiting step in denitrification [9,43,44]. The *norB* gene, encoding for the NO reductase, facilitates the reduction of NO to N₂O. Finally, the *nosZ* gene, encoding N₂O reductase, catalyzes the transformation of N₂O into N₂, representing the final step in denitrification [12,45]. These genetic markers provide insights into the complex processes of denitrification [9,31,40]. Understanding alterations in the abundance and diversity of these N functional genes provides insights into the biotic mechanisms mediating N₂O emissions [9].

Quantifying and characterizing these functional genes associated with the N biogeochemical cycle establishes a direct link between N-cycling microbial groups and the actual N processes, enriching our understanding of the ecological importance of N-cycling traits and soil N₂O emissions [27,35,40,46]. For example, nitrification inhibitors can limit the growth and activity of nitrifiers by deactivating the ammonia monooxygenase enzyme [30,47]. Elevated temperatures enhance nitrification and denitrification rates, thereby impacting soil N pools and N availability to plants [28]. These processes are intricately regulated through the interplay of various factors, including temperature, precipitation or soil moisture content, N availability, N fertilization rates, soil microbial biomass, pH, and oxygen supply [7,48] (Figure 1). The complexity of the mechanisms controlling N₂O emissions makes it challenging to predict how climate change and agricultural practices will influence these emissions. To gain deeper insights into the mechanisms responsible for soil N₂O emission under changing climate and evolving agricultural practices, it is essential to enhance our understanding of how soil microbial communities engaged in soil N cycling respond to these dynamic factors [9].

3. Methods of Study: Experimental Study, Meta-Analysis, and Mega-Analysis

To address the intricate impacts of climate change and agricultural practices on N processes, genes, and the regulation of soil N₂O emissions, numerous field and laboratory studies have been carried out [25,40,49]. For example, Xu et al. [50] reported that warmer and drier conditions have led to reduced N₂O emissions, with soil N₂O being closely associated with the abundance of *AOB*, *nirK*, and *nosZ* genes. Meanwhile, Huang et al. [51] demonstrated a 41.83% reduction of soil N₂O emission induced by a biological nitrification inhibitor, attributing the reduction to the promotion of bacteria with the *nosZ* gene, while the growth of bacteria with *nirS* and *nirK* genes was inhibited. While these studies have contributed valuable insights, the presence of disparities and contradictions among individual findings underscores the need for a systematic approach.

Meta-analysis, a statistical method that combines the results of multiple individual studies to generate an overall effect size, proves indispensable in resolving these discrepancies in ecological studies [52,53]. This approach provides a more robust estimation and helps reconcile conflicting results from individual studies. Meta-analysis has been widely applied in ecological studies to quantify the effects of climate change and agricultural practices on soil N₂O emissions [30,40]. However, the proliferation of meta-analyses has led to inconsistent findings and occasionally opposing conclusions, necessitating a critical evaluation and synthesis of existing meta-analyses [25,54–56].

Recently, Kaur et al. [25] conducted a critical assessment of 18 meta-analyses concerning the impacts of biochar application on soil N₂O emission. Their findings revealed that the reduction of soil N₂O emission ranges from no change to a substantial reduction of 58%. They further applied the approach of mega-analysis, estimating an overall reduction of 38% by synthesizing data from these 18 meta-analyses. Mega-analysis, or a meta-analysis of meta-analyses, offers a more precise estimate of the mean effect size by combining data from various meta-analyses [25,54]. This approach has been found to provide effect size quantification with low bias and high precision [55].

In this study, we conducted a comprehensive literature search for meta-analysis studies on the impacts of climate change and agricultural practices on soil N process, genes, and N₂O emission. Utilizing the keywords “meta-analysis” and “N₂O emission”, we identified 326 relevant papers through the Web of Science (Figure S1). We carefully filtered these papers, excluding those that were not true meta-analyses, such as individual field experiments, modeling studies, studies that investigated factors unrelated to climate change or agricultural practices, and those that did not cover the N processes, enzymes, and genes involved in N fixation, mineralization, nitrification, or denitrification. In total, we included 25 meta-analysis studies in this review. For studies addressing multiple meta-analyses on the impacts of climate change factors and agricultural practices, a mega-analysis was conducted to consolidate findings. Following Kaur et al. [25], the grand mean

of response ratio was calculated using the weighted mean of response ratio using the following equation:

$$RR = \frac{\sum(n_i * R Ri)}{\sum n_i} \quad (1)$$

where RR is the grand response ratio of response variables such as genes and soil N_2O emissions across all studies, $R Ri$ is the response ratio of these variables from the i th meta-analysis study, and n_i is the sample size. The 95% confidence interval of the grand mean of RR was calculated using the weighted standard error. The RR , standard error, and 95% CI were further converted to percentage change using the following equation:

$$RR (\%) = (e^{lnRR} - 1) \times 100 \quad (2)$$

4. Impacts of Climate Change on Gene, Enzyme, Nitrification and Denitrification Processes, and Soil N_2O Emissions

Climate change such as global warming, elevated CO_2 concentration, and precipitation change has the potential to exert significant influences on soil N processes and soil N_2O emissions [57–59]. Here, we collected data from meta-analyses exploring the impacts of climate change and quantitatively evaluated the response ratios concerning the effects of global warming, elevated CO_2 levels, alterations in precipitation patterns on various aspects of soil N dynamics, including soil N pools, abundance of genes, and soil N_2O emissions.

4.1. Impacts of Global Warming

Global warming has the potential to significantly influence microbially mediated N cycling processes, including N mineralization, nitrification, and denitrification. These alternations can lead to notable changes in soil N pool sizes, N availability, and soil N_2O emissions in terrestrial ecosystems [28]. A range of studies has demonstrated that increases in temperature can modify microbial N immobilization and mineralization rates [60–62]. Ecosystems and climate zones may have significant impacts on soil N processes. For example, those in forest soils can have different responses from those in grasslands. Cold regions may show more sensitive responses to warming than warm regions [48]. Additionally, several studies have reported that increased temperatures stimulate soil microbial metabolism, enhance soil enzyme activities, and accelerate the decomposition of organic matter [59]. Furthermore, elevated temperatures have been found to increase the abundances of genes like *nirK* and *nosZ* [63,64], with *nirS*-containing denitrifiers being more sensitive to temperature increases than those containing *nirK* and *nosZ* genes [57]. However, it is worth noting that several other studies have reported that elevated temperatures do not change the abundance of *amoA* genes or have found inconsistent responses of *AOA* and *AOB* to elevated temperatures [65,66].

To assess the overall impact of global warming on N processes, meta-analyses have been conducted on the impacts of temperature on soil N processes, enzyme activities, and soil functional genes involved in N_2O emission. In an early meta-analysis examining the influence of warming on soil N_2O emission, Bai et al. [67] collected 528 observations from 51 papers, revealing a non-significant mean effect size of 0.128 of soil N_2O emission by warming, based on 26 studies. Dai et al. [28] synthesized a comprehensive dataset of 1270 observations from 134 papers and revealed that elevated temperature significantly amplifies soil nitrification and denitrification rates, leading to a notable surge of up to 227% in N_2O emissions. The prevalence of the *nirS* gene increases in the presence of plants, whereas the *nosZ* gene becomes more predominant in the absence of plants at elevated temperatures. Conversely, the *AOA*, *AOB*, and *nirK* genes remain unaffected by the elevated temperature. More recently, Li et al. [58] analyzed 72 case studies from 46 papers and found that increased temperatures do not significantly affect the abundance of archaeal *amoA*, bacterial *amoA*, and *nosZ* genes, but they significantly decrease the abundances of *nirK* and *nirS* genes by 26% and 31%, respectively. Temperature increases N_2O emissions by 33%. Additionally, Salazar et al. [49] found that warming leads to increased N mineralization rates and N_2O

emissions in cold ecosystems due to heightened enzyme activity targeting relatively labile N sources rather than alternations in the abundance of N-relevant genes (e.g., *amoA* and *nosZ*). Liu et al. [68] synthesized 1845 measurements from 164 publications and found that warming significantly enhances soil N₂O emission. About 1.5 °C of experimental warming significantly stimulates N₂O emissions by 35.2%.

We quantified the impacts of warming on nitrification and denitrification genes and soil N₂O emission based on four meta-analyses (Tables 1 and 2). The results of this mega-analysis showed that warming did not influence AOA or AOB (Table 2), but reduced MBN (−15.1%), and stimulated soil N₂O emissions (147.9%) (Table 1). Warming did not change *nirK*, *nirS*, and *nosZ* (Table 2). It enhanced the mineralization rate by 153.0%, nitrification rate by 62.0%, and denitrification rate by 159.7% (Table 1). It also increased Protease by 38.7% and Urease by 216.5% (Table 1). While there was no significant impact on the abundance of AOA and AOB, warming led to a decrease in MBN, increased soil N₂O emissions, and stimulated rates of N cycling processes such as mineralization, nitrification, and denitrification. The increased enzyme activities further highlight the accelerated decomposition of organic matter and nutrient cycling under warming conditions.

Table 1. Effects of climate change factors and agricultural practices on the main microbial enzymes and nitrification and denitrification rates, based on results of meta- and mega-analyses. Values are effective size (RR, %) with 95% confidence intervals.

Treatment	Nitrogen Fixation		Mineralization			Nitrification			Denitrification		MBC	MBN	Soil N ₂ O Emission	References	
	Rate	Bacteria	Rate	Protease	Urease	NH ₄ ⁺ -N	Rate	Enzyme	NO ₃ ⁻ -N	Rate					Enzyme
Climate change															
Warming		−9.1% [−34.7%, 26.5%]	153.0% [106.9%, 209.4%]	38.7% [6.4%, 80.7%]	216.5% [59.5%, 528.0%]	−0.6% [−20.8%, 24.7%]	62.0% [33.2%, 97.1%]		8.6% [−14.0%, 37.4%]	159.7% [127.1%, 196.9%]		−15.1% [−27.4%, −0.7%]	147.9% [92.2%, 219.7%]	[28,49,58, 59]	
Elevated CO ₂					−0.3% [−7.1%, 7.0%]	32.7% [7.4%, 63.9%]	−18.4% [−32.7%, −1.0%]	13.1% [4.7%, 22.1%]	5.3% [−7.7%, 20.1%]	−20.8% [−34.5%, −4.2%]	1.6% [−20.1%, 29.1%]	27.8% [16.4%, 40.3%]	40.6% [25.3%, 57.8%]	[12,22,59]	
PPT+													54.2% [29.5%, 83.7%]	[58,59]	
PPT−													−45.9% [−55.9%, −33.6%]	[58,59]	
Agricultural practices															
Nitrogen fertilization			−3.0% [−11.3%, 6.2%]	27.1% [15.0%, 40.5%]	81.1% [67.3%, 96.2%]	33.9% [17.4%, 52.9%]	198.0% [98.4%, 347.8%] 216.0% [99.5%, 400.5%]		94.2% [68.4%, 123.9%]	42.0% [9.6%, 84.0%]	28.5% [2.4%, 61.5%]	113.7% [−87.2%, 3463%]	28.4% [7.4%, 53.6%]	153.2% [109.6%, 205.9%]	[9,27,35, 69–72]
Biochar addition	13.2% [2.9%, 24.6%]	44.4% [29.4%, 61.1%]	14.4% [1.2%, 29.3%]	4.0% [−18.2, 32.3%]	6.0% [1.0%, 11.3%]	40.8% [1.8%, 94.8%]		3.4% [0.4%, 6.5%]	13.3% [−2.9%, 32.2%]	27.3% [−0.4%, 62.7%]		13.2% [8.3%, 18.3%]	−15.8% [−20.5%, −10.8%]	[14,40,72]	
Nitrification inhibitor					87.8% [68.2%, 109.6%]	−21.3% [−26.6%, −15.7%]	−32.8% [−38.8%, −26.4%]	−37.7% [−41.9%, −33.3%]		−25.3% [−34.1, −15.2%]			−56.1% [−64.8%, −45.0%]	[5,30,72]	
Liming		−20.8% [−39.4%, 3.6%]			−3.1% [−24.3%, 23.8%]	62.7% [27.6%, 107.4%]		55.8% [35.2%, 79.6%]	201.1% [94.7%, 365.7%]				−37.9% [−48.2%, −25.5%]	[73]	
Microplastics		4.9% [−9.9%, 22.1]	149.2% [112.0%, 192.8%]	6.8% [2.1%, 11.7%]	−6.8% [−12.8%, −0.4%]	0.9% [−2.5%, 4.5%]		−22.3% [−28.3%, −16.0%]	17.8% [1.4%, 36.8%]			27.5% [4.4%, 55.8%]	140.4% [73.7%, 232.7%]	[74]	

Table 1. Cont.

Treatment	Nitrogen Fixation		Mineralization			Nitrification			Denitrification		MBC	MBN	Soil N ₂ O Emission	References	
	Rate	Bacteria	Rate	Protease	Urease	NH ₄ ⁺ -N	Rate	Enzyme	NO ₃ ⁻ -N	Rate					Enzyme
Crop diversity			−6.6% [−21.1%, 10.5%]			−11.4% [−23.4%, 2.4%]	−24.4% [−35.1%, −12.0%]			−6.8% [−24.6%, 15.1%]	−24.6% [−49.1%, 11.7%]		−19.9% [−32.0%, −5.6%]	[31]	
Grazing			−6.6% [−21.1%, 10.5%]			−11.4% [−23.4%, 2.4%]	−24.4% [−35.1%, −12.0%]			−6.9% [−24.6%, 15.1%]	−24.6% [−49.1%, 11.7%]		−19.9% [−32.0%, −5.6%]	[8]	
Earthworm			217.8% [45.5% , 594.1%]			71.2% [31.6% , 122.6%]	58.5% [−40.9%, 324.7%]			17.9% [0.4%, 38.6%]	38.6% [17.8% , 63.0%] 0.3 [0.2,0.5]		13.7% [0.3% , 28.8%]	28.9% [0, 68.0%]	[75]

Note: Bolded values indicate a significant effect. PPT+ indicates increased precipitation. PPT− indicates reduced precipitation. MBC: microbial biomass carbon. MBN: microbial biomass nitrogen.

Table 2. Effects of climate change factors and agricultural practices on the abundance of main genes involved in the processes of nitrification and denitrification based on the results of meta- and mega-analyses. Values are effective size (RR, %) with 95% confidence intervals.

Treatment	Nitrogen Fixation		Nitrification			Denitrification			Reference
	<i>nifH</i>	<i>AOA</i>	<i>AOB</i>	<i>narG</i>	<i>nirS</i>	<i>nirK</i>	<i>norB</i>	<i>nosZ</i>	
Climate change									
Warming	−5.1% [−11.3%, 6.7%]	0.7% [−15.5%, 20.0%]	−1.34% [−22.0%, 24.7%]		12.7% [−2.3%, 30.0%]	−9.7% [−23.5%, 6.7%]		50.9% [−6.9%, 144.6%]	[28,49,58,59]
Elevated CO ₂		5.7% [−9.8%, 23.9%]	12.8% [−9.5%, 40.6%]		18.0% [0.4% , 38.6%]	18.6% [−2.8%, 43.3%]		2.6% [−16.9%, 26.7%]	[12,22,59]
PPT+		−5.1% [−29.9%, 28.4%]	−33.2% [−44.7%, 9.4%]		31.2% [−9.0%, 89.1%]	−5.0% [−20.9%, 14.2%]		4.2% [−15.6%, 28.5%]	[58,59]
PPT−		28.5% [−20.0%, 106.6%]	23.4% [−4.1%, 58.8%]		173.6% [39.5% , 436.5%]	79.6% [−8.0%, 250.6%]		48.7% [−19.6%, 175.0%]	[58,59]
Agricultural practices									
Nitrogen fertilization	−5.9% [−23.5%, 15.6%]	15.0% [− 0.8% , 33.3%]	146.3% [108.9% , 190.4%]		31.8% [6.5% , 63.0%]	43.2% [12.2% , 82.8%]		38.2% [−6.9%, 105.1%]	[9,27,35,69–72]
Biochar addition	4.7% [−9.8%, 21.6%]	23.0% [10.1% , 37.4%]	11.0% [−4.3%, 28.7%]	17.2% [0.2% , 37.1%]	19.1% [6.2% , 33.7%]	28.3% [13.1% , 45.4%]		17.1% [7.5% , 27.6%]	[14,40,72]
Nitrification inhibitor		−6.3% [−14.0%, 2.1%]	− 51.9% [− 61.5% , −40.0%]	−4.3% [−52.5%, 93.1%]	− 22.7% [− 37.9% , −3.9%]	− 20.0% [− 28.7% , −10.3%]		−0.04% [−16.3%, 19.7%]	[5,30,72]
Liming		70.2% [16.2% , 149.2%]	132.6% [35.2% , 299.9%]		37.5% [9.9% , 72.0%]	142.0% [54.3% , 279.6%]		16.0% [−1.5%, 36.5%]	[73]
Microplastics		0.4% [−11.2%, 14.2%]		− 5.2% [− 8.7% , −1.5%]	18.6% [5.1% , 34.0%]		− 36.8% [− 55.5% , −10.1%]	10.6% [2.7% , 19.1%]	[74]
Crop diversity	33.8% [17.3% , 52.2%]	18.2% [−1.4%, 41.6%]	−2.8% [−12.7%, 8.3%]	17.8% [5.1% , 32.2%]	39.4% [18.5% , 63.9%]	14.0% [0.5% , 29.3%]		3.8% [−8.4%, 17.5%]	[31]
Grazing		− 34.9% [− 57.5% , −1.99%]	−28.5% [−57.3, 19.5%]	− 28.3% [− 40.5% , −14.0%]	− 35.3% [− 55.3% , −6.1%]	−3.4% [−61.8%, 142.7%]		−22.1% [−42.2%, 4.9%]	[8]
Earthworm	27.3% [−7.2%, 74.8%]	269.6% [−34.1%, 1974.4%]	10.8% [1.3% , 21.1%]			−12.3% [−32.6%, 14.2%]		5.1% [−8.0%, 21.0%]	[75]

Note: Bolded values indicate significant effects. PPT+ indicates increased precipitation. PPT− indicates reduced precipitation. *AOA*: *amoA* genes from ammonia-oxidizing archaea. *AOB*: *amoA* genes from ammonia-oxidizing bacteria. *narG*: nitrate reductase. *nirS* and *nirK*: nitrite reductase. *nosZ*: N₂O reductase.

4.2. Impacts of Elevated CO₂

It is well documented that elevated CO₂ enhances plant growth and biomass production and increases ecosystem carbon sequestration [13,53,58]. Elevated CO₂ can also have several other impacts, such as promoting organic C decomposition, enhancing microbial activity, and stimulating soil extracellular enzyme activity [59,76]. However, the impacts of elevated CO₂ on nitrification, denitrification, and associated functional genes are still a topic of ongoing research, and the results are not consistent. Various studies have reported diverse responses to elevated CO₂ in the nitrification and denitrification rates, with some showing negative, positive, or neutral effects of elevated CO₂ on these processes [13,77]. Variations also exist in the responses of nitrifying and denitrifying functional genes to elevated CO₂ conditions. Different studies have reported divergent outcomes, indicating that the amounts of *AOA*, *AOB*, *nirK*, *nirS*, and *nosZ* functional genes may exhibit increases, decreases, or remain unaffected under elevated CO₂ levels [77–79].

We found that four meta-analyses have been published on the impacts of elevated CO₂ on genes involved in N processes and N₂O emissions. Barnard et al. [80] reviewed the impacts of elevated CO₂, N, and temperature on nitrification, denitrification, and soil N₂O emission and found that elevated CO₂ enhanced net nitrification, reduced potential denitrification (−18%), increased net nitrification (33%), and did not significantly alter soil N₂O emissions. Li et al. [59] analyzed the impacts of multiple climate factors on N-cycling genes and found that elevated CO₂ increased N-cycling functional gene abundances (19.5%). In particular, elevated CO₂ increased *nirK* but did not change *AOB*. Du et al. [12] collected data from 50 publications and reported that elevated CO₂ enhanced N₂O emissions by 44%. Elevated CO₂ increases the abundance of *AOB* (21%), *nirK* (15%), and *nirS* (15%) but does not change *AOA* and *nosZ*. Gineyts and Niboyet [22] used 879 observations from 58 papers and found that elevated CO₂ increased *AOA* (62%), *nirK* (32%), and *nirS* (27%), leading to 26% increases in soil N₂O emission.

Synthesizing these meta-analyses, our results showed that elevated CO₂ increased the abundance of *nirS* (18.0%) and soil N₂O emission by 40.6% but did not significantly change *AOA*, *AOB*, *nirK*, and *nosZ* (Tables 1 and 2). Elevated CO₂ also increased MBN (27.8%), net nitrification rate (32.7%), and NO₃[−]-N (13.1%) but reduced nitrifying enzymes by 18.4% [12,22,59,80] (Table 1). While there was an increase in soil N₂O emissions, possibly associated with changes in denitrification (as indicated by increased *nirS* abundance), there were mixed effects on nitrification-related parameters. The increased MBN, net nitrification rate, and NO₃[−]-N levels indicate the stimulation of N cycling processes, but the reduction in nitrifying enzymes suggests a potential deceleration of ammonia oxidation. These findings highlight the need to consider multiple factors influencing soil N dynamics in the context of elevated CO₂.

4.3. Impacts of Precipitation

In terrestrial ecosystems, precipitation change can have multifaceted effects. These alterations influence soil microclimate and impact the soil water balance, soil aeration, nutrient availability, and microbial ecology [58,67]. Consequently, they play a role in shaping soil N₂O emissions. For example, Šťovíček et al. [81] found that soil microbial diversity tends to be high under dry conditions due to the fragmentation of niches in dry soils. However, drought can also reduce the genetic potential and stability of soil microbiomes [59]. Homyak et al. [82] investigated the effects of reduced precipitation on soil N₂O emissions and found that a reduction in precipitation significantly lowers soil N₂O emissions, suggesting that denitrification is more sensitive to drought than processes controlling N supply. Decreased precipitation appears to have minimal effects on the abundances of archaeal *amoA*, bacterial *amoA*, *nirK*, and *nosZ*, but it shows positive effects on the abundances of *nirS* [58]. Conversely, increased precipitation has little effect on the abundances of archaeal *amoA*, *nirK*, *nirS*, and *nosZ* while exhibiting negative effects on the abundances of bacterial *amoA* [58].

Several meta-analyses have been conducted to further understand the impacts of precipitation on soil N₂O emissions [58,83]. For example, Yan et al. [83] performed a meta-analysis incorporating 84 published studies and found that increased precipitation significantly increases N₂O emissions (+154.0%), whereas decreased precipitation significantly decreases N₂O emissions (−64.7%). They also found that precipitation increases enhanced soil N₂O emissions by 128.3% in temperate forests and by 179.6% in boreal forests but did not influence soil N₂O emissions in grasslands. The impacts of decreased precipitation also varied in different ecosystems, ranging from no effect in subtropical forests to −24.3% in temperate forests and −92.6% in grasslands. However, only two papers have synthesized the impacts of precipitation on N-cycling genes and soil N₂O emission. Li et al. [58] explored the effect of global climate change on N₂O emissions and the related N functional genes in terrestrial ecosystems. Their findings indicated that precipitation promoted N₂O emissions by 55%, while reduced precipitation inhibited N₂O emissions by 31%. Based on two meta-analyses, our results showed that increased precipitation did not influence *AOA*, *AOB*, *nirK*, *nirK*, or *nosZ* (Table 2) [58,59]. Reduced precipitation did not change *AOA* and *AOB* but increased *nirS* by 173.6% and reduced soil N₂O emission by 45.9% (Tables 1 and 2) [58,59]. It is worth noting that the sample sizes of two meta-analyses were also small (18 and 20 for soil N₂O emissions). More studies are needed to focus on the N-cycling genes. Nevertheless, the findings emphasize the importance of considering the direction and magnitude of precipitation changes when assessing its impact on soil N dynamics and greenhouse gas emissions.

5. Impacts of Agricultural Practices on Gene, Enzyme, Nitrification and Denitrification, and Soil N₂O Emissions

Various agricultural management practices, including N fertilizer application, conservation tillage, cover cropping, soil amendments like biochar, and the utilization of nitrification inhibitors, can significantly affect soil physiochemical factors and microbial activity. These practices, in turn, have the potential to influence soil nitrification and denitrification processes, ultimately leading to changes in soil N₂O emissions. Notably, the application of N fertilizers, the incorporation of biochar into the soil, and the use of nitrification inhibitors are of particular importance in this context. Here, we focused on understanding the effects of these and other agricultural practices on N processes, genes, and soil N₂O emissions.

5.1. Impacts of Nitrogen Application

Nitrogen (N), an essential element, serves as a critical determinant of plant growth and productivity in terrestrial ecosystems [9,84,85]. The application of N fertilizer, a fundamental agricultural practice, has made a substantial impact on plant biomass production, soil microbial activities, and soil N₂O emissions [35,86]. The response of N cycling gene abundances to N fertilization is influenced by several factors [27]. For example, different sources of N fertilization, such as mineral and organic N fertilization, often lead to distinct changes in N-cycling gene abundances [27,32,71,87]. Additionally, N deposition increases the availability of soil nutrients and can cause soil acidification. This, in turn, affects soil element stoichiometry, nutrient utilization, and limitation, thus influencing soil microbial communities [70,88]. N deposition also has effects on functional genes related to N cycling [70]. Different studies have drawn disparate conclusions regarding the impact of functional genes and external environmental factors on N₂O emissions following N addition. For example, Soares et al. [89] found that N₂O emissions are associated with the abundance of *AOA* but exhibit no correlation with the *nirK*, *nirS*, and *nosZ* genes in fertilized soil with diverse N sources. However, Domeignoz-Horta et al. [86] analyzed over 59,000 field measurements and concluded that the diversity of the *nosZ* is the most important factor explaining the variation in N₂O emissions. Hallin et al. [90] demonstrated that a 50-year period of N fertilizer application significantly reduces potential denitrification rates and *nirK*, *nirS*, and *nosZ* gene abundance [70]. Similarly, Liang et al. [91] showed that

over 100 years of N fertilization reduces the temporal turnover of functional communities involved in denitrification. In contrast, several studies have found that N fertilization significantly increases *nirK*, *nirS*, and *nosZ* gene abundances [27,70].

The impact of N fertilization on soil greenhouse gas emissions has been well investigated and synthesized. Six meta-analyses have synthesized the impacts of N application on N processes and soil N₂O emission. Carey et al. [69] examined 98 sets of measurement data from 33 articles to understand how N additions affect the abundances of AOA and AOB during nitrification and found that nitrification genes AOA and AOB responded differently to N fertilizer, with N applications having the most effect on AOA in croplands and on AOB in wildlands [35]. Ouyang et al. [27] investigated N functional genes (*nifH*, AOA, AOB, *nirS*, *nirK*, and *nosZ*) in response to N fertilization in agricultural ecosystems. They reported that, except for *nifH*, functional gene abundances increase during nitrification (*amoA*) and denitrification (*nirK*, *nirS*, and *nosZ*) when fertilized with N. In a meta-analysis by Li et al. [70], it was observed that prolonged N fertilization leads to a substantial 75.9% increase in potential denitrification activity. Additionally, there is an elevation in the proportions of *nirK*, *nirS*, and *nosZ* gene copies. Furthermore, the denitrification N₂O/(N₂O + N₂) ratio and *nir(K + S)/nosZ* ratio also experience an increase under prolonged N fertilization.

Our mega-analysis showed that N application tended to increase AOA (15%, 95% CI was [−0.8, 33.3]) and significantly enhanced AOB by 146.3%, *nirK* by 43.2%, and *nirS* by 31.8% (Table 2). It did not change MBC, *nifH*, and *nosZ* but increased MBN by 28.4%, soil NH₄⁺-N by 33.9%, soil NO₃[−]-N by 94.2%, and total N by 27.1% (Tables 1 and 2). All meta-analyses showed an increase in soil N₂O emission, ranging from 121% to 258.8%, with a grand mean of increase in soil N₂O emission by 153.2%. The strong enhancement of soil N₂O emission was attributed to the changes in microbial community composition, stimulated N transformation processes, and elevated soil nutrient levels.

5.2. Impacts of Biochar Applications

Biochar, a recalcitrant carbonaceous biomass product generated through pyrolysis, has attracted considerable attention in agriculture for its potential to mitigate soil N losses and enhance N use efficiency [14,92,93]. Its application can enhance soil aeration, increase soil pH, promote microbial N immobilization, modify enzyme activities, and potentially affect nitrifier and denitrifier communities [94]. As reported by Zhang et al. [14] and Kaur et al. [25], the introduction of biochar into the soil has been found to elevate soil NH₄⁺ and NO₃[−]-N levels. This addition is associated with increased N mineralization, nitrification, N₂ fixation, and enhanced plant N uptake. Moreover, it also decreases N loss and N₂O emissions. An increasing number of studies demonstrate that biochar amendments can alter soil microbial communities and N-cycling gene abundance [36,40,95,96]. For example, Ducey et al. [95] found that the incorporation of biochar leads to increased levels of *nifH* and *nirS*, and *amoA* and *nosZ* in soil [40,97]. However, some other studies have not reported significant effects of biochar addition on the abundance of N-cycling microbial genes, and a few have even reported a decrease in the abundance of *nifH* and N-cycling enzyme activity [98–100]. Recent research has highlighted how biochar amendment can modify soil pH, which, in turn, affects the abundance and diversity of N-cycling genes [40,90]. Van Zwieten et al. [101] suggested that biochar addition increases the abundance of *nosZ* transcripts, consequently reducing N₂O emissions.

Quite some meta-analyses have been conducted and confirmed that biochar application can effectively reduce N losses, including N leaching and N₂O emissions [14,102–104]. Kaur et al. [25] recently synthesized 18 meta-analyses and reported that biochar application reduces soil N₂O emissions by 38%. But, only three meta-analyses synthesized genes involved in N processes [14,40,72]. Xiao et al. [40] reviewed 36 papers and found that biochar addition significantly increases the abundance of AOA, *nirK*, *nirS*, and *nosZ* by an average of 25.3%, 32.0%, 14.6%, and 17.0%, respectively, and reduced soil NH₄⁺-N (15.5%), soil NO₃[−]-N (−12.9%), and soil N₂O emission (−14.6%). Zhang et al. [14] analyzed 131 field experiments and showed that biochar significantly enhances soil NH₄⁺-N (5.3%) and NO₃[−]-N

(3.7%) contents, N mineralization (15.3 or 13.5%), nitrification (48.5%), N₂ fixation (14.7%), and plant N uptake (18.3%) but reduced N₂O emissions by 14.9%. Biochar application also increased the abundance of soil denitrifying/nitrifying genes (*amoA*, *narG*, *nirS/nirK* + *S*, and *nosZ*), the proportion of N₂ fixation bacteria, and N-acetyl-glucosaminidase activity by 18.6–87.6%. Meng et al. [72] also analyzed the impacts of biochar application on AOA, AOB, and soil N₂O emission and found that biochar application enhanced AOB and reduced soil N₂O emission.

Synthesizing these three meta-analyses, our mega-analysis showed that biochar application did not influence *nifH* and AOB but enhanced AOA by 23.0%, *nirS* by 19.1%, *nirK* by 28.3%, and *nosZ* by 17.1% (Table 2). In addition, biochar application enhanced NH₄⁺-N by 6.0%, NO₃⁻-N by 3.4%, and total N by 11.1% and reduced NH₃ emission by 34.0%, resulting in an 11.4% reduction in N leaching and 15.8% reduction in soil N₂O emissions (Table 1). The reduction of soil N₂O emissions with biochar application, despite the stimulation of N transformation genes and the enrichment of soil N levels, may be attributed to the complex interplay of various factors. Biochar's influence on microbial community composition, soil physical and chemical properties, and N retention may collectively contribute to a more balanced and controlled N cycling, reducing soil N₂O emissions. The specific mechanisms likely depend on the unique conditions of the study site and the properties of the biochar used.

5.3. Impacts of Nitrification Inhibitor Usage

Nitrification inhibitors (NIs), such as dicyandiamide (DCD), 3,4-dimethylpyrazole phosphate (DMPP), and 2-chloro-6-(trichloromethyl) pyridine (nitrapyrin), have been commonly employed to mitigate N₂O emissions by delaying the microbial oxidation of NH₄⁺ to NO₃⁻ in the soil and limiting nitrification and denitrification [30,105]. Previous individual studies have explored the influence of NIs on soil N₂O emissions and associated functional gene and transcript abundances, and community structure. However, results have been inconsistent [30]. In general, AOB tends to dominate nitrification in neutral and alkaline soils, while AOA is more prevalent in acidic environments [106]. Increasing the NH₄⁺ concentrations can enhance the nitrification activity of AOB [106], while AOA prefers environments with lower NH₄⁺ concentrations. Furthermore, many studies showed that NIs effectively decreased the AOB population but not AOA [30,65,106]. In contrast, for alkaline paddy soil in China, nitrapyrin decreased the rates of nitrification and denitrification by limiting the abundances of AOA and *nirK*, respectively [30].

Recently, four meta-analyses have investigated the impact of NIs on the abundance of N-cycling genes and the release of N₂O from the soil. Yin et al. [30] conducted a synthesis of 88 studies, revealing that the use of NIs significantly decreases the number of AOB genes, *nirS*, and *nirK* genes. NIs contributed to a 34.5% reduction in the activity of soil nitrifying enzymes (potential nitrification) and a 27.0% decrease in denitrifying enzymes (potential denitrification). Consequently, there is a notable decline of 63.6% in soil N₂O emissions. Lei et al. [106] synthesized 48 papers and reported that NIs on average reduced 58.1% of N₂O emissions and increased 71.4% of soil NH₄⁺-N concentrations. The abundance of AOB *amoA* genes was dramatically reduced by about 50% with NI application in most soil types. Meng et al. [72] synthesized several studies related to NI on N pools and process genes and found that NI increased NH₄⁺-N and total N but reduced potential nitrification and soil N₂O emission. Guo et al. [5] synthesized 166 published papers, and N-cycling inhibitors decreased soil AOB *amoA* gene abundances (212%) and significantly decreased the *nirS* gene (39%). In general, NIs consistently exhibit a substantial reduction in the release of N₂O from the soil, showing a negative correlation with the amounts of *nirK* and *nirS* genes. This consistent decrease in N₂O emissions is a common finding across all meta-analyses.

Our mega-analysis showed that NIs did not change AOA but reduced AOB by 51.9% (ranged from −4.3 to −56.7% of four meta-analyses), as well as *nirK* (−20.0%) and *nirS* (−22.7%) (Table 2). NIs enhanced soil NH₄⁺-N (87.8%) but reduced NO₃⁻-N (−37.7%), leading to a reduction of soil N₂O emission by 56.1% (from −51.7% to −62.7%) (Table 1).

5.4. Effects of Liming

Liming has the potential to reduce soil N₂O emission through two primary mechanisms [73]. Firstly, it can achieve it by increasing the population of *nosZ*-type denitrifying bacteria while decreasing the ratio of fungi to bacteria. Both of these changes contribute to a lower N₂O:N₂ production ratio. Secondly, liming can also lower the amount of soil mineral N by promoting plant uptake. Zhang et al. [73] conducted a global meta-analysis using 1474 paired observations from 124 studies to explore the responses of GHG emissions to liming. They found that liming enhances nitrification by 62.7% and denitrification by 201.1%, increases soil NO₃⁻-N by 55.8%, and reduces soil N₂O emissions by 37.9%. Liming has been found to increase the abundance of AOA by 70.2%, AOB by 132.6%, *nirS* by 37.5%, and *nirK* by 142.0% (Table 2). Interestingly, liming does not change *nosZ* copy numbers. Given its significant influence on both N₂O emissions and soil microbes, liming represents a potential strategy for mitigating soil N₂O emissions.

5.5. Impacts of Microplastics

Microplastics pose a significant threat to ecosystem health, disrupting soil biological activities, and affecting biogeochemical cycles [74,107–109]. They can alter community structures of soil microorganisms and may ultimately impact the corresponding N process [9,72,74,110]. Functional microorganisms, particularly those with the *amoA* marker gene, play a crucial role in the denitrification process [74,111]. On the other hand, functional microorganisms with marker genes, including *narG*, *napA*, and *nirS*, are responsible for the denitrification process.

Several studies, including Gao et al. [112], Li et al. [110], and Zhang et al. [108], have explored the impact of microplastics on functional microorganisms involved in the N processes. The findings indicate varying effects on specific genes associated with nitrification and denitrification. With microplastics present, contrasting trends in the copy numbers of *nirS* have been observed, with some studies reporting an increase [113] and others a decrease [114]. Similarly, *amoA* gene sizes either remain constant or decrease. The divergent reactions of genes can complicate the understanding of how nitrification and denitrification impact N₂O emissions.

Su et al. [74] conducted a meta-analysis using 60 published studies and found that in the presence of microplastics, N₂O emissions surged by 140.4%, while nitrate reductase activities increased by 4.9%. The rate of denitrification rose by 17.8%, accompanied by a 10.6% increase in the number of genes responsible for denitrification (Tables 1 and 2). This suggests that microplastics may significantly enhance the genetic potential of microorganisms to carry out denitrification. Conversely, the nitrification rate and nitrifier genes exhibited minimal changes. The changes in N processes, especially the acceleration of denitrification, were identified as key contributors to increased N₂O emissions. Microplastics may also create microenvironments that favor the growth and activity of denitrifying microorganisms, leading to remarkably increased N₂O emissions.

5.6. Impacts of Crop Diversity

Diverse crops contribute significantly to both plant biomass and play a crucial role in shaping the functional microbial communities in the soil [115,116]. The introduction of crop diversity has been shown to increase the number of *nifH* gene copies in the soil and induce changes in microbial community structure [117,118]. In the context of intercropping legumes with non-legume plants, Chen et al. [119] found that it does not have a substantial impact on the amount of *nifH*. However, in mixed teak forests, the soil experiences a decrease in the amount of AOB alongside an increase in the number of *nosZ* genes. This phenomenon correlated with variations in total N and NH₄⁺-N in the soil [120]. Moreover, agricultural systems with crop cycles, especially supplemented with inorganic N fertilizers, tend to increase the abundance of soil AOB, *nirK*, and *nosZ* genes, ultimately contributing to increased N₂O emissions [121,122].

Using a meta-analysis, Hao et al. [31] found that soil *nifH*, *nirS*, *nirK*, and *narG* abundances were positively affected by the diversity of plant species, whereas the *amoA* and *nosZ* showed no response, based on 189 observations. In particular, crop diversity significantly reduced nitrification (−24.4%) and lowered soil $\text{NH}_4^+\text{-N}$ (−11.4%) and $\text{NO}_3^-\text{-N}$ (−6.8%), resulting in a reduction in soil N_2O emissions (−19.9%) (Table 1). In addition, crop diversity enhanced *nifH* (33.8%), *narG* (17.8%), *nirS* (39.4%), and *nirK* (14.0%) (Table 2). Crop diversity enhanced the abundance of genes associated with N denitrification, reduced nitrification, lowered soil N concentrations, and significantly reduced soil N_2O emissions. These findings highlight the potential of diverse plant species in agroecosystems to positively influence soil health and mitigate environmental impacts.

5.7. Impacts of Grazing

The response of N functional gene abundances to grazing has been inconsistent [8]. In an incubation experiment, Le Roux et al. [123] found that grazed soils exhibit higher levels of AOA and AOB, along with an elevated potential nitrification rate compared to control soils that have not been grazed [8]. In an alpine meadow in China, Zhang et al. [29] reported that grazing leads to increased N_2O emissions and higher abundances of AOA, AOB, *nirK*, *nirS*, and *nosZ*. Conversely, Zhong et al. [124] reported that the moderate grazing of arid grassland does not affect the abundances of *narG* and *nosZ* or nitrification or denitrification rates. In the context of light grazing, Yin et al. [8] observed no significant effect on N_2O emissions, the nitrification rate, or the denitrification rate compared to non-grazed land. However, under moderate to heavy grazing, there are notable changes in the abundance of key N functional genes. The amounts of AOB and AOA decrease, and there are sharp reductions in the amounts of *narG* and *nirS*. Interestingly, the abundance of *nosZ* remains unaffected. Using a laboratory incubation study, Pan et al. [122] demonstrated that heavy grazing decreases the abundance of AOB and the potential nitrification rate, while light grazing increases the abundance of AOA. In contrast, grazing over nine years decreases the abundance of AOA, AOB, *narG*, *nirS*, and *nirK* in semi-arid grassland [124]. Ding et al. [125] also reported that grazing reduces the abundance of AOA, AOB, and *nirK* in loamy-sand soil.

Yin et al. [8] conducted a meta-analysis with 83 published studies and found that heavy and moderate grazing reduced N_2O emissions by 22–25%, nitrification rate by 23–37%, and denitrification rate by 44–48%, respectively, compared to the ungrazed condition. Moderate to heavy grazing intensities decreased the abundances of AOB by 40–47%. Heavy grazing also simultaneously decreased AOA by 34.9% (Table 2). Additionally, grazing significantly decreased the abundance of *narG* (−28%) and *nirS* (−35%) but did not affect the abundance of *nosZ* (Table 2). Livestock grazing at an appropriately moderate intensity is important for sustaining livestock production while contributing to greenhouse gas mitigation.

5.8. Impacts of Earthworms

While earthworms naturally inhabit the soil, incorporating sustainable agricultural practices such as increasing the return of organic matter to the soil, reducing tillage, adopting crop rotation, and avoiding the application of harmful chemicals can substantially enhance the presence of earthworms. Earthworms, often regarded as ecosystem engineers, play a crucial role in shaping soil health and ecosystem processes such as the N cycle [75]. The extent of their impact on the N cycle is closely related to their N-rich metabolic byproducts, the turnover of the N pool within the earthworm biomass, and the contribution of their deceased tissues [126,127]. Earthworms exert an indirect influence on the N cycle by changing the distribution of soil particles and incorporating pre-decomposed organic matter. Their activities, such as burrowing, contribute to an increase in the amount of macroaggregates, a factor that plays a crucial role in regulating N-cycling microorganisms in the soil [20]. Their burrowing activity further enhances N transformation through the input of organic materials into the root bioprocess [128]. Furthermore, earthworm intestinal tracts create a conducive environment for the survival of N-cycling microorganisms, stimu-

lating various N-cycle processes [129]. Earthworms have been found to exert significant effects on soil N-cycling microorganisms, such as the abundance of *amoA* gene of soil AOB, and significantly promote soil N-cycle processes, including denitrification, mineralization, and plant assimilation. In laboratory studies and experiments involving legume plants and in clay soils, earthworms are observed to significantly increase soil N₂O emissions [75].

Xue et al. [75] conducted a meta-analysis with 130 publications and found that earthworms significantly enhanced soil NH₄⁺-N (71.2%), NO₃⁻-N (17.9%), MBN (13.7%), and soil N₂O emission by 28.9% (Table 1). Earthworms also affected soil N-cycling microorganisms, including the *amoA* gene abundance of AOB (10.8%), and significantly promoted denitrification (38.6%) and mineralization (217.8%) (Tables 1 and 2). The presence of earthworms in the soil had complex effects on N dynamics and soil N₂O emission. While earthworms actively enhanced soil N levels and MBN and promoted mineralization, they also led to an increase in soil N₂O emissions. This meta-analysis reveals the positive impact of earthworms on the abundance of soil N and the available N content to soil microbes. These observed effects have the potential to alter the functions and services of ecosystems related to N cycling.

6. Regulating Factors on Soil N₂O Emissions

Several factors have been identified as influencers of soil N₂O emissions [9,29,40,58,72]. Notably, among these meta-analyses, three studies found that soil N₂O was associated with climate factors such as mean annual temperature (MAT) and mean annual precipitation MAP (Table 3). In particular, soil N₂O emission demonstrated a linear increase with MAT, while a concave relationship with soil MAP and soil moisture. Additionally, most of the studies found soil properties, especially soil available N (NH₄⁺-N and NO₃⁻-N) and C:N ratio, were associated with soil N₂O emissions. Both positive and negative relationships between soil N₂O emission and soil pH were found. Concerning N functional genes, soil N₂O was predominantly found to have a negative correlation with AOA and a positive correlation with AOB. Furthermore, *nikS* and *nikS* were identified in some studies as significantly linked to soil N₂O emissions (Table 3). However, studies also found that soil N₂O emission is not closely related to nitrifier and denitrifier abundances [29]. It is important to note that additional studies are required to establish consensus results in these areas.

Table 3. Cont.

	Elevated CO ₂		Nitrogen Fertilization		Biochar Addition		Nitrification Inhibitor		Liming	Grazing	
	Du et al. [12]	You et al. [9]	Zhang et al. [14]	Meng et al. [72]	Zhang et al. [14]	Xiao et al. [40]	Guo et al. [5]	Lei et al. [106]	Yin et al. [30]	Zhang et al. [73]	Yin et al. [8]
N functional gene											
AOA		11.6 *		−1.623 *	−0.070	−0.28 *	−0.37	−0.06	0.29	−0.90 *	
AOB		10.7 *		−0.653 *	0.849 *	−0.001	0.07	0.42 **	0.31 *	0.99 *	
<i>nosZ</i>		30.7 *			0.363	−0.12			0.29	−1.43 *	
<i>nirK</i>					0.965 *	0.20			0.50 *		
<i>nirS</i>					0.885 *	0.41 *			0.13		
<i>narG</i>					0.808 *				0.69		
Nitrification				Nonlinear *							
Denitrification							0.24			−0.25 *	
Yield-scaled NH ₃							0.48				
Treatment											
N application rate		12.1 *		Linear increase							
Nitrogen form		8.2									
Grazing intensity											13.26% *
Grazing duration											0.58%

Note: * indicates $p < 0.05$ and ** $p < 0.01$. MAP: mean annual precipitation; MAT: mean annual temperature. SOM: Soil organic matter. AOA: *amoA* genes from ammonia-oxidizing archaea. AOB: *amoA* genes from ammonia-oxidizing bacteria. *narG*: nitrate reductase. *nirS* and *nirK*: nitrite reductase. *nosZ*: N₂O reductase.

7. Conclusions and Future Research Directions

N₂O, a potent greenhouse gas originating from soil microbial processes like nitrification and denitrification, is significantly influenced by diverse factors that are shaped by climate change and agricultural practices. In this mega-analysis, we quantified their impacts on soil N₂O emissions based on 25 meta-analyses. The findings revealed that global warming substantially increased soil N₂O emissions by 159.7%, attributed to enhanced rates of both soil nitrification and denitrification. Elevated CO₂ stimulated soil N₂O emissions by 40.6%, potentially linked to changes in denitrification. The application of N fertilization emerged as a significant contributor, increasing soil N₂O emissions by 153.2%, largely associated with elevated abundances of AOB, *nirS*, and *nirK*, along with higher soil N levels. On the other hand, biochar addition reduced soil N₂O emission by 15.8%. Interestingly, microplastics had adverse impacts, stimulating soil N₂O emissions by 140.4%, primarily due to an intensified denitrification process. Additionally, earthworm activity was found to enhance soil N₂O emissions by 28.9% as earthworms actively enhanced soil N levels and MBN and promoted mineralization.

Despite the valuable insights provided by numerous previous studies for mitigating soil N₂O emissions and formulating effective management practices, critical knowledge gaps persist. The spatial and temporal variations in soil N₂O emissions, ranging from small plots to ecosystems, on regional and global scales, remain unclear. Establishing a comprehensive network for monitoring soil N₂O emissions across diverse ecosystems at large spatial scales over extended periods is crucial. This initiative not only aids in accurate emission estimates but also contributes valuable data for regional, national, and global ecosystem modeling.

Microbial process studies play a pivotal role in understanding the mechanisms of soil N₂O emissions and developing reduction strategies. Focused investigations into the microbial processes involved in N₂O production can pinpoint key microorganisms responsible for N₂O emissions. Utilizing molecular techniques like DNA sequencing and metagenomics reveals microbial community composition and diversity, shedding light on changes in soil N₂O emissions. Additionally, identifying N₂O-reducing microorganisms capable of converting N₂O to N₂ holds promise for emission mitigation. Other potential strategies include optimizing fertilizer application rates, employing nitrification inhibitors, promoting specific microbial communities, implementing sustainable practices such as biochar application, and fostering a harmonious balance between agricultural productivity and environmental sustainability in the face of evolving climate challenges.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/agriculture14020240/s1>, Figure S1: Number of publications based on the topic search of “meta-analysis” and “N₂O emission” using Web of Science.

Author Contributions: Conceptualization, D.H.; methodology, D.H., A.R., L.K. and J.C.; writing—original draft preparation, D.H.; writing—review and editing, D.H., A.R., L.K. and J.C.; funding acquisition, D.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the U.S. Department of Agriculture (USDA) CBG project TENX12899 (2022-38821-37341) and the National Science Foundation (NSF) EiR project (2000058).

Data Availability Statement: No new data were created. Data are contained within this article.

Conflicts of Interest: The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

1. Foley, J.; Ramankutty, N.; Brauman, K.; Cassidy, E.; Gerber, J.; Johnston, M.; Mueller, N.; O’Connell, C.; Ray, D.; West, P.; et al. Solutions for a cultivated planet. *Nature* **2011**, *478*, 337–342. [[CrossRef](#)]
2. Tian, H.; Lu, C.; Ciais, P.; Michalak, A.; Canadell, J.; Saikawa, E.; Huntzinger, D.; Gurney, K.; Sitch, S.; Zhang, B.; et al. The terrestrial biosphere as a net source of greenhouse gases to the atmosphere. *Nature* **2016**, *531*, 225–228. [[CrossRef](#)]
3. Cassman, K.; Grassini, P. A global perspective on sustainable intensification research. *Nat. Sustain.* **2020**, *3*, 262–268. [[CrossRef](#)]

4. Chen, J.; Liu, X.; Li, L.; Zheng, J.; Qu, J.; Zheng, J.; Zhang, X.; Pan, G. Consistent increase in abundance and diversity but variable change in community composition of bacteria in topsoil of rice paddy under short term biochar treatment across three sites from South China. *Appl. Soil Ecol.* **2015**, *91*, 68–79. [[CrossRef](#)]
5. Guo, T.; Bai, S.; Omidvar, N.; Wang, Y.; Chen, F.; Zhang, M. Insight into the functional mechanisms of nitrogen-cycling inhibitors in decreasing yield-scaled ammonia volatilization and nitrous oxide emission: A global meta-analysis. *Chemosphere* **2023**, *338*, 139611. [[CrossRef](#)]
6. Kroeze, C.; Mosier, A.; Bouwman, L. Closing the global N₂O budget: A retrospective analysis 1500–1994. *Glob. Biogeochem. Cycles* **1999**, *13*, 1–8. [[CrossRef](#)]
7. Tian, H.; Xu, R.; Canadell, J.; Thompson, R.; Winiwarter, W.; Suntharalingam, P.; Davidson, E.; Ciais, P.; Jackson, R.; Janssens-Maenhout, G.; et al. A comprehensive quantification of global nitrous oxide sources and sinks. *Nature* **2020**, *586*, 248–256. [[CrossRef](#)]
8. Yin, M.; Gao, X.; Kuang, W.; Tenuta, M. Soil N₂O emissions and functional genes in response to grazing grassland with livestock: A meta-analysis. *Geoderma* **2023**, *436*, 116538. [[CrossRef](#)]
9. You, L.; Ros, G.; Chen, Y.-L.; Yang, X.; Cui, Z.; Liu, X.; Jiang, R.; Zhang, F.; Vries, W. Global meta-analysis of terrestrial nitrous oxide emissions and associated functional genes under nitrogen addition. *Soil Biol. Biochem.* **2021**, *165*, 108523. [[CrossRef](#)]
10. Ravishankara, A.; Daniel, J.; Portmann, R. Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century. *Science* **2009**, *326*, 123–125. [[CrossRef](#)] [[PubMed](#)]
11. Ollivier, Q.; Maher, D.; Pitfield, C.; Macreadie, P. Winter emissions of CO₂, CH₄, and N₂O from temperate agricultural dams: Fluxes, sources, and processes. *Ecosphere* **2019**, *10*, e02914. [[CrossRef](#)]
12. Du, Y.; Guo, X.; Li, J.; Liu, Y.; Luo, J.; Liang, Y.; Li, T. Elevated carbon dioxide stimulates nitrous oxide emission in agricultural soils: A global meta-analysis. *Pedosphere* **2022**, *32*, 3–14. [[CrossRef](#)]
13. Hui, D.; Deng, Q.; Tian, H.; Luo, Y. Global climate change and greenhouse gases emissions in terrestrial ecosystems. In *Handbook of Climate Change Mitigation and Adaptation*; Lackner, M., Sajjadi, B., Chen, W.-Y., Eds.; Springer International Publishing: Cham, Switzerland, 2022; pp. 23–76. [[CrossRef](#)]
14. Zhang, L.; Jing, Y.; Chen, C.; Xiang, Y.; Rashti, M.; Li, Y.; Deng, Q.; Zhang, R. Effects of biochar application on soil nitrogen transformation, microbial functional genes, enzyme activity, and plant nitrogen uptake: A meta-analysis of field studies. *GCB Bioenergy* **2021**, *13*, 1859–1873. [[CrossRef](#)]
15. Soussana, J.; Allard, V.; Pilegaard, K.; Ambus, P.; Amman, C.; Campbell, C.; Ceschia, E.; Clifton-Brown, J.; Czobel, S.; Domingues, R.; et al. Full accounting of the greenhouse gas (CO₂, N₂O, CH₄) budget of nine European grassland sites. *Agric. Ecosyst. Environ.* **2007**, *121*, 121–134. [[CrossRef](#)]
16. Guenet, B.; Gabrielle, B.; Chenu, C.; Arrouays, D.; Balesdent, J.; Bernoux, M.; Bruni, E.; Caliman, J.; Cardinael, R.; Chen, S.; et al. Can N₂O emissions offset the benefits from soil organic carbon storage? *Glob. Chang. Biol.* **2021**, *27*, 237–256. [[CrossRef](#)] [[PubMed](#)]
17. Garvin, J.; Buick, R.; Anbar, A.; Arnold, G.; Kaufman, A. Isotopic evidence for an aerobic nitrogen cycle in the latest Archean. *Science* **2009**, *323*, 1045–1048. [[CrossRef](#)] [[PubMed](#)]
18. Iqbal, S.; Xu, J.; Allen, S.D.; Khan, S.; Nadir, S.; Arif, M.S.; Yasmeen, T. Unraveling consequences of soil micro- and nano-plastic pollution on soil-plant system: Implications for nitrogen (N) cycling and soil microbial activity. *Chemosphere* **2020**, *260*, 127578. [[CrossRef](#)] [[PubMed](#)]
19. Duan, P.; Zhang, Q.; Xi, Z.; Xiong, Z. Mechanisms of mitigating nitrous oxide emissions from vegetable soil varied with manure, biochar and nitrification inhibitors. *Agric. For. Meteorol.* **2019**, *278*, 107672. [[CrossRef](#)]
20. Zhang, W.; Wang, C.; Liu, M.; Yu, Y. Integrated reclamation of saline soil nitrogen transformation in the hyphosphere by earthworms and arbuscular mycorrhizal fungus. *Appl. Soil Ecol.* **2019**, *135*, 137–146. [[CrossRef](#)]
21. Tang, S.; Rao, Y.; Huang, S.; Xu, Y.; Zeng, K.; Liang, X.; Ling, Q.; Liu, K.; Ma, J.; Yu, F.; et al. Impact of environmental factors on the ammonia-oxidizing and denitrifying microbial community and functional genes along soil profiles from different ecologically degraded areas in the Siding mine. *J. Environ. Manag.* **2023**, *326*, 116641. [[CrossRef](#)]
22. Gineyts, R.; Niboyet, A. Nitrification, denitrification, and related functional genes under elevated CO₂: A meta-analysis in terrestrial ecosystems. *Glob. Chang. Biol.* **2023**, *29*, 1839–1853. [[CrossRef](#)]
23. Deng, Q.; Hui, D.; Wang, J.; Iwuozo, S.; Yu, C.; Jima, T.; Smart, D.; Reddy, C.; Dennis, S. Corn yield and soil nitrous oxide emission under different fertilizer and soil management: A three-year field experiment in middle Tennessee. *PLoS ONE* **2015**, *10*, e0125406. [[CrossRef](#)]
24. Hassan, M.; Amer, M.; Mahmood, A.; Awan, M.; Barbanti, L.; Seleiman, M.; Bakhsh, G.; Alkharabsheh, H.; Babur, E.; Shao, J.; et al. Management strategies to mitigate N₂O emissions in agriculture. *Life* **2022**, *12*, 439. [[CrossRef](#)] [[PubMed](#)]
25. Kaur, N.; Kieffer, C.; Ren, W.; Hui, D. How much is soil nitrous oxide emission reduced with biochar application? An evaluation of meta-analyses. *GCB Bioenergy* **2023**, *15*, 24–37. [[CrossRef](#)]
26. Borchard, N.; Schirrmann, M.; Cayuela, M.L.; Kammann, C.; Wrage-Mönnig, N.; Estavillo, J.M.; Fuertes-Mendizábal, T.; Sigua, G.; Spokas, K.; Ippolito, J.A.; et al. Biochar, soil and land-use interactions that reduce nitrate leaching and N₂O emissions: A meta-analysis. *Sci. Total Environ.* **2019**, *651*, 2354–2364. [[CrossRef](#)]
27. Ouyang, Y.; Evans, S.; Friesen, M.; Tiemann, L. Effect of nitrogen fertilization on the abundance of nitrogen cycling genes in agricultural soils: A meta-analysis of field studies. *Soil Biol. Biochem.* **2018**, *127*, 71–78. [[CrossRef](#)]

28. Dai, Z.; Yu, M.; Chen, H.; Zhao, H.; Huang, Y.; Su, W.; Xia, F.; Chang, S.; Brookes, P.; Dahlgren, R.; et al. Elevated temperature shifts soil N cycling from microbial immobilization to enhanced mineralization, nitrification and denitrification across global terrestrial ecosystems. *Glob. Chang. Biol.* **2020**, *26*, 5267–5276. [[CrossRef](#)] [[PubMed](#)]
29. Zhang, L.; Wang, X.; Wang, J.; Wan, Q.; Liao, L.; Liu, G.; Zhang, C. Grazing exclusion reduces soil N₂O emissions by regulating *nirK*– and *nosZ*– type denitrifiers in alpine meadows. *J. Soils Sediments* **2021**, *21*, 3753–3769. [[CrossRef](#)]
30. Yin, M.; Gao, X.; Kuang, W.; Zhang, Y. Meta-analysis of the effect of nitrification inhibitors on the abundance and community structure of N₂O-related functional genes in agricultural soils. *Sci. Total Environ.* **2023**, *865*, 161215. [[CrossRef](#)]
31. Hao, J.; Feng, Y.; Wang, X.; Yu, Q.; Zhang, F.; Yang, G.; Ren, G.; Han, X.; Wang, X.; Ren, C. Soil microbial nitrogen-cycling gene abundances in response to crop diversification: A meta-analysis. *Sci. Total Environ.* **2022**, *838*, 156621. [[CrossRef](#)]
32. Yang, Y.; Feng, X.; Hu, Y.; Ren, C.; Zeng, Z. Effects of legume-oat intercropping on abundance and community structure of soil N₂-fixing bacteria. *Chin. J. Appl. Ecol.* **2017**, *28*, 957–965. [[CrossRef](#)]
33. Vadakattu, G.; Roper, M.; Roget, D. Potential for non-symbiotic N₂-fixation in different agroecological zones of southern Australia. *Aust. J. Soil Res.* **2006**, *44*, 343–354. [[CrossRef](#)]
34. Collavino, M.; Tripp, H.; Frank, I.; Vidoz, M.; Calderoli, P.; Donato, M.; Zehr, J.; Aguilar, O. *nifH* pyrosequencing reveals the potential for location-specific soil chemistry to influence N₂-fixing community dynamics. *Environ. Microbiol.* **2014**, *16*, 3211–3223. [[CrossRef](#)] [[PubMed](#)]
35. Dong, J.; Zhang, J.; Liu, Y.; Jing, H. How climate and soil properties affect the abundances of nitrogen-cycling genes in nitrogen-treated ecosystems: A meta-analysis. *Plant Soil* **2022**, *477*, 389–404. [[CrossRef](#)]
36. Gul, S.; Whalen, J. Biochemical cycling of nitrogen and phosphorus in biochar-amended soils. *Soil Biol. Biochem.* **2016**, *103*, 1–15. [[CrossRef](#)]
37. Ouyang, Y.; Norton, J. Short-term nitrogen fertilization affects microbial community composition and nitrogen mineralization functions in an agricultural soil. *Appl. Environ. Microbiol.* **2020**, *86*, e02278-19. [[CrossRef](#)] [[PubMed](#)]
38. Rosenkranz, S.; Wilcke, W.; Eisenhauer, N.; Oelmann, Y. Net ammonification as influenced by plant diversity in experimental grasslands. *Soil Biol. Biochem.* **2012**, *48*, 78–87. [[CrossRef](#)]
39. Pester, M.; Maixner, F.; Berry, D.; Rattei, T.; Koch, H.; Lückner, S.; Nowka, B.; Richter, A.; Spieck, E.; Lebedeva, E.; et al. NxrB encoding the beta subunit of nitrite oxidoreductase as functional and phylogenetic marker for nitrite-oxidizing Nitrospira. *Environ. Microbiol.* **2014**, *16*, 3055–3071. [[CrossRef](#)] [[PubMed](#)]
40. Xiao, Z.; Rasmann, S.; Yue, L.; Lian, F.; Zou, H.; Wang, Z. The effect of biochar amendment on N-cycling genes in soils: A meta-analysis. *Sci. Total Environ.* **2019**, *696*, 133984. [[CrossRef](#)]
41. Martínez-Espinosa, C.; Sauvage, S.; Al Bitar, A.; Green, P.; Vörösmarty, C.; Sánchez-Pérez, J. Denitrification in wetlands: A review towards a quantification at global scale. *Sci. Total Environ.* **2021**, *754*, 142398. [[CrossRef](#)]
42. Canfield, D.; Glazer, A.; Falkowski, P. The evolution and future of Earth's nitrogen cycle. *Science* **2010**, *330*, 192–196. [[CrossRef](#)]
43. Kandler, E.; Deiglmayr, K.; Tschirko, D.; Bru, D.; Philippot, L. Abundance of *narG*, *nirS*, *nirK*, and *nosZ* genes of denitrifying bacteria during primary successions of a glacier foreland. *Appl. Environ. Microbiol.* **2006**, *72*, 5957–5962. [[CrossRef](#)] [[PubMed](#)]
44. Kuypers, M.M.M.; Marchant, H.K.; Kartal, B. The microbial nitrogen-cycling network. *Nat. Rev. Microbiol.* **2018**, *16*, 263–276. [[CrossRef](#)]
45. Henry, S.; Bru, D.; Stres, B.; Hallet, S.; Philippot, L. Quantitative detection of the *nosZ* gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, *narG*, *nirK*, and *nosZ* genes in soils. *Appl. Environ. Microbiol.* **2006**, *72*, 5181–5189. [[CrossRef](#)] [[PubMed](#)]
46. Morales, S.E.; Cosart, T.; Holben, W.E. Bacterial gene abundances as indicators of greenhouse gas emission in soils. *ISME J.* **2010**, *4*, 799–808. [[CrossRef](#)] [[PubMed](#)]
47. McCarty, G. Modes of action of nitrification inhibitors. *Biol. Fertil. Soils* **1999**, *29*, 1–9. [[CrossRef](#)]
48. Deng, Q.; Hui, D.; Wang, J.; Yu, C.; Li, C.; Reddy, C.; Dennis, S. Assessing the impacts of tillage and fertilization management on nitrous oxide emissions in a cornfield using the DNDC model. *J. Geophys. Res. Biogeosci.* **2016**, *121*, 337–349. [[CrossRef](#)]
49. Salazar, A.; Rousk, K.; Jónsdóttir, I.; Bellenger, J.-P.; Andr sson,  . Faster nitrogen cycling and more fungal and root biomass in cold ecosystems under experimental warming: A meta-analysis. *Ecology* **2020**, *101*, e02938. [[CrossRef](#)] [[PubMed](#)]
50. Xu, X.; Ran, Y.; Li, Y.; Zhang, Q.; Liu, Y.; Pan, H.; Guan, X.; Li, J.; Shi, J.; Dong, L.; et al. Warmer and drier conditions alter the nitrifier and denitrifier communities and reduce N₂O emissions in fertilized vegetable soils. *Agric. Ecosyst. Environ.* **2016**, *231*, 133–142. [[CrossRef](#)]
51. Huang, X.; Zou, Y.; Qiao, C.; Liu, Q.; Liu, J.; Kang, R.; Ren, L.; Wu, W. Effects of biological nitrification inhibitor on nitrous oxide and *nosZ*, *nirK*, *nirS* denitrifying bacteria in Paddy soils. *Sustainability* **2023**, *15*, 5348. [[CrossRef](#)]
52. Curtis, P. A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide. *Plant Cell Environ.* **1996**, *19*, 127–137. [[CrossRef](#)]
53. Luo, Y.; Hui, D.; Zhang, D. Elevated CO₂ stimulates net accumulations of carbon and nitrogen in land ecosystems: A meta-analysis. *Ecology* **2006**, *87*, 53–63. [[CrossRef](#)]
54. Fox, J. How much does the typical ecological meta-analysis overestimate the true mean effect size? *Ecol. Evol.* **2022**, *12*, e9521. [[CrossRef](#)]
55. Makowski, D.; Catarino, R.; Chen, M.; Bosco, S.; Montero-Casta o, A.; P rez-Soba, M.; Schievano, A.; Tamburini, G. Synthesising results of meta-analyses to inform policy: A comparison of fast-track methods. *Environ. Evid.* **2023**, *12*, 16. [[CrossRef](#)]

56. Hui, D.; Kaur, N.; Ray, A.; Kasrija, L.; Li, Q. Literature Review, Meta-Analysis, and Mega-Analysis in Ecological and Agricultural Sciences. *Agric. Sci.* **2023**, *14*, 474–484. [[CrossRef](#)]
57. Cui, P.; Fan, F.; Yin, C.; Song, A.; Huang, P.; Tang, Y.; Zhu, P.; Peng, C.; Li, T.; Wakelin, S.; et al. Long-term organic and inorganic fertilization alters temperature sensitivity of potential N₂O emissions and associated microbes. *Soil Biol. Biochem.* **2015**, *93*, 131–141. [[CrossRef](#)]
58. Li, L.; Zheng, Z.; Wang, W.; Biederman, J.; Xu, X.; Ran, Q.; Qian, R.; Xu, C.; Zhang, B.; Wang, F.; et al. Terrestrial N₂O emissions and related functional genes under climate change: A global meta-analysis. *Glob. Chang. Biol.* **2020**, *26*, 931–943. [[CrossRef](#)] [[PubMed](#)]
59. Li, Y.; Ma, J.; Yu, Y.; Li, Y.; Shen, X.; Huo, S.; Xia, X. Effects of multiple global change factors on soil microbial richness, diversity and functional gene abundances: A meta-analysis. *Sci. Total Environ.* **2022**, *815*, 152737. [[CrossRef](#)] [[PubMed](#)]
60. Niu, S.; Sherry, R.; Zhou, X.; Wan, S.; Luo, Y. Nitrogen regulation of the climate–carbon feedback: Evidence from a long-term global change experiment. *Ecology* **2010**, *91*, 3261–3273. [[CrossRef](#)]
61. Weedon, J.; Kowalchuk, G.; Aerts, R.; van Hal, J.; van Logtestijn, R.; Taş, N.; Röling, W.; van Bodegom, P. Summer warming accelerates sub-arctic peatland nitrogen cycling without changing enzyme pools or microbial community structure. *Glob. Chang. Biol.* **2012**, *18*, 138–150. [[CrossRef](#)]
62. Mooshammer, M.; Wanek, W.; Hämmerle, I.; Fuchslueger, L.; Hofhansl, F.; Knoltsch, A.; Schneckner, J.; Takriti, M.; Watzka, M.; Wild, B.; et al. Adjustment of microbial nitrogen use efficiency to carbon:nitrogen imbalances regulates soil nitrogen cycling. *Nat. Commun.* **2014**, *5*, 3694. [[CrossRef](#)] [[PubMed](#)]
63. Xue, K.; Xie, J.; Zhou, A.; Liu, F.; Li, D.; Wu, L.; Deng, Y.; He, Z.; Van Nostrand, J.; Luo, Y.; et al. Warming alters expressions of microbial functional genes important to ecosystem functioning. *Front. Microbiol.* **2016**, *7*, 668. [[CrossRef](#)] [[PubMed](#)]
64. Xue, K.; Yuan, M.; Shi, Z.; Qin, Y.; Deng, Y.; Cheng, L.; Wu, L.; He, Z.; Van Nostrand, J.; Bracho, R.; et al. Tundra soil carbon is vulnerable to rapid microbial decomposition under climate warming. *Nat. Clim. Chang.* **2016**, *6*, 595–600. [[CrossRef](#)]
65. Liu, Y.; Zhou, H.; Wang, J.; Liu, X.; Cheng, K.; Li, L.; Zheng, J.; Zhang, X.; Pan, G.-X. Short-term response of nitrifier communities and potential nitrification activity to elevated CO₂ and temperature interaction in a Chinese paddy field. *Appl. Soil Ecol.* **2015**, *96*, 88–98. [[CrossRef](#)]
66. Waghmode, T.; Chen, S.; Li, J.; Sun, R.; Liu, B.; Hu, C. Response of nitrifier and denitrifier abundance and microbial community structure to experimental warming in an agricultural ecosystem. *Front. Microbiol.* **2018**, *9*, 474. [[CrossRef](#)]
67. Bai, E.; Li, S.; Xu, W.; Li, W.; Dai, W.; Jiang, P. A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics. *New Phytol.* **2013**, *199*, 441–451. [[CrossRef](#)]
68. Liu, S.; Zheng, Y.; Ma, R.; Yu, K.; Han, Z.; Xiao, S.; Li, Z.; Wu, S.; Li, S.; Wang, J.; et al. Increased soil release of greenhouse gases shrinks terrestrial carbon uptake enhancement under warming. *Glob. Chang. Biol.* **2020**, *26*, 4601–4613. [[CrossRef](#)] [[PubMed](#)]
69. Carey, C.; Dove, N.; Beman, J.; Hart, S.; Aronson, E. Meta-analysis reveals ammonia-oxidizing bacteria respond more strongly to nitrogen addition than ammonia-oxidizing archaea. *Soil Biol. Biochem.* **2016**, *99*, 158–166. [[CrossRef](#)]
70. Li, L.; Yang, M.; Li, J.; Roland, B.; Du, Z.; Wu, D. Potential denitrification activity response to long-term nitrogen fertilization—A global meta-analysis. *J. Clean. Prod.* **2022**, *336*, 130451. [[CrossRef](#)]
71. Gao, Y.; Tan, Z.; Wang, H.; Zhu, Y. Nitrogen fertilization and the rhizosphere effect on nitrogen cycling: A meta-analysis. *Appl. Soil Ecol.* **2023**, *186*, 104788. [[CrossRef](#)]
72. Meng, X.; Yu, H.; Zhang, X.; Li, Y.; Zamanien, K.; Yao, H. Fertilization regimes and the nitrification process in paddy soils: Lessons for agricultural sustainability from a meta-analysis. *Appl. Soil Ecol.* **2023**, *186*, 104844. [[CrossRef](#)]
73. Zhang, H.-M.; Liang, Z.; Li, Y.; Chen, Z.-X.; Zhang, J.-B.; Cai, Z.-C.; Elsgaard, L.; Cheng, Y.; van Groenigen, K.; Abalos, D. Liming modifies greenhouse gas fluxes from soils: A meta-analysis of biological drivers. *Agric. Ecosyst. Environ.* **2022**, *340*, 108182. [[CrossRef](#)]
74. Su, P.; Gao, C.; Zhang, X.; Zhang, D.; Liu, X.; Xiang, T.; Luo, Y.; Chu, K.; Zhang, G.; Bu, N.; et al. Microplastics stimulated nitrous oxide emissions primarily through denitrification: A meta-analysis. *J. Hazard. Mater.* **2023**, *445*, 130500. [[CrossRef](#)]
75. Xue, R.; Wang, C.; Liu, X.; Liu, M. Earthworm regulation of nitrogen pools and dynamics and marker genes of nitrogen cycling: A meta-analysis. *Pedosphere* **2022**, *32*, 131–139. [[CrossRef](#)]
76. Jansson, J.; Hofmockel, K. Soil microbiomes and climate change. *Nat. Rev. Microbiol.* **2020**, *18*, 35–46. [[CrossRef](#)] [[PubMed](#)]
77. Shen, L.; Yang, Y.; Liu, J.; Hu, Z.; Liu, X.; Tian, M.; Yang, W.; Jin, J.; Wang, H.; Wang, Y.; et al. Different responses of ammonia-oxidizing archaea and bacteria in paddy soils to elevated CO₂ concentration. *Environ. Pollut.* **2021**, *286*, 117558. [[CrossRef](#)] [[PubMed](#)]
78. Simonin, M.; Le Roux, X.; Poly, F.; Lerondelle, C.; Hungate, B.; Nunan, N.; Niboyet, A. Coupling between and among ammonia oxidizers and nitrite oxidizers in grassland mesocosms submitted to elevated CO₂ and nitrogen supply. *Microb. Ecol.* **2015**, *70*, 809–818. [[CrossRef](#)] [[PubMed](#)]
79. Yang, S.; Zheng, Q.; Yuan, M.; Shi, Z.; Chiariello, N.; Docherty, K.; Dong, S.; Field, C.; Gu, Y.; Gutknecht, J.; et al. Long-term elevated CO₂ shifts composition of soil microbial communities in a Californian annual grassland, reducing growth and N utilization potentials. *Sci. Total Environ.* **2019**, *652*, 1474–1481. [[CrossRef](#)] [[PubMed](#)]
80. Barnard, R.; Leadley, P.; Hungate, B. Global change, nitrification, and denitrification: A review. *Glob. Biogeochem. Cycles* **2005**, *19*, GB1007. [[CrossRef](#)]

81. Šťovíček, A.; Kim, M.; Or, D.; Gillor, O. Microbial community response to hydration-desiccation cycles in desert soil. *Sci. Rep.* **2017**, *7*, 45735. [[CrossRef](#)] [[PubMed](#)]
82. Homyak, P.; Allison, S.; Huxman, T.; Goulden, M.; Treseder, K. Effects of drought manipulation on soil nitrogen cycling: A meta-analysis. *J. Geophys. Res. Biogeosci.* **2017**, *122*, 3260–3272. [[CrossRef](#)]
83. Yan, G.; Mu, C.; Xing, Y.; Wang, Q. Responses and mechanisms of soil greenhouse gas fluxes to changes in precipitation intensity and duration: A meta-analysis for a global perspective. *Can. J. Soil Sci.* **2018**, *98*, 591–603. [[CrossRef](#)]
84. Vitousek, P.; Aber, J.; Howarth, R.; Likens, G.; Matson, P.; Schindler, D.; Schlesinger, W.; Tilman, D. Human alteration of the global nitrogen cycle: Sources and consequences. *Ecol. Appl.* **1997**, *7*, 737–750. [[CrossRef](#)]
85. Lebauer, D.; Treseder, K. Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology* **2008**, *89*, 371–379. [[CrossRef](#)] [[PubMed](#)]
86. Domeignoz-Horta, L.; Philippot, L.; Peyrard, C.; Bru, D.; Breuil, M.; Bizouard, F.; Justes, E.; Mary, B.; Léonard, J.; Spor, A. Peaks of in situ N₂O emissions are influenced by N₂O-producing and reducing microbial communities across arable soils. *Glob. Chang. Biol.* **2018**, *24*, 360–370. [[CrossRef](#)]
87. Kelly, J.; Policht, K.; Grancharova, T.; Hundal, L. Distinct responses in ammonia-oxidizing archaea and bacteria after addition of biosolids to an agricultural soil. *Appl. Environ. Microbiol.* **2011**, *77*, 6551–6558. [[CrossRef](#)] [[PubMed](#)]
88. Fierer, N.; Lauber, C.; Ramirez, K.; Zaneveld, J.; Bradford, M.; Knight, R. Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients. *ISME J.* **2012**, *6*, 1007–1017. [[CrossRef](#)]
89. Soares, J.; Cassman, N.; Kielak, A.; Pijl, A.; Carmo, J.; Lourenço, K.; Laanbroek, H.; Cantarella, H.; Kuramae, E. Nitrous oxide emission related to ammonia-oxidizing bacteria and mitigation options from N fertilization in a tropical soil. *Sci. Rep.* **2016**, *6*, 30349. [[CrossRef](#)]
90. Hallin, S.; Jones, C.; Schloter, M.; Philippot, L. Relationship between N-cycling communities and ecosystem functioning in a 50-year-old fertilization experiment. *ISME J.* **2009**, *3*, 597–605. [[CrossRef](#)]
91. Liang, Y.; Ning, D.; Lu, Z.; Zhang, N.; Hale, L.; Wu, L.; Clark, I.; McGrath, S.; Storkey, J.; Hirsch, P.; et al. Century long fertilization reduces stochasticity controlling grassland microbial community succession. *Soil Biol. Biochem.* **2020**, *151*, 108023. [[CrossRef](#)]
92. Lehmann, J.; Rillig, M.; Thies, J.; Masiello, C.; Hockaday, W.; Crowley, D. Biochar effects on soil biota—A review. *Soil Biol. Biochem.* **2011**, *43*, 1812–1836. [[CrossRef](#)]
93. Liu, Q.; Liu, B.; Zhang, Y.; Hu, T.; Lin, Z.; Liu, G.; Wang, X.; Ma, J.; Wang, H.; Jin, H.; et al. Biochar application as a tool to decrease soil nitrogen losses (NH₃ volatilization, N₂O emissions, and N leaching) from croplands: Options and mitigation strength in a global perspective. *Glob. Chang. Biol.* **2019**, *25*, 2077–2093. [[CrossRef](#)] [[PubMed](#)]
94. Hui, D. Effects of biochar application on soil properties, plant biomass production, and soil greenhouse gas emissions: A mini-review. *Agric. Sci.* **2021**, *12*, 213–236. [[CrossRef](#)]
95. Ducey, T.; Ippolito, J.A.; Cantrell, K.; Novak, J.; Lentz, R. Addition of activated switchgrass biochar to an aridic subsoil increases microbial nitrogen cycling gene abundances. *Appl. Soil Ecol.* **2013**, *65*, 65–72. [[CrossRef](#)]
96. Yu, M.; Meng, J.; Yu, L.; Su, W.; Afzal, M.; Li, Y.; Brookes, P.; Redmile-Gordon, M.; Luo, Y.; Xu, J. Changes in nitrogen related functional genes along soil pH, C and nutrient gradients in the charosphere. *Sci. Total Environ.* **2019**, *650*, 626–632. [[CrossRef](#)] [[PubMed](#)]
97. Liu, Q.; Liu, B.; Zhang, Y.; Lin, Z.; Zhu, T.; Sun, R.; Wang, X.; Ma, J.; Bei, Q.; Liu, G.; et al. Can biochar alleviate soil compaction stress on wheat growth and mitigate soil N₂O emissions? *Soil Biol. Biochem.* **2017**, *104*, 8–17. [[CrossRef](#)]
98. Edwards, J.; Pittelkow, C.; Kent, A.; Yang, W. Dynamic biochar effects on soil nitrous oxide emissions and underlying microbial processes during the maize growing season. *Soil Biol. Biochem.* **2018**, *122*, 81–90. [[CrossRef](#)]
99. Bai, S.; Reverchon, F.; Xu, C.; Xu, Z.; Blumfield, T.; Zhao, H.; Van Zwieten, L.; Wallace, H. Wood biochar increases nitrogen retention in field settings mainly through abiotic processes. *Soil Biol. Biochem.* **2015**, *90*, 232–240. [[CrossRef](#)]
100. Liang, F.; Li, G.-T.; Lin, Q.-M.; Zhao, X.-R. Crop yield and soil properties in the first 3 years after biochar application to a Calcareous soil. *J. Integr. Agric.* **2014**, *13*, 525–532. [[CrossRef](#)]
101. Van Zwieten, L.; Singh, B.; Kimber, S.; Murphy, D.; Macdonald, L.; Rust, J.; Morris, S. An incubation study investigating the mechanisms that impact N₂O flux from soil following biochar application. *Agric. Ecosyst. Environ.* **2014**, *191*, 53–62. [[CrossRef](#)]
102. Cayuela, M.; van Zwieten, L.; Singh, B.; Jeffery, S.; Roig, A.; Sánchez-Monedero, M. Biochar's role in mitigating soil nitrous oxide emissions: A review and meta-analysis. *Agric. Ecosyst. Environ.* **2014**, *191*, 5–16. [[CrossRef](#)]
103. He, Y.; Zhou, X.; Jiang, L.; Li, M.; Du, Z.; Zhou, G.; Shao, J.; Wang, X.; Xu, Z.; Bai, S.; et al. Effects of biochar application on soil greenhouse gas fluxes: A meta-analysis. *GCB Bioenergy* **2017**, *9*, 743–755. [[CrossRef](#)]
104. Nguyen, T.; Xu, C.; Tahmasbian, I.; Che, R.; Xu, Z.; Zhou, X.; Wallace, H.; Bai, S. Effects of biochar on soil available inorganic nitrogen: A review and meta-analysis. *Geoderma* **2017**, *288*, 79–96. [[CrossRef](#)]
105. Akiyama, H.; Yan, X.; Yagi, K. Evaluation of effectiveness of enhanced-efficiency fertilizers as mitigation options for N₂O and NO emissions from agricultural soils: Meta-analysis. *Glob. Chang. Biol.* **2010**, *16*, 1837–1846. [[CrossRef](#)]
106. Lei, J.; Fan, Q.; Yu, J.; Ma, Y.; Yin, J.; Liu, R. A meta-analysis to examine whether nitrification inhibitors work through selectively inhibiting ammonia-oxidizing bacteria. *Front. Microbiol.* **2022**, *13*, 962146. [[CrossRef](#)] [[PubMed](#)]
107. Rillig, M.; Lehmann, A. Microplastic in terrestrial ecosystems. *Science* **2020**, *368*, 1430–1431. [[CrossRef](#)] [[PubMed](#)]
108. Zhang, Y.; Zhang, X.; Li, X.; He, D. Interaction of microplastics and soil animals in agricultural ecosystems. *Curr. Opin. Environ. Sci. Health* **2022**, *26*, 100327. [[CrossRef](#)]

109. Zhu, F.; Yan, Y.; Doyle, E.; Zhu, C.; Jin, X.; Chen, Z.; Wang, C.; He, H.; Zhou, D.; Gu, C. Microplastics altered soil microbiome and nitrogen cycling: The role of phthalate plasticizer. *J. Hazard. Mater.* **2022**, *427*, 127944. [[CrossRef](#)]
110. Li, H.; Liu, L.; Xu, Y.; Zhang, J. Microplastic effects on soil system parameters: A meta-analysis study. *Environ. Sci. Pollut. Res.* **2022**, *29*, 11027–11038. [[CrossRef](#)]
111. Song, L.; Niu, S. Increased soil microbial AOB *amoA* and *narG* abundances sustain long-term positive responses of nitrification and denitrification to N deposition. *Soil Biol. Biochem.* **2022**, *166*, 108539. [[CrossRef](#)]
112. Gao, H.; Liu, Q.; Yan, C.; Mancl, K.; Gong, D.; He, J.; Mei, X. Macro-and/or microplastics as an emerging threat effect crop growth and soil health. *Resour. Conserv. Recycl.* **2022**, *186*, 106549. [[CrossRef](#)]
113. Yu, Y.; Li, X.; Feng, Z.; Xiao, M.; Ge, T.; Li, Y.; Yao, H. Polyethylene microplastics alter the microbial functional gene abundances and increase nitrous oxide emissions from Paddy soils. *J. Hazard. Mater.* **2022**, *432*, 128721. [[CrossRef](#)] [[PubMed](#)]
114. Seeley, M.; Song, B.; Passie, R.; Hale, R. Microplastics affect sedimentary microbial communities and nitrogen cycling. *Nat. Commun.* **2020**, *11*, 2372. [[CrossRef](#)] [[PubMed](#)]
115. Chen, X.; Yang, J.; Zhu, X.; Liang, X.; Lei, Y.; He, C. N-fixing trees in wetland restoration plantings: Effects on nitrogensupply and soil microbial communities. *Environ. Sci. Pollut. Res.* **2016**, *23*, 24749–24757. [[CrossRef](#)] [[PubMed](#)]
116. Cai, S.; Zhang, H.; Zhu, H.; Xu, S.; Zhang, D.; Lv, W. Rice rotation system affects the spatial dynamics of the diazotrophic community in Paddy soil of the Yangtze Delta, China. *Eurasian Soil Sci.* **2019**, *52*, 696–706. [[CrossRef](#)]
117. Zou, J.; Yao, Q.; Liu, J.; Li, Y.; Song, F.; Liu, X.; Wang, G. Changes of diazotrophic communities in response to cropping systems in a Mollisol of Northeast China. *PeerJ* **2020**, *8*, e9550. [[CrossRef](#)] [[PubMed](#)]
118. Barros, F.; Fracetto, F.; Lira Junior, M.; Bertini, S.; Fracetto, G. Spatial and seasonal responses of diazotrophs and ammonium-oxidizing bacteria to legume-based silvopastoral systems. *Appl. Soil Ecol.* **2021**, *158*, 103797. [[CrossRef](#)]
119. Chen, J.; Arafat, Y.; Wu, L.; Xiao, Z.; Li, Q.; Khan, M.; Khan, M.; Lin, S.; Lin, W. Shifts in soil microbial community, soil enzymes and crop yield under peanut/maize intercropping with reduced nitrogen levels. *Appl. Soil Ecol.* **2018**, *124*, 327–334. [[CrossRef](#)]
120. Reverchon, F.; Bai, S.; Liu, X.; Blumfield, T. Tree plantation systems influence nitrogen retention and the abundance of nitrogen functional genes in the Solomon Islands. *Front. Microbiol.* **2015**, *6*, 1439. [[CrossRef](#)]
121. Chu, H.; Morimoto, S.; Fujii, T.; Yagi, K.; Nishimura, S. Soil ammonia-oxidizing bacterial communities in paddy rice fields as affected by upland conversion history. *Soil Sci. Soc. Am. J.* **2009**, *73*, 2026–2031. [[CrossRef](#)]
122. Pan, H.; Xie, K.; Zhang, Q.; Jia, Z.; Xu, J.; Di, H.; Li, Y. Archaea and bacteria respectively dominate nitrification in lightly and heavily grazed soil in a grassland system. *Biol. Fertil. Soils* **2018**, *54*, 41–54. [[CrossRef](#)]
123. Le Roux, X.; Poly, F.; Currey, P.; Commeaux, C.; Hai, B.; Nicol, G.; Prosser, J.; Schloter, M.; Attard, E.; Klumpp, K. Effects of aboveground grazing on coupling among nitrifier activity, abundance and community structure. *ISME J.* **2008**, *2*, 221–232. [[CrossRef](#)]
124. Zhong, L.; Zhou, X.; Wang, Y.; Li, F.; Zhou, S.; Bai, Y.; Rui, Y. Mixed grazing and clipping is beneficial to ecosystem recovery but may increase potential N₂O emissions in a semi-arid grassland. *Soil Biol. Biochem.* **2017**, *114*, 42–51. [[CrossRef](#)]
125. Ding, K.; Zhong, L.; Xin, X.; Xu, Z.; Kang, X.; Liu, W.; Rui, Y.; Jiang, L.; Tang, L.; Wang, Y. Effect of grazing on the abundance of functional genes associated with N cycling in three types of grassland in Inner Mongolia. *J. Soils Sediments* **2015**, *15*, 683–693. [[CrossRef](#)]
126. Willems, J.; Marinissen, J.; Blair, J. Effects of earthworms on nitrogen mineralization. *Biol. Fertil. Soils* **1996**, *23*, 57–63. [[CrossRef](#)]
127. Blouin, M.; Hodson, M.; Delgado, E.; Baker, G.; Brussaard, L.; Butt, K.; Dai, J.; Dendooven, L.; Peres, G.; Tondoh, J.; et al. A review of earthworm impact on soil function and ecosystem services. *Eur. J. Soil Sci.* **2013**, *64*, 161–182. [[CrossRef](#)]
128. Hoang, D.; Pausch, J.; Razavi, B.; Kuzyakova, I.; Banfield, C.; Kuzyakov, Y. Hotspots of microbial activity induced by earthworm burrows, old root channels, and their combination in subsoil. *Biol. Fertil. Soils* **2016**, *52*, 1105–1119. [[CrossRef](#)]
129. Lubbers, I.; van Groenigen, K.; Fonte, S.; Six, J.; Brussaard, L.; Van Groenigen, J. Greenhouse gas emissions from soils increased by earthworms. *Nat. Clim. Chang.* **2013**, *3*, 187–194. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.