

Supplementary Materials

Hydro-Saline Dynamics of a Shallow Mediterranean Coastal Lagoon: Complementary Information from Short and Long Term Monitoring

Olivier Boutron ^{1,*}, Caroline Paugam ², Emilie Luna-Laurent ¹, Philippe Chauvelon ¹, Damien Sous ³, Vincent Rey ², Samuel Meulé ⁴, Yves Chérain ⁵, Anais Cheiron ⁵ and Emmanuelle Migne ⁵

¹ Tour du Valat Research Institute, Le Sambuc, 13200 Arles, France; lunalaurent@tourduvalat.org (E.L.-L.); philou13ch@hotmail.com (P.C.)

² Mediterranean Institute of Oceanography (MIO), Université de Toulon AMU, CNRS/INSU, IRD, UM 110 Toulon, France; caroline.paugam@mio.osupytheas.fr (C.P.); vincent.rey@univ-tln.fr (V.R.)

³ Université de Pau, Pays Adour-E2S UPPA, SIAME, Anglet, France; damien.sous@mio.osupytheas.fr

⁴ Université Aix Marseille, CNRS, IRD, INRA, Coll France, CEREGE, Aix-en-Provence, France; meule@cerege.fr

⁵ National Nature Reserve of Camargue, National Society for Nature Protection (SNPN), La Capeliere, 13200 Arles, France; ycherain@gmail.com (Y.C.); anais.cheiron@espaces-naturels.fr (A.C.); emmanuelle.migne@espaces-naturels.fr (E.M.)

* Correspondence: boutron@tourduvalat.org; Tel.: +33-4-90-97-63-73




Figure S1. Percentage of volume contained in the i) Vaccarès, ii) Impériaux and iii) Lion/Dame lagoons, in comparison with the total volume contained in the Vaccarès Lagoon System.




Figure S2. Percentage of mass of dissolved salt contained in the i) Vaccarès, ii) Impériaux and iii) Lion/Dame lagoons, in comparison with the total mass of dissolved salt contained in the Vaccarès Lagoon System.