

## Supplementary Materials


### Can zeolite-supporting acridines boost their anticancer performance?

Maja Ranković<sup>a</sup>, Anka Jevremović<sup>a</sup>, Aleksandra Janošević Ležaić<sup>b</sup>, Aleksandar Arsenijević<sup>c</sup>, Jelena Rugar<sup>b</sup>, Vladimir Dobričić<sup>b</sup>, Bojana Nedić Vasiljević<sup>a</sup>, Nemanja Gavrilov<sup>a</sup>, Danica Bajuk-Bogdanović<sup>a</sup>, Maja Milojević-Rakić<sup>a</sup>

<sup>a</sup>University of Belgrade–Faculty of Physical Chemistry, 11000 Belgrade, Serbia.

<sup>b</sup>University of Belgrade–Faculty of Pharmacy, Department of Physical Chemistry and Instrumental Methods, 11221 Belgrade, Serbia

<sup>c</sup>University of Kragujevac–Faculty of Medical Sciences, Department of Pharmacy and Center for molecular medicine and stem cells research, 34000 Kragujevac, Serbia



**Figure S1.** MolView presentation (H – white, C – grey, Cl – green, N – blue) and assessment of acridines dimensions.

**Table S1.** Solvation enthalpy for tested acridines calculated from heat of formation with and without solvation sphere.

	Heat of formation/kcal per mol		
	Acr	9ClAcr	9NAcr
No solvation	71.7	62.5	69.2
Solvation	63.0	56.2	53.4
Dielectric part / eV	-0.44	-0.34	-1.03