

Supporting information

Dual pH- and GSH-Responsive Degradable PEGylated Graphene Quantum Dot-Based Nanoparticles for Enhanced HER2-Positive Breast Cancer Therapy

Na Re Ko ^{1,2,†}, Se Young Van ^{1,2,†}, Sung Hwa Hong ³, Seog-Young Kim ⁴, Miran Kim ², Jae Seo Lee ⁵, Sang Ju Lee ², Yong-kyu Lee ⁶, Il Keun Kwon ⁵ and Seung Jun Oh ^{2,*}

¹ Biomedical Research Center, Asan Institute for Life Sciences, 88 Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Korea; nare.ko@amc.seoul.kr (N.R.K.); sey.van88@gmail.com (S.Y.V.)

² Department of Nuclear Medicine, Asan Medical Center, University of Ulsan College of Medicine, 88 Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Korea; ranaranmi@naver.com (M.K.); atlas425@amc.seoul.kr (S.J.L.)

³ Department of Chemical Engineering and Applied Chemistry, University of Toronto, 200 College Street, Toronto, ON M5S 3E5, Canada; sunghwa.hong@mail.utoronto.ca (S.H.H.)

⁴ Department of Convergence Medicine, University of Ulsan College of Medicine, 88 Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Korea; ksy0222@gmail.com (S.-Y.K.)

⁵ Department of Dental Materials, School of Dentistry, Kyung Hee University, 26 Kyungheedaero, Dongdaemun-gu, Seoul 02447, Korea; leejs8433214@naver.com (J.S.L.); kwoni@khu.ac.kr (I.K.K.)

⁶ Department of Chemical & Biological Engineering, Korea National University of Transportation, 50 Daehak-ro, Chungcheongbuk-do, Cheongju 27469, Korea; leeyk@ut.ac.kr (Y.-k.L.)

* Correspondence: sjoh@amc.seoul.kr; Tel.: +82-2-3010-4595

† These authors contributed equally to this work.




Figure S1. (a) expression of HER2 in SK-BR-3 and MDA-BM-231 breast cancer cell lines. β -actin served as a loading control. (b) Quantification of HER2 protein expression is given as intensity of β -actin-standardized HER2 levels in the cells.