



SUPPORTING INFORMATIONS

Pb-Free Cs₃Bi₂I₉ Perovskite as a Visible-Light-Active Photocatalyst for Organic Pollutant Degradation

Bianca-Maria Bresolin ^{1,*}, Carsten Günnemann ², Detlef W. Bahnemann ^{2,3,4} and Mika Sillanpää ^{5,6,7}

- ¹ Department of Separation Science, School of Engineering Science, Lappeenranta University of Technology, Sammonkatu 12, 50130 Mikkeli, Finland
 - ² Institute of Technical Chemistry, Leibniz University Hannover, Callinstraße 3, D-30167 Hannover, Germany; guennemann@iftc.uni-hannover.de (C.G.); Bahnemann@iftc.uni-hannover.de (D.W.B.)
 - ³ Laboratory of Nano- and Quantum-Engineering (LNQE), Gottfried Wilhelm Leibniz University Hannover, Schneiderberg 39, D-30167 Hannover, Germany
 - ⁴ Laboratory “Photoactive Nanocomposite Materials”, Saint-Petersburg State University, Ulyanovskaya str. 1, Peterhof, 198504 Saint-Petersburg, Russia
 - ⁵ Institute of Research and Development, Duy Tan University, De Nang 550000, Vietnam; mikaesillanpaa@gmail.com
 - ⁶ Faculty of Environmental and Chemical Engineering, Duy Tan University, De Nang 550000, Vietnam
 - ⁷ School of Civil Engineering and Surveying, Faculty of Health, Engineering and Science, University of Southern Queensland, West Street, Toowoomba 4350, QLD, Australia
- * Correspondence: biancabresolin@yahoo.it or Bianca.Maria.Bresolin@lut.fi




Figure S1. Irradiance spectrum of a Halogen lamp (3200K) compared to the solar irradiance on a surface per unit wavelength with emphasis on visible light range.




Figure S2. XRD patterns of CsI and BiI₃.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).