

Supplementary Materials

A High-Throughput NanoBiT-Based Serological Assay Detects SARS-CoV-2 Seroconversion

Taha Azad ^{1,2,†}, Reza Rezaei ^{1,2,†}, Ragnath Singaravelu ^{1,2,†}, Taylor R. Jamieson ^{1,2}, Mathieu J. F. Crupi ^{1,2}, Abera Surendran ^{1,2}, Joanna Poutou ^{1,2}, Parisa Taklifi ³, Juthaporn Cowan ⁴, D. William Cameron ⁴ and Carolina S. Ilkow ^{1,2,*}

¹ Ottawa Hospital Research Institute, Ottawa, ON K1H 8L6, Canada

² Department of Biochemistry, Microbiology and Immunology, University of Ottawa, Ottawa, ON K1H 8M5, Canada

³ Department of Biotechnology, University of Tehran, Tehran, Iran

⁴ Department of Medicine, Division of Infectious Disease, University of Ottawa at The Ottawa Hospital/Research Institute, Ottawa, ON, Canada

* Correspondence: cilkow@ohri.ca; Tel.: +1-613-737-8899 (ext. 75208)

† These authors contributed equally




Figure S1. High signal intensity and detection of different amount of the SARS-CoV-2 neutralizing antibodies compared to low signal for the control antibody.