



Supplementary Materials

Photosensitized Thermoplastic Nano-Photocatalysts Active in the Visible Light Range for Potential Applications Inside Extraterrestrial Facilities

Lidia Mezzina ¹, Angelo Nicosia ¹, Fabiana Vento ¹, Guido De Guidi ¹ and Placido Giuseppe Mineo ^{1,2,3,*}

¹ Department of Chemical Sciences and INSTM UdR of Catania, University of Catania, V.le A. Doria 6, I-95125 Catania, Italy; lidia.mezzina@phd.unict.it (L.M.); angelo.nicosia@unict.it (A.N.); fabiana.vento@phd.unict.it (F.V.); guido.deguidi@unict.it (G.D.G.)

² Institute for Chemical and Physical Processes, National Research Council (IPCF-CNR), Viale F. Stagno d'Alcontres 37, I-98158 Messina, Italy

³ Institute of Polymers, Composites and Biomaterials, National Research Council (IPCB-CNR), Via P. Gaifami 18, I-95126 Catania, Italy

* Correspondence: placido.mineo@unict.it




Figure S1. UV-Vis spectra of Acetaminophen (red line), Paraquat (green line), Rhodamine B (magenta line) and Methylene Blue (blue line).




Figure S2. ^1H - ^1H COSY spectrum of the MAP macromonomer (range, 7–9 ppm).