



Supplementary Materials

Facile and Green Synthesis of Highly Fluorescent Carbon Quantum Dots from Water Hyacinth for the Detection of Ferric Iron and Cellular Imaging

Pei Zhao ¹, Qin Zhang ^{1,2,*}, Juanjuan Cao ¹, Cheng Qian ¹, Jing Ye ¹, Siyuan Xu ¹, Yonggui Zhang ¹ and Yanbin Li ^{1,2,*}

¹ College of Biological and Food Engineering, Anhui Polytechnic University, Wuhu 241000, China; lvxmzp@163.com (P.Z.); cjj18890973009@163.com (J.C.); qiancheng9981@163.com (C.Q.); a2235672022@163.com (J.Y.); shanyazi0605@163.com (S.X.); xjagcdzyg@163.com (Y.Z.)

² Anhui Engineering Laboratory for Industrial Microbiology Molecular Breeding, College Biological and Food Engineering, Anhui Polytechnic University, Wuhu 241000, China

* Correspondence: jhtabszq@163.com (Q.Z.), lyb@ahpu.edu.cn (Y.L.)

Table S1. Wh-CQDs and Quinine Sulfate Concentration for Measuring QY.

Number	Concentration of Quinine Sulfate ($\mu\text{g/mL}$)	Absorbance	Concentration of Wh-CQDs ($\mu\text{g/mL}$)	Absorbance
1	1	0.016	3.17	0.036
2	2	0.032	6.36	0.048
3	3	0.043	9.22	0.062
4	4	0.057	12.41	0.079
5	5	0.074	15.53	0.092




Figure S1. Selectivity of the wh-CQDs toward different metal ions.