

Thioredoxin Reductase Activity Predicts Gold Nanoparticle Radiosensitization Effect

Sébastien Penninckx ¹, Anne-Catherine Heuskin ¹, Carine Michiels ^{2,*} and Stéphane Lucas ¹

¹ Research center for the Physics of Matter and Radiation (PMR-LARN), Namur Research Institute For Life Sciences (NARILIS), University of Namur, Rue de Bruxelles 61, B-5000 Namur, Belgium; sebastien.penninckx@unamur.be (S.P.); anne-catherine.heuskin@unamur.be (A.-C.H.); stephane.lucas@unamur.be (S.L.)

² Unité de Recherche en Biologie Cellulaire (URBC), Namur Research Institute For Life Sciences (NARILIS), University of Namur, Rue de Bruxelles 61, B-5000 Namur, Belgium

* Correspondence: carine.michiels@unamur.be; Tel.: +32-81-724131; Fax +32-81-724135

Supplementary information




Figure S1. Toxicity of GNPs across the cell lines. All cells were incubated 24 h with 50 $\mu\text{g}/\text{mL}$ GNPs prior the clonogenic assay. Results were reported as mean plating efficiency \pm SEM. Results were statistically analyzed using a one-way ANOVA evidencing no significant difference (Tukey test).




Figure S2. Box plot of TXRND1 mRNA expression across the two expression groups for (A) brain (GSE42669), (B) breast (TCGA-BRCA), (C) head & neck (TCGA-HNSC), (D) lung (GSE 30219) and (E) pancreas (PACA-AU) cancer datasets. P-values were calculated using t-test.