

Supplementary

Yeast-Based Fluorescent Sensors for the Simultaneous Detection of Estrogenic and Androgenic Compounds, Coupled with High-Performance Thin Layer Chromatography

Liat Moscovici ¹, Carolin Riegraf ^{2,3}, Nidaa Abu-Rmailah ¹, Hadas Atias ¹, Dror Shakibai ¹, Sebastian Buchinger ², Georg Reifferscheid ² and Shimshon Belkin ^{1,*}

¹ Department of Plant and Environmental Sciences, Institute of Life Sciences, Hebrew University of Jerusalem, Jerusalem 91904, Israel; liatmosc@savion.huji.ac.il (L.M.); nidaan8@gmail.com (N.A.-R.); hadas.atias@mail.huji.ac.il (H.A.); drors1987@gmail.com (D.S.)

² Department Biochemistry, Ecotoxicology, Federal Institute of Hydrology (BfG), Am Mainzer Tor 1, 56068 Koblenz, Germany; carolin.riegraf@gmail.com (C.R.); Buchinger@bafg.de (S.B.); reifferscheid@bafg.de (G.R.)

³ RWTH Aachen University, Department of Ecosystem Analysis, Worringerweg 1, D-52074 Aachen, Germany

* Correspondence: shimshon.belkin@mail.huji.ac.il; Tel.: +972-2-6584192

Table S1. Oligonucleotide primer sequences used for the construction of the plasmids in this study.

	Primer	Primer Sequence 5' to 3'
1.	ADH1_SpeI_EG FP_For	GGGGGTCTAGTAAAGAATTAATTA CATGTCAAAA
2.	ADH1_Kpn1_E GFP_Rev	GCTCGGTACCAATCTAGGCGCGCCTTA CTTGATAATTCA
3.	ADH1_SpeI_BF P_For	TTTTTCCCGGACTAGTAAAGAAATG AGTGAGTTGATTAAGGA
4.	ADH1_EcoRI_B FP_Rev	TTTTTGAATTTCGAGCTCGGTACCAATC TATTAGTTTAGTTTGTGCCCA
5.	ADH1_SpeI_Ru by_For	GGTGGTGAATTCTTACTTATACAATTC ATCCA
6.	ADH1_EcoRI_R uby_Rev	GGTGGTGAATTCTTACTTATA CAATTCATCCA